to assess the true extent of the vascular lesions from many of these case reports as post-mortem examination was frequently incomplete.

More recently, Heptinstall *et al.* (1954) have reported 14 cases of giant-cell arteritis, three of which came to necropsy. Vascular changes were widespread in the three fatal cases, and included histologically confirmed lesions in the right femoral artery and left popliteal artery in one case, and in the right popliteal artery in a second case. Pains in the limbs and joints were noted in 5 of the 14 cases—the authors suggested the possibility that the limb pains may be due to specific changes in the arteries—but peripheral gangrene was not encountered. Reference, however, is made to a case reported by Frangenheim (1951) in which a 72-year-old man had suddenly become blind and in which amputation of a leg for gangrene revealed the changes of giant-cell arteritis in the femoral artery. Reference is also made to a case described by Atlas (1943) in which a 68-year-old woman had a leg amputated for gangrene. The case was recorded as one of Buerger's disease, but we agree with Heptinstall *et al.* (1954) that the histological findings are consistent with giant-cell arteritis. In this case the patient died shortly after operation, but whether a necropsy was performed is not recorded.

It is thus apparent that giant-cell arteritis can involve the arteries of the lower limbs and in rare instances be complicated by gangrene. Giant-cell arteritis is a disease of the middle-aged and elderly, and it is therefore very probable that affected limb arteries will also show some degree of incidental atherosclerosis; but unless the latter is extensive it is reasonable to suggest that ischaemic symptoms which may develop are due mainly to the obliterative arteritis which is a well-recognized feature of giant-cell arteritis.

Giant-cell arteritis should therefore be considered in the differential diagnosis of any unexplained case of peripheral vascular disease occurring in the middle-aged or elderly, and if such a diagnosis is substantiated corticotrophin or cortisone therapy should be given before gangrene supervenes and surgery becomes imperative.

Summary

A case of giant-cell arteritis apparently localized to the arteries of the lower limbs and producing gangrene is reported. Both legs had to be amputated, and histological examination showed the typical features of giant-cell arteritis in the tibial and peroneal arteries. One limb also showed involvement of the pedal arteries, and essentially similar changes were noted in some of the small arteries supplying voluntary muscles.

A review of the literature indicates that peripheral gangrene is a rare complication of giant-cell arteritis, but the real incidence of involvement of the leg arteries in this disease is not known. The apparent rarity of this condition may account for its being overlooked in other cases, and it is suggested that giant-cell arteritis should be considered in the differential diagnosis of any unexplained case of peripheral vascular disease occurring in the middle-aged or elderly.

We are indebted to Mr. E. G. Tuckwell for permission to publish this case and to Professor J. W. S. Blacklock for his help and advice. We thank Mr. K. W. Iles for the photomicrographs and Miss Audrey Dixon for the cutting and staining of many sections.

REFERENCES

- Atlas, L. N. (1943). *Amer. Heart J.*, 26, 120.
 Cardell, B. S., and Hanley, T. (1951). *J. Path. Bact.*, 63, 587.
 Frangenheim, H. (1951). *Zbl. allg. Path. path. Anat.*, 88, 81.
 Harrison, C. V. (1947-8). *J. clin. Path.*, 1, 197.
 Heptinstall, R. H., Porter, K. A., and Barkley, H. (1954). *J. Path. Bact.*, 67, 507.
 Horton, B. T., Magath, T. B., and Brown, G. E. (1934). *Arch. intern. Med.*, 53, 400.
 Hutchinson, J. (1890). *Arch. Surg. (Lond.)*, 1, 323.
 Robertson, K. (1947). *British Medical Journal*, 2, 168.

NASAL GRANULOMA AND PERIARTERITIS NODOSA

REPORT OF A CASE

BY

PAUL F. MILNER, M.B., B.Ch.

Captain, R.A.M.C.; formerly Resident in Pathology,
Northampton General Hospital

[WITH SPECIAL PLATE]

Since Kussmaul and Maier (1866) gave the name "periarteritis nodosa" to a disease which they were the first to describe, many cases differing in certain clinical and pathological details have been reported. In 1903 Veszprémi and Jancsó made the diagnosis from histological material in a case presenting no gross evidence of the disease; and Mönckeberg (1905) for the first time described lesions of periarteritis in the lungs. The second case with pulmonary-artery involvement, reported by Ophüls (1923), had many distinguishing features, among them granulomatous lesions in serous membranes. Gradually it came to be appreciated that the microscopic form of the disease, affecting small arteries and veins, was the commoner and could be distinguished from the classical disease (Zeek *et al.*, 1948; Zeek, 1952).

Cases of periarteritis nodosa with granulomatous lesions in the upper respiratory tract are uncommon. Godman and Churg (1954), who reported seven cases of their own, were able to find 22 cases in the world literature. Of these, one reported by Howells and Friedmann (1950) and two reported by Stratton *et al.* (1953) are the only British cases in their list. Klinger (1931) was the first to describe a syndrome of severe destructive sinusitis, nephritis, and uraemia. Wegener (1936, 1939), describing three cases of ulcerative granulomata in the upper respiratory tract with periarteritis nodosa, wrote: "The point of origin of the disease was the inner nose, where a putrid, necrotic inflammation was found with granulation tissue and nodular granulomata. This evidently led to sensitization of the body as a whole, resulting in allergic, hyperergic reactions, especially in the vascular system." Stratton *et al.* (1953) stressed that "the essence of this syndrome is that the upper respiratory tract lesions are the first and, at the onset, the dominating feature of the condition." Pulmonary lesions, both vascular and granulomatous, are almost a constant feature of this syndrome; in only one case of the series reviewed by Godman and Churg were they absent. In this case, reported by Lindsay *et al.* (1944), there was a progressive destructive nasal lesion for a year before acute fatal periarteritis nodosa developed.

The case now reported appears unique in two respects: (1) the acute nasal and gingival giant-cell granulomatous lesions regressed and healed before the fatal illness; and (2) no vascular or granulomatous lesions could be found in the lungs at necropsy.

Case Report

A 41-year-old housewife had an upper right premolar extracted in May, 1954. The socket healed well till about June 17, when a bright-red swelling appeared on the gum, which became painful. Two lower left molars were extracted at this time, the sockets failing to heal and becoming swollen and painful. The face then began to swell and she had headaches and nasal discharge. She felt weak

and lethargic, but had not lost weight and had no cough or other symptoms. Until her present illness her general health had always been good.

On admission to an E.N.T. ward on June 26 she had a sessile, bluish-grey, nodular soft swelling of the gum behind the upper right canine. A similar swelling was seen in the $\overline{123}$ region. There was severe oral sepsis. The nose and cheeks were swollen and there was conjunctival and pericorneal injection of the left eye. The right nostril was obstructed and there was a yellow serous discharge. The submandibular lymph nodes were enlarged, firm, and tender, the right one feeling about the size of a golf ball. The upper nodes of the deep cervical chain on the right were slightly enlarged, but there was no other lymphadenopathy. There were no other clinical findings at this time. She had been receiving 1,000,000 units of penicillin daily for ten days without benefit and was now running a low-grade fever. X-ray examination revealed both antra clear and no abnormality in upper or lower jaws.

Progress

A piece of gum lesion was removed for histological examination. The report was: "A granuloma with pleomorphic cytology including multinucleate giant cells of the Langhans type; hyperplasia of the covering squamous epithelium with micro-abscesses at the apices of some of the papillae. The appearances are not unlike those seen in hypertrophic gingivitis of pregnancy." Sections were stained by Ziehl-Neelsen and periodic-acid-Schiff techniques without revealing anything further (Special Plate, Fig. 1).

By July 6 the patient was worse. The submandibular nodes were larger, the nose was painfully swollen, and completely blocked by congested mucosa and a shaggy ulcerating granulomatous lesion involving the inferior turbinates. Antral radiographs showed a triangular density in the lower lateral angle of the right antrum. A further biopsy was performed; the histology was the same as before. Aureomycin, 250 mg. four-hourly, was given.

From the beginning of July the urine contained a trace of albumin and the deposit occasional red cells. On July 14 it reduced Benedict's solution; the blood sugar was 340 mg. per 100 ml. A glucose tolerance test showed a diabetic curve: fasting, 220 mg. per 100 ml.; at 30 minutes, 240 mg.; at one hour, 250 mg.; at 90 minutes, 300 mg.; and at two hours, 350 mg. At this time blood examination showed a rising leucocytosis and falling haemoglobin level and red cell count. Repeated blood culture failed to grow any organism; the Wassermann and Kahn reactions were negative; a Mantoux test was negative with 1 in 100 old tuberculin; and the serum was negative for antibodies to enteric organisms and *Brucella abortus*. A vesicular eruption broke out in a small area over the right deltoid, and a biopsy of this was taken. Sections showed subcutaneous haemorrhage beginning to organize, but no evidence of vascular disease. Again Ziehl-Neelsen and periodic-acid-Schiff staining techniques gave no further information (Plate, Fig. 2).

By July 20 the granulomatous lesions were diminishing and the secondary infection was subsiding. On July 22 the serum proteins were 6.15 g. per 100 ml., with albumin 1.73 g. and globulin 4.42 g. per 100 ml. Serum bilirubin was 0.9 mg. per 100 ml. By July 26 the mouth and nose lesions had almost completely resolved, the nose was clear, the facial swelling had subsided, and the conjunctivitis was no longer present. X-ray appearances of skull, chest, and pelvis were normal, and the right antrum was now clear. The patient was still pyrexial and complained of great lassitude and weakness; she was depressed and wept frequently.

On August 5 she was transferred to a medical ward. The mouth was now clean, all signs of the granuloma having disappeared. She was obviously very ill, with fever rising to 101° F. (38.3° C.) at times, and her skin was pale and sallow without icterus. She complained of weakness, of headache, and of discomfort on swallowing. The blood pressure was 190/120 mm. Hg and there was slight oedema

of the ankles. All tendon reflexes were greatly diminished. The blood urea was found to be 200 mg. per 100 ml., but the glucose tolerance curve had altered: fasting, 160 mg. per 100 ml.; at 30 minutes, 180 mg.; at one hour, 190 mg.; at 90 minutes, 210 mg.; and at two hours, 190 mg. Serum protein values were the same as before. On August 13 she complained of pain and tenderness in the left loin and became drowsy and disoriented. Blood urea was 300 mg. per 100 ml. Next day she became comatose and died, eight weeks and three days from the onset of the gum lesions.

Blood Picture

Haemoglobin values fell from 15 g. per 100 ml. on admission to 9 g. the day before death. Leucocytosis reached 24,000 per c.mm. during the sixth week; eosinophils never exceeded 5% of the total count. Bone marrow was studied on two occasions without any pathological findings.

Post-mortem Examination

The gross findings were as follows. Small right hydrothorax, no pleurisy, moderate oedema of the lungs. Normal-sized heart with small grey nodules in the myocardium. Moderately enlarged, grossly fatty liver. Moderately enlarged spleen with gross perisplenitis, and cut surface showing many grey nodules of varying sizes, some with central thrombosed vessels, standing out against a dull-red hyaline background. Small fibrosed nodules in the pancreas with haemorrhages. Retroperitoneal haemorrhage around the left kidney. Both kidneys swollen, with tense capsules which on stripping left a glistening mottled surface with petechial haemorrhages. The cut surface showed complete destruction of architecture, small thrombosed vessels surrounded by grey nodules standing out against a glistening background. No gum or nasal lesions could be found.

Histology.—In the lungs there was oedema and mild bronchopneumonic changes. No active arteritis was seen, and in sections stained with orcein-methylene-blue for elastic tissue no healed arteritis, as indicated by segmental rupture of elastic fibres, was demonstrable. In the heart a large branch of a coronary artery showed acute arteritis and was partially blocked by thrombus. There were multiple foci of muscle necrosis in varying stages of replacement fibrosis, some of which contained many inflammatory cells and new capillaries. The liver showed extensive peripheral fatty change with pleomorphic cellular infiltration in the portal tracts without any marked changes in the walls of the branches of the hepatic artery. The spleen was extensively necrosed with little unaffected tissue. There were gross necrotizing changes in the vessel walls and an extensive inflammatory exudate. In the pancreas one area showed acute arteritis with necrosis and another a hyalinized nodule. Post-mortem change made interpretation of the lesions difficult. The salivary gland was traversed by intersecting bands of fibrosis with infiltration, mainly lymphocytic and plasmacell, of remaining parenchyma.

In the kidney there were areas of acute necrosis, and elsewhere oedema and pleomorphic cellular infiltration, mainly polymorphs, eosinophils, and histiocytes, which varied in intensity from place to place and was associated with foci of acute arteritis in medium- and large-sized vessels. There was necrotizing glomerulitis with progressive shrinkage of glomerular tufts and periglomerular fibrosis. Necrosis of tubular epithelium appeared mainly in the proximal convoluted tubules, but owing to the extent and severity of the renal lesions the region of maximum severity was difficult to determine (Plate, Fig. 3).

Discussion

This case was essentially one of periarteritis nodosa with predominance of upper respiratory lesions at onset. These lesions were not invasive or destructive, and resolved before the periarteritis terminated in uraemia. They were,

however, of a granulomatous nature with numerous giant-cells—a picture that has been associated with “malignant granuloma” and “Wegener’s granulomatosis.”

Godman and Churg (1954) are of the opinion that “Wegener’s granulomatosis” and “granuloma gangraenescens” (malignant granuloma) are closely related conditions. These authors were able to study one case of the latter disease of slow progressive course in which at necropsy recent arteritic lesions were found. They also note with interest that in the case of “Wegener’s granulomatosis” of longest duration—48 months—only minimal vascular lesions were present *post mortem*. One of their patients is receiving maintenance doses of cortisone and antibiotics nearly three years since he presented with destructive sinusitis and pulmonary granulomata and nephritis.

Summary

The case history of a woman of 41 years who presented with granulomatous rhinitis and gingivitis and died in uraemia after a seven-weeks illness is reported. Necropsy and histological examinations revealed necrotizing glomerulonephritis and lesions of “periarteritis” in many organs, but absent from the lungs.

The relationship between “periarteritis nodosa” and periarteritis with granulomatous lesions is briefly discussed. The general occurrence of pulmonary lesions in the latter syndrome is noted.

Attention is drawn to a recent survey of “Wegener’s granulomatosis” and the possible relationship of this condition to “malignant granuloma.”

I am indebted to Dr. D. H. G. MacQuaide, Mr. W. C. Gledhill, and Dr. R. H. Heggie for permission to report this case; to Dr. G. Hollingsworth for radiological reports; to Dr. W. E. Bryan for biopsy reports and post-mortem histology, and for assistance and advice; and to Dr. R. B. Coles for his interest and encouragement in the preparation of this report.

REFERENCES

- Godman, G. C., and Churg, J. (1954). *Arch. Path.*, **58**, 533.
 Howells, G. H., and Friedmann, I. (1950). *J. clin. Path.*, **3**, 220.
 Klinger, H. (1931). *Frankfurt. Z. Path.*, **42**, 455.
 Kussmaul, A., and Maier, R. (1866). *Dtsch. Arch. klin. Med.*, **1**, 484.
 Lindsay, S., Aggeler, P. M., and Lucia, S. P. (1944). *Amer. J. Path.*, **20**, 1057.
 Mönckeberg, J. C. (1905). *Beitr. path. Anat.*, **38**, 101.
 Ophüls, W. (1923). *Arch. intern. Med.*, **32**, 870.
 Stratton, H. J. M., Price, T. M. L., and Skelton, M. O. (1953). *British Medical Journal*, **1**, 127.
 Veszprémi, D., and Jancsó, M. (1963). *Beitr. path. Anat.*, **34**, 1.
 Wegener, F. (1936). *Verh. dtsch. path. Ges.*, **29**, 202.
 — (1939). *Beitr. path. Anat.*, **102**, 36.
 Zeek, P. M., Smith, C. M., and Weeter, J. C. (1948). *Amer. J. Path.*, **24**, 889.
 — (1952). *Amer. J. clin. Path.*, **22**, 777.

The fortieth annual report of St. Dunstan’s, *Forty Years On*, gives an account of the work of St. Dunstan’s for the war-blinded since its inception. Some 5,000 Servicemen and a few Home Guards and Civil Defence workers have come to St. Dunstan’s for training and rehabilitation. To-day a total of 2,600 St. Dunstaners are living—1,450 from the first world war and 1,150 from the second. 58 new cases, including 27 from the second world war, recent casualties from Malaya, Korea, and East Africa, and men from 1914–1918 who have gradually gone blind from mustard gas poisoning, were received last year. Experience has proved that those who come to St. Dunstan’s in the early days of their blindness readjust themselves quickest to their new condition, and that neither home nor hospital is the best place for such adjustment. As many blind people have administrative and executive posts and help in the teaching and welfare departments at St. Dunstan’s, newcomers are taught by example how to reconcile themselves to their deprivation. To-day, 120 war-blinded are chartered physiotherapists, 140 telephone operators, 100 and more shopkeepers, 550 industrial workers, and some are doctors, solicitors, company directors, and welfare officers.

MALIGNANT CHANGE IN ERYTHEMA AB IGNE

BY

G. A. GRANT PETERKIN, M.B.E., M.B.
F.R.C.P.Ed.

Physician-in-charge, Skin Department, Royal Infirmary,
Edinburgh

[WITH SPECIAL PLATE]

The reddish-brown marbling on the skin of the lower legs often found in women who sit long in front of a fire is so common in Britain that it is taken as a matter of course and little attention is paid to it in our textbooks of dermatology. In these it is dismissed in a few lines as erythema ab igne or ephelis ab igne, though the word *epheles* (freckle) is wrongly used. Hartzell (1912) pointed out that the disease is essentially an inflammatory change and not simply a discoloration of the skin with blood pigment.

Possibly because of the ubiquity of central heating, the dermatosis appears to be rare in the United States, for Hartzell (1912) remarked that cases he has shown at meetings had not been diagnosed by competent dermatologists. He stated that “English dermatologists are apparently the only ones who see it with any degree of frequency.” In the same paper he described four cases of this disease and observed that none of the patients were in good health—a significant point. It seems likely that the condition is still often unrecognized in the U.S.A., for Hyman (1954) in discussing a case shown at the New York Academy of Medicine as purpura annularis telangiectodes (Majocchi’s disease) remarked that “the lesions are neither annular nor telangiectatic. So the two important features of Majocchi’s disease are lacking. In England one often sees this peculiar kind of mottling of the skin, with nothing telangiectatic about it at all, known as erythema ab igne and due to sitting in front of the fire.”

In the standard French textbooks, such as the *Précis de Dermatologie*, no reference to the disease can be found, though Erasmus Wilson (1863) quoted Rayer, who described “epheles ignealis in women sitting over charcoal burners, the histological appearances being those of congestion followed by chronic inflammatory thickening, and even vesication and considerable pigmentation is produced.”

In Algiers, Montpellier (1921) described this same reticulate condition as occurring on the inner aspect of the thighs in Moslem women from the use of a little charcoal heater held between the thighs in winter.

But erythema ab igne is probably most common in the colder parts of Asia; for instance, in Kashmir, in cold weather the inhabitants carry beneath their clothing a warming-pan or *kangri*, the temperature of which may reach 150° F. (65.5° C.). Neve (1910) described *kangri* cancer which followed heat burns similar to erythema ab igne. Within a period of fifty years Neve (1929) saw more than 2,000 cases of such cancers. Similar to these is the reaction seen in some parts of China where the people sleep on a heated brick bed or *kang*, so that the skin over the trochanter is burned and *kang* cancer, as described by Laycock (1948), may be provoked. An editorial in the *British Medical Journal* (1954) on “Skin Cancer and Burns” refers to these cancers, and to others such as the well-known carcinoma of the lips in clay-pipe smokers.

R. FINLAYSON AND J. O. ROBINSON: GIANT-CELL ARTERITIS OF LEGS

FIG. 1.—Left posterior tibial artery: occlusion of lumen by connective tissue containing moderate numbers of cells, with diffuse cellular infiltration of media. (Haematoxylin and eosin. $\times 24.5$.)

FIG. 2.—Left anterior tibial artery: proliferation, splitting, and bulging of internal elastic lamina. Small segmental deficiency of lamina adjacent to calcified medial plaque. (Weigert-van Gieson. $\times 28$.)

FIG. 3.—Part of wall of left anterior tibial artery: intense chronic inflammatory cellular infiltration of media, with foci simulating tuberculous follicles. Smooth-muscle cells clearly identified only by adventitia on right. (H. & E. $\times 75$.)

FIG. 4.—Left posterior tibial artery: infiltration of media with giant cells, lymphocytes, plasma cells, and histiocytes. Irregular staining of internal elastic lamina with calcification. (Haematoxylin and eosin. $\times 165$.)

FIG. 5.—Small artery from right tibialis anterior muscle: focal infiltration of media, adventitia, and perivascular connective tissue with lymphocytes and giant cells, the latter related in part to small areas of calcification. (Haematoxylin and eosin. $\times 80$.)

P. F. MILNER: NASAL GRANULOMA AND PERIARTERITIS NODOSA -6-4

FIG. 1.—Biopsy of gingival tumour, showing multinucleate giant cells in the dermis. (Haematoxylin and eosin. $\times 60$.)

FIG. 2.—Skin over right deltoid: normal vessel in subcutaneous fat surrounded by organizing haemorrhage. (H. & E. $\times 65$.)

FIG. 3.—Kidney, showing hyaline necrosis and fibrosis of glomeruli with oedema and inflammatory exudate. (H. & E. $\times 60$.)