EMERGENCIES IN GENERAL PRACTICE

CORONARY THROMBOSIS

BY

A. MORGAN JONES, M.B., M.Sc., F.R.C.P.*Director of the University Department of Cardiology, Manchester Royal Infirmary*

One of the physician's principal responsibilities in this condition is to recognize when it is likely that coronary thrombosis has occurred. It is a serious mistake to believe that the onset is always marked by a dramatic attack of prolonged chest pain. Indeed, there are at least four other modes of onset: sudden onset of angina of effort; sudden exacerbation of existing angina of effort; the appearance of short attacks of cardiac pain at rest; and sudden onset of heart failure without apparent cause.

In any of these circumstances it is now commonplace to find electrocardiographic evidence of recent myocardial infarction, but if the electrocardiogram does not show any evidence of recent infarction this should not be regarded as proof of the innocence of the symptoms. The clinical assessment of the case must be made independently of the electrocardiogram. In fact, the investigation is useful only when the result is positive, and then its value probably lies as much in the handling of the patient and his relatives as in clarifying the clinical diagnosis. In the absence of a positive electrocardiogram, assessment turns principally on whether the pain is definitely cardiac. Pain at rest is clearly more difficult to assess than effort pain. Inquiry should therefore always be made for previous effort pain, for if typical angina of effort is followed by rest pain of similar character and distribution (though possibly more severe or widespread) it is very likely that the rest pain is also cardiac.

The site of cardiac pain is very variable: it may occur anywhere between the mouth and the umbilicus, anteriorly or posteriorly, or in either arm. Contrary to some published assertions, it may be confined to the left mammary region or to the back of the chest. The character is more helpful than the site, for it is usually described as tight, heavy, constricting, pressing, aching, or burning but not as stabbing or shooting. It is not usually most severe or widespread at the onset, but tends to increase in severity and to radiate to a wider area simultaneously. It more commonly starts centrally in the chest and radiates peripherally to the neck, jaw, or arms, but the reverse can occur.

Among the more important differential diagnoses are acute pericarditis, dissecting aneurysm of the aorta, pulmonary embolism, hiatus hernia, and spontaneous pneumothorax, though in my experience the last only occasionally simulates coronary thrombosis. Both acute pericarditis and massive pulmonary embolism may be associated with T-wave changes in the electrocardiogram but not with the typical pattern of infarction. The sudden onset of an arrhythmia, such as paroxysmal tachycardia, is sometimes associated with quite severe cardiac pain without myocardial infarction, but if such pain lasts more than half an hour infarction should be suspected. T-wave changes may follow a paroxysmal arrhythmia although no infarction has occurred.

Severity of the Infarction

The severity of the pain is of no significance in assessing the severity of the myocardial infarction, but intractable or

recurrent pain over a period of more than forty-eight hours is, in my experience, a somewhat ominous feature. Maintenance of the blood pressure is a good sign, but an immediate considerable fall is not necessarily serious provided, as is often the case, evidence of shock is absent, for frequently there will be recovery from this early fall within twenty-four or forty-eight hours. A steady progressive fall over the first few days is more important, for in these circumstances the pressure is apt to remain low. Considerable fever and a leucocytosis of 20,000 or more are usually associated with extensive myocardial necrosis. Most of these criteria are concerned with the severity of the acute lesion, but, in fact, the prognosis really depends upon the total cardiac damage, old and recent. For this reason a history of old infarction, the presence of cardiac enlargement, and especially previous or present heart failure are important. The appearance of presystolic gallop rhythm or an attack of cardiac asthma may indicate the presence of left ventricular failure, and examination will then often reveal moist sounds at the lung bases. Right heart failure will be marked by engorged neck veins or oedema of the ankles and sacrum. Such signs are ominous.

Pericardial friction, usually present only for twenty-four to forty-eight hours, is probably an indication of a fairly large infarct, but in my experience it is not so ominous a sign as has sometimes been suggested. Cardiac arrhythmias, such as auricular flutter or fibrillation, and ventricular paroxysmal tachycardia are serious and require prompt treatment, but occasional premature beats are exceedingly common and not often followed by more serious arrhythmias; so they are probably better left untreated to disappear spontaneously in a few days. Very frequent ventricular premature beats have greater significance.

Management of the Acute Attack

The objects of immediate treatment are to allay anxiety, to relieve pain, to treat shock or heart failure, and to control serious arrhythmias. Nothing which the physician can say will allay anxiety so effectively as an unruffled and matter-of-fact approach to the patient, and if the doctor is sometimes himself anxious it is worth remembering that recovery is the rule if the patient is still alive when he arrives. The relief of pain is of immediate importance. It is traditional to give large doses of morphine, usually $\frac{1}{2}$ gr. (32 mg.). I doubt if it is desirable, however, to give repeated large doses of morphine in an attempt to remove all pain, for although this may induce sleep, complete relief of pain without unconsciousness is not, in my experience, attainable except by natural remission. Furthermore, very large and especially repeated morphine injections may lead to vomiting and later to troublesome constipation and lack of appetite. I therefore prefer the minimum dose which will make the pain tolerable, rather than rigid adherence to the large doses traditionally recommended in textbooks. If morphine in moderate doses does not make the pain tolerable, oxygen, preferably by tent, will sometimes help. Once the pain has eased, it is better to avoid morphine altogether.

There is no really satisfactory treatment for shock due to extensive myocardial damage, and most of the measures suggested, such as 50% glucose intravenously or small transfusions, are positively dangerous because they may precipitate acute heart failure. The best plan is to leave the patient where he is and keep him reasonably warm, but not to heap blankets upon him to near suffocation. If severe hypotension (80 mm. Hg) develops it may be life-saving to raise the blood pressure to 100–110 mm. Hg

by a noradrenaline drip (1:10,000), for this will maintain the myocardial and renal circulations. This technique requires very careful observation, for the blood pressure can be varied almost instantaneously by altering the drip rate, and readings must therefore be taken every fifteen minutes and the rate varied correspondingly.

Treatment of acute left heart failure with pulmonary oedema follows the usual lines, but I would like to emphasize the value of the slow intravenous injection of 0.5 g. of aminophylline in 20 ml. of solution.

Arrhythmias such as auricular flutter and fibrillation are treated by the usual methods, but the most rapid method of control should be chosen—that is, by intravenous digoxin.

Admission to Hospital

It is very doubtful whether any community will in the future be able to provide enough hospital beds for the treatment of all cases of coronary thrombosis, even if this were desirable. In my view, cases in which anticoagulant therapy is to be used should be admitted to hospital at the present time, for at home satisfactory laboratory control of this treatment is exceedingly difficult, though perhaps not impossible. In general, therefore, the present criterion for hospital admission is the desirability of anticoagulant treatment.

So far there is no uniformity of opinion about the desirability of anticoagulants, for all shades of opinion exist in this country from that of William Evans—"That anticoagulant treatment of coronary occlusion will go the way of other discarded remedies is certain. Let it go now"—to that of Rae Gilchrist—"Anticoagulants are justified by a halving of the death rate and by an even greater reduction in thrombo-embolic episodes." In controlled series published throughout the world the mortality in cases treated with anticoagulants has averaged 16.6% in 1,091 cases, whereas in the corresponding control series treated by conservative methods the mortality averaged 32.4% in 1,344 cases. If, however, we separate off what are termed "good risk" cases we get a somewhat different picture. A patient with recent coronary thrombosis is generally regarded as a "good risk" if there has been no previous infarction, if there is no intractable pain, no severe shock, and no cardiac enlargement, gallop rhythm, heart failure, or significant arrhythmia. In addition there should be no indication of a liability to thrombo-embolism, such as varicose veins in the legs or a previous history of thrombophlebitis or pulmonary embolism.

These criteria may appear to eliminate a large proportion of cases, but in fact they do allow about half the cases to be regarded as "good risks." In 489 such cases recently reviewed the mortality was only 3.1% and the reduction of mortality by anticoagulants only 1%. In such cases the gain from the use of anticoagulants is relatively small and against that gain must be set the risk of transport to hospital and the risk of anticoagulant therapy. It is true, of course, that a "good risk" case at the outset may, owing to further infarction, become a "bad risk" case. These patients include, of course, the 3% who die. It has often been held that uncertainty of the outcome is one of the outstanding clinical features of coronary thrombosis, but there must be few serious conditions in which we can at the onset separate off clinically so large a proportion of cases with so hopeful and encouraging a prognosis. My experience has certainly been in conformity with this view, and my own practice is, after careful assessment, to encourage the treatment of "good risk" cases at home by conservative methods. In the "bad risk" case, on the other hand, I very strongly advise anticoagulant treatment in hospital, for I believe that in these cases all the available evidence indicates that anticoagulants substantially reduce the mortality.

Control of Anticoagulant Therapy

Anticoagulant treatment can be controlled *only* by regular estimation of prothrombin activity in the blood, and this

is usually necessary daily at first. Even if a patient's anticoagulant requirement is known on one occasion it may be quite different in a subsequent illness. The institution of anticoagulant therapy should therefore always be undertaken in hospital unless reliable daily laboratory facilities are available at home. Once it has been shown that the patient's requirement of anticoagulants is steady, continuation may be possible at home with weekly or twice weekly prothrombin estimations.

In my department anticoagulant treatment in coronary thrombosis follows the same general lines as those laid down by Rob for peripheral arterial occlusion in his article earlier in this series,* but heparin is used at the onset only if there is initial shock. Prothrombin activity is estimated on admission and a low initial reading is regarded as an indication of liver damage; if below 50% anticoagulants should not be given, and if between 50 and 75% caution is necessary. A history of recent spontaneous bleeding (for instance, from a peptic ulcer) is a contraindication. The initial dose of phenindione is 200 mg. by mouth if the preliminary prothrombin level is normal. All subsequent doses are controlled entirely by the prothrombin level, which should be reduced to 20%. The stabilized daily requirement varies from 25 to 250 mg. and may fluctuate from time to time. For subsequent out-patient treatment it is therefore safer to aim at a level of 30%. In the early days of anticoagulants spontaneous and sometimes fatal bleeding from bowel, kidneys, or into the tissues often occurred, but with careful control haemorrhage is now very uncommon and almost never fatal. The most effective antidote is vitamin K₁ orally (20 mg.), but this is still difficult to obtain, and injected "synkavit" (50–100 mg.) is useful but less effective. Anticoagulants once started should be continued until the patient is up for part of the day.

General Management

The following observations apply whether the patient is treated conservatively at home or with anticoagulants in hospital.

Reassurance.—Once a patient knows he has had a coronary thrombosis—and this cannot usually be concealed from him—anxieties crowd upon him. His heart, to which he may never have given a thought, has let him down, and this is, in the popular view, the vital organ; he fears, first, that he may lose his life; then that he may be permanently crippled; and, finally, although he may survive free from symptoms, he fears a recurrence. He may say little of these fears, yet he wants them allayed. In my view it is his physician's duty to reassure him, and in great part this is medically justified. The majority do survive and return to a normal, if slightly less hectic, life; if the condition recurs, it may not do so for very many years, and he may again make as good a recovery as before. Older views on prognosis are unjustifiably gloomy. The technique of reassurance is important, for exaggerated or over-dogmatic statements will be discounted by an intelligent patient. His confidence will lie in the physician who himself believes and acts and talks at all times with a quiet confidence in ultimate recovery; who assumes that, after a space of time for complete healing, resumption of a normal life is the rule. In this atmosphere those who recover will do so with patience and determination, and without unnecessary uncertainty and fear.

Rest.—Rest for four to six weeks is still generally regarded as desirable in coronary thrombosis, but the days of rest flat in bed—rigidly immobile, being fed, and being uncomfortably balanced on bedpans—are, if not gone, at least, I hope, numbered. If this spirit of tense apprehension dominates the sickroom the patient's future may well look as blank to him as the ceiling which is made to form almost the whole of his field of vision. We know, both from clinical experience and from scientific measurement, that the

*"Sudden Occlusion of a Major Peripheral Artery," by Professor C. G. Rob (*Journal*, February 12, 1955, p. 407).

physical effort of using a commode is much less than that of maintaining a precarious balance upon an apparently razor-edged bedpan which tilts unpredictably upon a yielding mattress. Rigid immobility in bed predisposes to thrombo-embolism, as does over-sedation. In the more severe cases, and especially those with shock, the patient may be fed and shaved for the first week, but whatever level of rest is imposed at first it should be progressively, though gradually, relaxed so that each week the patient feels that he is making some progress. In the last (fourth to sixth) week before the patient gets out of bed, breathing and leg exercises will help to restore his flabby muscles and build up his morale. In old people some period out of bed in a comfortable chair may well be instituted from the end of the first week. The keynote of the management of activity is gradual increase from the end of the initial phase of shock (if present), lack of appetite, and general malaise, which usually lasts only a few days. After that time most patients feel so well that it is necessary to explain to them the need for a sufficient period of comparative rest to ensure sound healing.

Drugs.—Apart from a gentle sedative during the day and perhaps a soporific at night, anticoagulants are generally the only medicinal treatment necessary. Too much attention to the bowels is to be discouraged in the first week, for a few days of "constipation" are physiological following the days at the onset when little food is desired.

Heart Failure.—Constant watch should be kept for early signs of heart failure, especially sacral oedema, cervical venous engorgement, a tendency to breathlessness in bed, and basal pulmonary congestion. The legs should be examined regularly for oedema or thrombophlebitis. Treatment for early heart failure, if necessary, follows the usual lines. If there is any tendency to heart failure the patient should be nursed sitting—oedema in the lower back or ankles never killed anyone, but fluid in the lungs is much more dangerous. For nocturnal breathlessness, associated with early left ventricular failure, "nepenthe"—30 min. (1.7 ml.)—at night is often an effective immediate prophylactic, but treatment should be by mercurial diuretics and salt restriction.

Visitors.—Some limitation is desirable in the first three weeks, and generally this can be effected only by confining the privilege to the family. But if a single urgent business matter needs attention, it is better to allow a visit for this purpose rather than permit the patient to worry about it. After the third week a telephone may be made available and short visits from friends encouraged.

Diet.—Light, small meals should be the rule, and this is especially true if the patient is overweight, but in any case there is a tendency to gain weight in bed and this should be guarded against. From the third week the patient should be weighed weekly at the bedside.

Tobacco and Alcohol.—Although there is no satisfactory evidence that tobacco is harmful in coronary thrombosis, it is probably wiser to prohibit it during the first two weeks and then, if the patient clearly feels the need for it, to allow not more than ten cigarettes a day. There is no contra-indication to the moderate use of alcohol, except that allowance must be made for the additional calorie intake.

Details of management can be described without end, but all decisions finally depend upon a common-sense approach which recognizes the patient's need for outward signs of steady progress and his dependence upon a background of quiet confidence.

Extension of Infarction

It is my practice to have a weekly electrocardiogram, for if a recent reading is available for comparison this assists in the interpretation of any recurrence of cardiac pain. The degree of change in serial records, however, has never seemed to me to bear much relation to the clinical progress of the patient. Pathological evidence from our recent

studies indicates that extension of an infarct is usually due to necrosis of ischaemic muscle related to the original occlusion and not due to fresh coronary occlusion. This tendency for extension exists for about four weeks, and since it does not imply a new coronary thrombosis I doubt if it should be allowed to increase the period of rest very much.

Convalescence and Subsequent Life

If the management of the period of bed rest has been psychologically sound the patient should come to the time of ambulation with an expectation of ultimate satisfactory recovery, but a realization that this will take some weeks or months. He will not expect to get up and go about his business, for he will know that owing to the period of bed rest his muscles will be flabby and will require progressive use to recover, and he will have been warned that some two months must elapse before the final stages of cardiac healing are complete and he can resume his life in a more or less ordinary way. Although the pace at which convalescence proceeds may be modified by the total cardiac damage and the temperament (either apprehensive or optimistic) of the patient, it is a good general rule to prescribe a month of gradually increasing activity at home followed by a quiet holiday of three or four weeks in a flat place. It is only during the first week at home that stairs need be avoided, provided they are taken quietly. A month after he returns home he should be able to drive his car locally, but not on long journeys. Abstinence from sexual intercourse during the period of convalescence of three to six months is probably advisable.

In the majority of cases in which heart failure has not occurred the patient will be able to resume the sort of life he has led before, provided this was a reasonable one. He will need only to avoid excesses of any kind: smoking, drinking, eating, work, idleness, worry, long hours, etc. If he drives a car for long distances, he should do so in easy stages. The golden rule for him is to allow plenty of time for anything he does, from catching a train to eating a meal. But a coddled and cosseted life of inactivity is as bad, or worse, than too busy a life, with snatched meals and hardly caught trains. In fact, he should lead the sort of life we all ought to lead—in full health we may escape retribution if we ignore the rules of a sensible life, but we cannot afford this indulgence after a coronary thrombosis.

Except for the heaviest manual job and work involving responsibility for the lives of others—for example, bus-drivers, engine-drivers, aircraft pilots—there are few occupations which a man is justified in leaving because he has had a coronary thrombosis, provided this has not been accompanied by heart failure. But he may have to plan his work more efficiently, especially if it is largely administrative, and learn to depute responsibility. Amongst manual workers, experience shows that, in the coronary age group, attempts to move men to lighter occupations often fail because they either return to their old work or give up altogether.

Conclusion

Finally, all that I have seen of coronary disease has convinced me that the management of the patient is even more important than the treatment of the disease: there is often far more residual disability from fear than from myocardial damage—and it is largely within our power to prevent this.

Next article on Emergencies in General Practice.—"Birth Trauma," by Dr. A. P. Norman.

Refresher Course Book.—Copies of the second volume of collected articles from the Refresher Course for General Practitioners are still available at 25s. (postage—inland 1s. 6d., overseas 1s.) each from the Publishing Manager, B.M.A. House, Tavistock Square, London, W.C.1, or through any bookseller. The first volume is now sold out.