gest as an answer the two infections which are most akin to infantile gastroenteritis—namely, salmonella enteritis and bacillary dysentery.—I am, etc.,

A. G. IRONSIDE.

Department of Infectious Diseases,
Monsall Hospital,
Manchester 10.

Chemotherapy of Burkitt's Lymphoma

SIR,—An observation in the treatment of a patient with Kaposi's sarcoma led to my combining potassium iodide with cyclophosphamide in the treatment of Burkitt's lymphoma. The results to date suggest that potassium iodide potentiates the action of cyclophosphamide.

The Figure shows the survival curves for 24 cases of Burkitt's lymphoma, of which 23 were treated with cyclophosphamide alone and one with methotrexate; and for 10 cases given the combined treatment. Although sufficient time has not yet elapsed to be sure that the levelling out of the survival curve will remain at 65%, yet the difference is so great when compared with the 21% survival for the 24 cases that it would seem that potassium iodide may be a valuable adjuvant to cyclophosphamide in the treatment of this and other tumours. Clinically the recession of the growths is more rapid, and complete remissions have been seen with large and multiple tumours. Of the 24 cases 12 are stage 1 and 2, and 12 stage 3 and 4. Of the 10 cases five are stage 1 and 2, and five stage 3 and 4.

All the patients did not start treatment at the same time, and because of the intervals between entrance to the inquiry some of the patients have been under observation for a much longer time than others. In order to surmount this difficulty a life-table method of obtaining survivors has been employed.

Study of the available literature suggests that potassium iodide may inhibit cell glycolysis, rendering the cell membrane more permeable to cyclophosphamide.¹ But it is also noted that iodides activate dihydrofolate reductase in purine synthesis² and so it is felt that caution is called for, since potassium iodide administered alone may reactivate a tumour. I have therefore adopted the following treatment regimen. The dosage of oral cyclophosphamide is calculated at 40 mg. per pound (500 g.) of body weight. This is given at the rate of 300 mg. per day, and at the same time potassium iodide 600 mg. twice daily is given, but the administration of potassium iodide must cease the day before the total dose of cyclophosphamide is achieved. Concurrent leucocyte counts must

be performed, as there is frequently a sharp fall in these.

As my series is small and would take a long time to become large I am asking you to publish this letter so that others who may be interested in carrying out trials may do so without delay.—I am, etc.,

E. H. WILLIAMS.

Kuluva Hospital,
Arua, Uganda.

REFERENCES

- ¹ Bergel, F., in *The Treatment of Cancer*, ed. by J. S. Mitchell, 1965, p. 9. London, Cambridge University Press.
- ² Shelley, W. B., *Journal of the American Medical Association*, 1967, 201, 1009.

Mononucleosis and Burkitt's Lymphoma

SIR,—I was interested in the medical memorandum by Dr. L. A. Price and Mr. D. Garfield Davies (5 April, p. 31), reporting a case of lymphosarcoma with a positive Paul-Bunnell test. Could this have been an atypical case of Burkitt's lymphoma?

Some authors feel that Burkitt's tumour is a form of lymphosarcoma.¹ Moreover, there is evidence to suggest that Epstein-Barr virus causes infectious mononucleosis in Britain and Burkitt's lymphoma in Africa. The reason for this is obscure, but the following explanation is suggested.

It is now well established that immunosuppression predisposes to malignant disease.^{2,3} In addition, it has also been noted that patients suffering from the coeliac syndrome have an increased incidence of lymphomata. Furthermore, these cases can often be identified by hypoproteinaemia and occasionally oedema.^{4,5} It may be that in malabsorption immunoglobulin synthesis is impaired, thereby leaving these patients unable to combat oncogenic virus or dispose of neoplastic cells. Such virus might attack the diseased bowel in the first instance, but extra-abdominal lymphomata have also been noted. In the case of Burkitt's tumour, it may be that African children, if not suffering from frank kwashiorkor and hypoproteinaemia, have at least an element of malnutrition leading to defective synthesis of immunoglobulin, which need not necessarily be quantitative. Again, Burkitt's lymphoma is unknown during the first year of life, when presumably these children still have passive immunity.

Finally, I would agree with your authors' explanation that their case is not a coincidence, but rather a true lymphoma presenting with a positive Paul-Bunnell test. At the same time it would be interesting to know how many other cases of lymphosarcoma have a positive Paul-Bunnell test in the very early stages of the disease. Certainly, in a recent case of reticulum cell sarcoma the Paul-Bunnell was negative some weeks prior to the correct diagnosis.⁶—I am, etc.,

W. S. B. LOWRY.

Royal Victoria Hospital,
Belfast.

REFERENCES

- ¹ Iversen, O. H., and Harket, R., *European Journal of Cancer*, 1968, 4, 383.
- ² *Lancet*, 1969, 1, 505.
- ³ Lowry, W. S. B., *Lancet*, 1969, 1, 731.
- ⁴ Brunt, P. W., Sircus, W., and Maclean, N., *Lancet*, 1969, 1, 180.
- ⁵ Eidelman, S., Parkins, R. A., and Rubin, C. E., *Medicine*, 1966, 45, 111.
- ⁶ Lowry, W. S. B., *Postgraduate Medical Journal*, 1968, 44, 926.

Diverticulosis of the Colon and Diet

SIR,—Mr. M. N. Morgan and Professor H. Ellis did not find diverticula in rats fed on a low residue diet for 50 to 70 weeks (5 April, p. 53) in contrast to the experience of Carlson and Hoelzel,¹ who found that such a diet produced the disease in rats in life-span studies. Their timely warning obviously was given so as to save other workers from terminating the existence of their experimental rats too soon, but it would be a pity if their letter fostered the belief that the clue to the causation of diverticular disease is most likely to be found in rats and animals. Furthermore, it is not true that the "aetiology of the disease is essentially unknown." The following facts are known:

Diverticular disease occurs in people who eat a low residue diet in which both flour and sugar are refined. The disease is not found in people living tribally who eat unrefined maize, but it appears in these people or their descendants if they change to a "civilized" diet lacking natural fibre. As would be expected in a disease which occurs in the latter half of life, this does not occur until after a latent period of some years.

Diverticulosis has been related to an increase in localized intrasigmoid pressures.²⁻⁴ These high localized pressures occur more easily in a narrow colon which is segmenting vigorously. A refined diet results in a more viscous stool that passes through the gut more slowly than the bulky stool of a high-residue diet. This can be proved by anyone who is prepared to eat either an unrefined diet or is willing to add unprocessed bran to his present diet. Is it not possible that this alteration in the faeces due to diet has an adverse effect on the intracolonic pressures, and hence leads to diverticula appearing in those who consume such a diet?

The incidence of the disease in England and Wales is increasing steadily since the beginning of this century, and, allowing for a latent period of about 40 years, this rise corresponds to the increased consumption of refined flour and sugar that has occurred in this country. The same is true of several other common diseases that attack civilized man.⁵

It is interesting to note that despite the fall in the mortality that follows perforated diverticulitis that resulted from the advent of the antibiotics, advances in anaesthesia, and better surgical management, the death rate from the disease steadily increased in England and Wales except for the years 1939-52.

FIG. 1.—Accrued death rate from diverticular disease in England and Wales, 1923-1966. From *Diabetes, Coronary Thrombosis, and Saccharine Disease*, by T. L. Cleave, G. D. Campbell, and N. S. Painter, reproduced by permission of John Wright & Sons Ltd., Bristol.