days for one to two weeks; they noted no toxicity. Sutton (1938) also remarked on its lack of toxicity when used as a local application. However, Slotkowski (1957) reported the case of a child who developed ulceration of his oral mucosa as a result of excessive crystal violet treatment. He remarked on the need to be aware of this possible complication of treatment and suggested that it might previously have been mistaken for a result of the candidial infection.

At other sites crystal violet is well recognized as a potential irritant. When vaginal candidiasis is treated with local applications of crystal violet some patients have to discontinue treatment because of its irritant effects (Jennison and Llywelyn-Jones, 1957). If the dye is in accidental contact with the conjunctival epithelium it will cause oedema, hyperaemia, and necrosis of the epithelium within a few hours (Hosford and Smith, 1952). When taken orally as a vermifuge, crystal violet may cause diarrhoea, nausea, and vomiting, suggesting a local effect on the gastrointestinal mucosa (Beckman, 1961; Martindale, 1967). It might therefore be expected that similar reactions could occur when crystal violet is applied to the oral mucosa.

It seems likely that many oral mucosal reactions to crystal violet have gone unrecognized, and it is clear that if crystal violet is used it should be limited to a maximum course of three

days with twice-daily applications. As nystatin suspension now provides a satisfactory method of treating oral candidiasis, there would seem to be a substantial case for the abandonment of local crystal violet applications to the oral mucosa. It might be that many sore mouths associated with thrush would disappear along with the crystal violet.

I wish to thank Dr. B. D. R. Wilson for his advice and for permission to report these cases.

R. W. JOHN, M.B., M.R.C.P., D.C.H.,
Registrar, Children's Department,
St. Thomas's Hospital, London S.E.1.

REFERENCES

- Beckman, H. (1961). *Pharmacology*, 2nd ed., p. 576. Philadelphia and London.
Faber, H. K., and Clark, E. B. (1927). *Amer. J. Dis. Child.*, 34, 408.
— and Dickey, L. B. (1925). *J. Amer. med. Ass.*, 85, 900.
Hosford, G. N., and Smith, J. G. (1952). *Ibid.*, 150, 1482.
Jennison, R. F., and Llywelyn-Jones, J. D. (1957). *Brit. med. J.*, 1, 145.
Martindale, W. (1967). *Extra Pharmacopoeia*, 25th ed., edited by R. G. Todd, p. 561. London.
Slotkowski, E. L. (1957). *J. Pediat.*, 51, 652.
Sutton, R. L. (1938). *J. Amer. med. Ass.*, 110, 1733.

Case of Toxoplasma Polymyositis

Brit. med. J., 1968, 1, 158-159

In the few reports of severe acquired toxoplasmosis the predominating features have been meningoencephalitis, myocarditis, and pneumonitis. Though myalgia, fatigue, and muscle weakness have been noted in several patients and the parasite has been demonstrated in muscle biopsies showing muscle necrosis and inflammatory cellular reaction, in only one case have symptoms of polymyositis dominated other manifestations of the infection (Rowland and Greer, 1961). So far as we are aware, the case of polymyositis reported here, which in our opinion is due to toxoplasmosis, is the second of its kind in the literature and the first to be reported in this country.

CASE REPORT

The patient, a 7-year-old boy, was taken ill in the second week of September 1966 with sore throat, malaise, and fever. The sore throat lasted a week but the malaise and fever persisted. During the second week of illness discrete and tender swellings appeared on both sides of his neck. A diagnosis of glandular fever was made and the boy was advised to rest at home. A fortnight after the onset of illness he felt weak and began to tire easily. His parents noticed that he walked with difficulty and fell several times when going to the lavatory. At the same time he complained of pain in his arms and thighs, and his mother commented that his arms appeared swollen. He was admitted to hospital on 11 October.

On admission his temperature was 99.6° F. (37.6° C.), pulse 128, and respiration 20. The lymph nodes in both posterior cervical triangles were enlarged, discrete, and tender. He was fully conscious and well orientated. There was no neck stiffness and Kernig's sign was negative. The cranial nerves were all intact. There was no weakness of facial or bulbar muscles. There was, however, pronounced weakness of the flexor and extensor muscles of the neck and those of the trunk. He could not hold his head forward or horizontally when lifted off the bed, nor was he able to lift his head from the bed. He could not sit up from the recumbent posture

unaided or stand up from the sitting position, and, when helped to stand, assumed a lordotic posture but was unable to walk. There was pronounced weakness proximally and moderate weakness distally in the muscles of all four extremities. Both arms appeared swollen and there was slight tenderness of both arm and thigh muscles. The deep reflexes in the upper limbs were markedly depressed but were normal in the lower limbs, and the plantar responses were flexor. There was no sensory abnormality. The rest of the systemic examination showed no abnormality. There was no family history of myopathy or any other neurological disorder.

Investigations.—Haemoglobin 12.4 g./100 ml.; white blood cells 8,700/cu. mm. (polymorphs 65%, lymphocytes 29%, monocytes 6%); E.S.R. 26 mm./hr. (Wintrobe); platelets normal; total plasma proteins 7.5 g./100 ml., with a normal electrophoretic pattern; thymol turbidity 4 units; serum aspartate aminotransferase 550 units (normal, 35-107 units); serum alanine aminotransferase 680 units (normal, 22-92 units); serum creatine kinase 2,550 i.u./l. (normal, 0-60 i.u.); total creatine content of 24-hour urine 101 mg.; edrophonium chloride test negative; C.S.F. normal; Paul-Bunnell reaction, agglutination test for *Brucella abortus*, and latex test for rheumatoid arthritis and lupus erythematosus negative; no virus isolated from throat swab, faeces, or C.S.F.

Electromyography of the left deltoid showed no spontaneous discharge on needle insertion. On maximal effort, only small, short-acting, high-frequency muscle potentials were seen with occasional high-incidence polyphasic potentials.

Muscle biopsy was taken from the left deltoid and the histological report was as follows: "The overall architecture of the muscle looks normal, but there is definite pathological change in focal areas. This takes the form of degeneration of isolated fibres with associated cellularity around them. There is some round-cell infiltration and also excess fibroblastic activity. The only conclusion one can draw from this biopsy is that it is abnormal, and that the changes are focal in nature, consistent with polymyositis, and certainly not suggestive of a muscular dystrophy or general neurogenic atrophy type of pattern."

The symptoms and signs of glandular fever at the beginning of the illness and the subsequent negative Paul-Bunnell reaction raised the possibility of acquired toxoplasmosis. The toxoplasma dye test, carried out seven days after admission, was positive at a titre of 1:4,096.

Treatment and Course.—A presumptive diagnosis of toxoplasma polymyositis was made and treatment with pyrimethamine 12.5 mg. once daily and sulphadimidine 0.5 g. and folic acid 5 mg. thrice

daily was started on 20 October and continued for 28 days. Within two days his temperature, which had never been high, appeared to settle. Five weeks after starting treatment he was able to walk slowly unaided and could lift his head 20 to 30 degrees from the horizontal, though he was still unable to sit up straight. His serum creatine kinase 45.5 i.u./l. and serum alanine aminotransferase 40 units were within normal limits, but serum aspartate aminotransferase 148 units was still slightly raised. The dye test titre remained unchanged. Three months after onset the patient was discharged fairly well. He was able to sit and stand unaided, though he still walked with a stamping gait. After a further three months recovery was complete. He could play games and keep up with other children.

EPIDEMIOLOGY

The patient's family comprised his parents and an 11-year-old brother and a 12-year-old sister. The family kept a cat which had no history of illness. Samples of blood were taken on 11 November 1966 from members of the family for toxoplasma dye tests. Negative results were obtained from the father and brother. The mother was positive at 1:32 and the sister, who appeared perfectly well, was positive at 1:32,000. The mother now recalled that small lumps had suddenly appeared behind her daughter's left ear about two weeks before her son was taken ill. She had taken the girl to her family doctor, who confirmed the presence of enlarged lymph nodes. The sister was examined by one of us (K. C.), and though asymptomatic she was found to have a mild pyrexia, a generalized lymphadenopathy, and a palpable spleen. The pet cat was bled and found to have a dye test titre of 1:16. It was not possible to obtain the cat for further examination.

The patient and his sister were seen again in October 1967, 12 months after his admission to hospital. Both were perfectly well and neither showed any trace of the earlier lymphadenopathy. The boy's dye test was still positive at 1:4,096, but the girl's titre had fallen sharply to 1:512.

DISCUSSION

The diagnosis of polymyositis was based on the clinical picture supported by the creatinuria, the abnormally high serum enzyme levels, and the result of the muscle biopsy. The electromyographic findings, while not conclusive, were not incompatible with the diagnosis, since action potentials of diminished amplitude and duration may be the only abnormal changes seen in some patients with polymyositis (Dowben *et al.*, 1965).

The finding of a positive toxoplasma dye test titre of 1:4,096 in a previously healthy 7-year-old boy must be regarded as significant. According to Beattie (1967) antibodies to toxoplasma are rare in children over 6 months and under 10 years of age, and their presence strongly suggests recent infection. He is of the opinion that when evidence of a rising titre is not obtainable it is reasonable to consider a dye test titre of 1,000 or more as significant.

The "glandular fever" which preceded the polymyositis is a common manifestation of acquired toxoplasmosis. According to Beverley and Beattie (1958) glandular toxoplasmosis has accounted for about 7% of cases clinically diagnosed as "glandular fever" but giving a negative Paul-Bunnell reaction. The fact that the patient's sister, who had a positive dye test titre of 1:32,000, also developed a lymphadenopathy about two weeks before her brother fell ill suggests that both may have been infected at or about the same time. The girl, whose dye test titre fell sharply to 1:512 within 11 months, appeared to have dealt more efficiently with the organism than had the patient, whose dye test titre remained raised over the same period.

After the diagnosis of toxoplasmosis was made the original biopsy section was examined again but no lesions could be seen within the muscle fibre or in the perimysium suggestive of the

cystic forms of toxoplasmosis, nor could any of the vegetative forms be detected. However, what appeared to be a terminal colony was seen within the cytoplasm of a multinucleate cell just beneath the perimysium (see Fig.).

Showing what seems to be a terminal colony of *Toxoplasma gondii* in a multinucleate cell. (Haematoxylin and eosin. $\times 1,500$.)

The patient's response to treatment with pyrimethamine and sulphonamides, a combination of drugs found to be the most effective remedy for toxoplasmosis, lends some support to the diagnosis.

The source of infection remains uncertain. Toxoplasmosis has often been found in patients closely associated with animals. In the present case, though the pet cat showed serological evidence of infection it was never ill.

The cause of polymyositis is obscure. Because of the fever and pain and the pathological findings of inflammatory cellular reaction in the muscles an infective cause is held responsible by many workers, though bacteriological studies have not yielded convincing results. *Toxoplasma gondii* is known to produce a parasitaemia which may involve skeletal muscle as well as other tissues. As a rule invasion is not followed by any obvious harm, but the occasional emergence of a strain of high virulence results in clinical evidence of severe illness in the form of encephalitis, myocarditis, pneumonitis, or hepatitis. We would suggest that polymyositis may also be a clinical manifestation of toxoplasmosis and that evidence of infection with the parasite should be sought for in all patients with polymyositis of obscure origin.

We wish to thank Dr. G. B. Ludlam for carrying out the dye tests and Dr. V. Dubowitz and Dr. J. K. A. Beverley for the histological reports.

KRISHAN CHANDAR, M.B., B.S., M.R.C.P.,
Lately Medical Registrar, Groby Road Hospital, Leicester.

HÉLÈNE J. MAIR, M.D., M.C.PATH., DIPL.BACT.,
Consultant Virologist, Public Health Laboratory Service,
Leicester.

N. S. MAIR, M.B., F.C.PATH., D.C.H., D.P.H., DIPL.BACT.,
Director, Public Health Laboratory Service, Leicester.

REFERENCES

- Beattie, C. P. (1967). In *Recent Advances in Medical Microbiology*, edited by A. P. Waterson, London.
- Beverley, J. K. A., and Beattie, C. P. (1958). *Lancet*, 2, 379.
- Dowben, R. M., Vawter, G. F., Brandfonbrenner, A., Sniderman, S. P., and Kaegy, R. D. (1965). *Arch. intern. Med.*, 115, 584.
- Rowland, L. P., and Greer, M. (1961). *Neurology (Minneapolis)*, 11, 367.