We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer. Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.

LETTERS

CHANGE PAGE

Way forward is not obvious at asthma crossroads

Barnes argues that using a combination inhaler (budesonide plus formoterol) as rescue therapy improves asthma control,¹ and several studies support its use in the SMART (Symbicort (budesonide/formoterol) maintenance and reliever therapy) regimen.²⁻³ They have shown longer time to first exacerbation, reduced rate of severe exacerbations, and less inhaled corticosteroid dose, although with similar improvement in symptoms, peak flow rates, and quality of life in some studies.

However, the main concern is that all the above studies were conducted by the manufacturer (study design, data interpretation, data analysis, and publication). Three studies from the rival manufacturers of fluticasone/salmeterol have shown the opposite effect—stable dosing reducing exacerbation rate and improving symptom free days compared with the SMART regimen.⁴

Secondly, in all SMART studies patients needing more than 10 inhalations were excluded.

Thirdly, there is no convincing evidence that in patients who are well controlled with stable dose inhaler therapy a change to the SMART regimen will be cost effective.

Of greater interest is the debate on once daily asthma maintenance therapy compared with twice daily regimens.⁵ Certainly, asthma treatment is at a crossroads awaiting a change of direction.

Syed F Hussain consultant respiratory physician, Kettering General Hospital, Kettering, Northamptonshire NN16 8UZ
Syed.Hussain@kgh.nhs.uk

Competing interests: SFH has received sponsorships for organising a CME programme, participating in conferences, and speaking from Astra Zeneca, GSK, and MSD.

- 1 Barnes PJ. Using a combination inhaler (budesonide plus formoterol) as rescue therapy improves asthma control. *BMJ* 2007;335:513. (8 September.)
- 2 Rabe KF, Atienza T, Magyar P, et al. Effect of budesonide in combination with formoterol for reliever therapy in asthma exacerbations: a randomised controlled, double-blind study. *Lancet* 2006;368:744-53.
- 3 Kuna P, Peters MJ, Manjra AI, et al. Effect of budesonide/formoterol maintenance and reliever therapy on asthma exacerbations. *Int J Clin Pract* 2007;61:725-36.
- 4 Price DB, Williams AE, Yoxall S. Salmeterol/fluticasone stable-dose treatment compared with formoterol/budesonide adjustable maintenance dosing: impact on health-related quality of life. *Respir Res* 2007;8:46.
- 5 American Lung Association Asthma Clinical Research Centers, Peters SP, Anthonisen N, Castro M, et al. Randomized comparison of strategies for reducing treatment in mild persistent asthma. *N Engl J Med* 2007;356:2027-39.

CARDIOVASCULAR PREVENTION

Risk factors inform screening

First degree relatives of patients with coronary heart disease are at higher risk than the general population.¹ Screening first degree relatives would be an improvement on unselected screening. However, family history is only one of several risk factors for cardiovascular disease. Selecting patients for screening on the basis of multiple risk factors is better than selecting patients on the basis of one risk factor. For most patients, many risk factors are known. All patients have an electronic record of their age and sex. Most have an electronic record of their smoking status and blood pressure, and some have a cholesterol measurement. If diagnosed, patients with diabetes have an electronic record of this diagnosis. If they have a family history of premature coronary heart disease, some patients have a record of this fact.

Earlier in 2007 I collected electronic data from three large general practices. Of 19 552 patients aged 35 and older, 35% had full risk factor information, and 76% lacked only a cholesterol measurement. Of these, 2.2% (437) were aged 35-74, untreated but at a greater than 20% 10 year cardiovascular risk and hence eligible for treatment. When patients whose cholesterol concentrations were not known were included, 6.7% (1307) were untreated patients at high risk. Only 14% (181) of these patients had a record of a family history of premature coronary heart disease.

Since patients over 35 are about half the practice population, this means that

in a typical practice about 3-4% of the population are currently untreated but have recorded risk factors indicating they are at high risk. These patients are identifiable by searching the practice database, without the need to wait until their relatives present to secondary care with heart disease.

Tom P Marshall senior lecturer in public health
University of Birmingham, Birmingham B15 2TT
T.P.Marshall@bham.ac.uk

Competing interests: None declared.

- 1 Chow CK, Pell ACH, Walker A, O'Dowd C, Dominiczak AF, Pell JP. Families of patients with premature coronary heart disease: an obvious but neglected target for primary prevention. *BMJ* 2007;335:481-5. (8 September.)

Here's to risk assessment

I was delighted to read the editorial suggesting that we should move away from the primitive "one threshold fits all" mentality for starting antihypertensive treatment and take a view based on the overall cardiovascular risk.¹ We already do this when treating cholesterol for the purposes of primary prevention, so it is inconsistent not to use this approach for blood pressure, which is another continuous variable. The recent Joint British Societies' guidelines recognise this as the predicted cardiovascular risk rises with systolic blood pressure to 160 mm Hg,² yet they are not used as a tool for assessing whether to treat hypertension.

This is part of the general problem that occurs when we assign arbitrary values to continuous and often fluctuating biological variables to create boundaries for disease labels. For example, bronchial hyper-reactivity can change quite notably over time, and it can be very difficult to decide whether the label of asthma is appropriate. Our target driven culture encourages the use of these labels, but I think that they are often not very helpful, and I often use asthma drugs in those whom I would not label as asthmatic.

Glucose metabolism represents another such variable, and I look forward to the day when I read an editorial suggesting that we abandon the World Health Organization's criteria for diagnosing diabetes, in favour of a decision tool for the treatment of abnormal glucose metabolism based on risk.

Jonathan D Sleath general practitioner, Hereford HR2 9HN
jonathan.sleath@nhs.net

Competing interests: None declared.

- 1 Nash IS. Reassessing normal blood pressure. *BMJ* 2007;335:408-9. (1 September.)
- 2 JBS 2: Joint British Societies' guidelines on prevention of cardiovascular disease in clinical practice. *Heart* 2005;91(suppl V):v1-52.

DRUG PRICING

Misconceptions about PPRS

Iheanacho argues that the Office of Fair Trading may have pinpointed some areas of improvement in the existing Pharmaceutical Price Regulation Scheme (PPRS).¹

However, its proposals for introducing a value based system of drug pricing are inherently flawed and, if implemented, would lead to a significant erosion of biopharmaceutical investment in the UK.

It is not true that "other countries have long been able to design, implement, and benefit from effective methods of value based pricing." Many European countries base their pricing on international reference pricing where the UK's pricing has a central role. In Germany therapeutic reference pricing is used, which has led to a severe erosion of that country's research and development over the past decade. In Australia, which has been put forward as an example of value based pricing, the authorities have now decided to overhaul the system and drop their approaches to value based pricing.

What any serious assessment of drug pricing and reimbursement will uncover is that the real problem is in the detail of developing a system of value based pricing and the risk of replacing a system that works with an untried and poorly understood system that effectively undermines the balance between research investment and market return. The UK has traditionally been a leader in research in pharmaceuticals, and great care must be taken to ensure that it is not turned into a follower, which would undermine patients' access to new treatments.

Aisling Burnand chief executive, BioIndustry Association, London SW1X 8PS kmurphy@bioindustry.org

Competing interests: None declared.

- 1 Iheanacho I. Slowly, the monster dies. *BMJ* 2007;335:452. (1 September.)

Author's reply

Burnand gives a predictable, if disappointing, response from an industry that has done very well out of the unequivocally flawed Pharmaceutical

Price Regulation Scheme. In seeking to challenge the basic principles of value based systems, the response attempts to downgrade some instances of where such pricing has provided tangible societal benefits, while completely sidestepping other examples (such as the systems in Canada and Sweden). The dig at the Pharmaceutical Benefits Scheme in Australia is superficial and potentially misleading, and those seeking a more balanced view of the regulation of drug prices in that country (including recent changes) would do well to look elsewhere.¹⁻⁴

Reforming the pricing system in the UK will require constructive contributions from the pharmaceutical industry, for which Burnand's comments are a poor model.

Ike Iheanacho editor, *Drug and Therapeutics Bulletin* BMA House, London WC1H 9JR ike.iheanacho@bmjgroup.com

Competing interests: None declared.

- 1 Australian Government, Department for Health and Ageing, 2006. Strengthening your PBS—preparing for the future www.health.gov.au/internet/wcms/publishing.nsf/Content/A2F23E7630B8F9F3CA25727007F1EC7/\$File/strengthening-your-PBS161106.pdf
- 2 Australian Government, Department for Health and Ageing, 2006. The pharmaceutical benefits scheme (PBS) reform. www.health.gov.au/internet/wcms/publishing.nsf/Content/24693658DD49E286CA2572750081DB74/\$File/PBS%20Reform%202Feb07.pdf
- 3 Searles A, Jefferys S, Doran E, Henry DA. Reference pricing, generic drugs and proposed changes to the pharmaceutical benefits scheme. *MJA* 2007;187:236-9.
- 4 Harvey KJ, Harris AH, Bullone L. The national health amendment (pharmaceutical benefits scheme) bill 2007: reform or fracture? *MJA* 2007;187:206-7.

PPRS is not NICE

Some of Burnand's comments on Iheanacho's article are astonishing.¹ If the mission of any national health system is maximising health, this mission is pursued by assuming that spending on a drug buys not the milligrams of the active substance but the clinical benefit resulting from it. From an ethical point of view, it is bad to use systems not based on clinical benefit (such as the Pharmaceutical Price Regulation Scheme, PPRS).

Outside the UK, many people consider the work done by the National Institute for Health and Clinical Excellence (NICE) in the area of value for money to be an extraordinary example of ethical progress. At last, there is a rationale to support the reimbursement of expensive innovative agents (for example, adjuvant trastuzumab for early breast cancer), as well as a rationale to explain why the reimbursement of poorly innovative agents is subjected to

restrictions (such as anti-Alzheimer drugs).

Drug pricing systems not based on the value for money approach (such as the Italian one) can pay from €0.84 (£0.58; \$1.17) for lansoprazole to €2.32 for esomeprazole, with no specific rationale behind these choices. They can also decide to pay for agents with very poor cost effectiveness profiles.

Andrea Messori coordinator, Laboratory of Pharmacoeconomics, Azienda Careggi, 50134 Florence, Italy
andreamessori@interfree.it

Competing interests: None declared.

- 1 Iheanacho I. Slowly, the monster dies. *BMJ* 2007;335:452. (1 September.)

GOING APE

Unwitting experimentation?

I was intrigued by the choice of antidepressant for the gorilla treated by Pop.¹ Is escitalopram on the zoo's formulary? It is not on our local NHS formulary. Escitalopram is an s-enantiomer of citalopram, which is now off-patent, and a recent *Drug and Therapeutics Bulletin* highlighted how such drugs are developed and marketed after the patent expiry of the originator drug, to extend profits.² In humans, as pointed out by *DTB*, there is little evidence that it is any better than any other serotonin reuptake inhibitors, but it is many times more expensive than generic alternatives. Has the gorilla unwittingly been experimented on in this way as a result of drug company marketing?

Martin Duerden general practitioner, Meddygfa Gyffin, Conwy, North Wales LL32 8LT martin@theduerdens.co.uk

Competing interests: MD is on the editorial board for *Drug and Therapeutics Bulletin*.

- 1 Pop IC. A depressed gorilla. *BMJ* 2007;335:445. (1 September.)
- 2 Anon. New drugs from old? *Drug Therapeutics Bull* 2006;44(10):73-7.

SHOs in the mist?

This filler piece about a senior house officer (SHO) treating a depressed gorilla seems to raise more questions than it answers.¹ Do SHOs in psychiatry have the time to spend two hours assessing a troubled gorilla? Presumably this costs more than peanuts. Was the NHS trust reimbursed for this consultation? Do SHOs have their own secretaries now?

Eamonn J Clarke general practitioner, Wisbech PE14 9BT
eamonn.clarke@nhs.net

Competing interests: None declared.

- 1 Pop IC. A depressed gorilla. *BMJ* 2007;335:445. (1 September.)