A case of fatal bleeding from a jejunal diverticulum, and a second which withstood bowel resection for acute diverticulitis of jejunum and later presented with haematemesis, are described.

The clinical features and findings at follow-up examinations in 90 cases of colonic diverticula are summarized, and one case of acute diverticulitis of caecum is recorded.

The relative incidence of the various complications of colonic diverticula given, and the findings at operation and on later pathological study of material from these cases are described. The principles which should govern management of these complications are discussed.

Diverticulosis has been found to be an important cause of chronic or recurrent dyspepsia. It should be regarded as a disease of the whole length of the intestine although the occurrence of a complication may direct attention mainly to one segment of the bowel.

I am indebted to my colleagues on the staff of the Departments of Pathology and X-ray Diagnosis for their kind co-operation in the investigation of the patients on whom this article is based. Mr. C. R. Strother-Stewart, Mr. J. R. S. Paterson, and Mr. A. R. Kirby were surgical registrars at this hospital during the period covered in this review, and it is a pleasure to record my gratitude for their enthusiasm.

REFERENCES

- Hughes, E. S. R., and Shaw, H. M. (1952). *Med. J. Aust.*, **1**, 259.
Lloyd-Davies, O. V. (1953). *Proc. roy. Soc. Med.*, **46**, 407.
Phillips, J. H. C. (1953). *Brit. J. Surg.*, **40**, 350.

HEPARIN-LIKE ANTICOAGULANT OCCURRING IN ASSOCIATION WITH CHRONIC NEPHRITIS

BY

A. I. CETINGIL, Professor

O. N. ULUTIN, M.D.

AND

M. KARACA, M.D.

From the Second Internal Clinic of Istanbul University,
Guraba Hospital, Istanbul, Turkey

In the case here reported a heparin-like circulating anticoagulant developed during the course of chronic nephritis and caused a haemorrhagic diathesis.

Circulating anticoagulants occur either spontaneously or as a result of transfusions. The presence of antihæmophilic globulin (A.H.G.) was noted in approximately 40 cases of hæmophilia A reported in the literature. A case has been described in Turkey (Frank and Peçikyan, 1954), and at our clinic we have a case of hæmophilia A in which A.H.G. occurred as a result of transfusion (Ulutin, Karaca, and Gürson in press).

Six cases of Christmas disease showing anti-plasma-thromboplastin component (anti-P.T.C.) have been described. Hörder (1955) drew attention to the existence of an inhibitor belonging to factor V. A case of anti-ac-globulin was reported by Ferguson *et al.*

(1957). Alexander (1955). Wagner, Brannan, and Brinkhous (1955), and Jügens (1955) described cases in which anti proconvertinaemia could be demonstrated. Brünimann (1954) described the formation of an antibody against fibrinogen caused by repeated transfusions in a case of congenital afibrinogenaemia.

Search of the literature has brought to light descriptions of five cases of circulating anticoagulant, with haemorrhagic diathesis, in normal subjects without any apparent cause. Anticoagulants have been reported during pregnancy or parturition in 13 cases. Other diseases during the course of which anticoagulants have been reported to occur are syphilis, tuberculosis, rheumatism, myocardial infarction, pemphigus (while being treated with arsenic), and lupus erythematosus. Conley *et al.* (1948) described the occurrence of a circulating anticoagulant, with haemorrhagic diathesis, in the course of chronic nephritis. A similar case is reported below.

Case Report

The patient, a 32-year-old man, had acute nephritis eight years before admission to hospital. For the last two years he complained of headache and pallor of the skin. A week before admission his nose and gums began to bleed. On the day before admission he had a severe headache and his temperature rose to 37.6° C. He was admitted to hospital with the diagnosis of meningitis.

On examination his skin was pale, his gums were bleeding, and there were hæmatomata in the subcutaneous tissues. His blood pressure was 190/120. A systolic murmur was heard in all cardiac areas, and the second aortic sound was accentuated. Nothing abnormal was found in the respiratory, gastro-intestinal, and urogenital systems. Neck stiffness was present, Kernig's sign was positive, and there was some impairment of consciousness. The spinal fluid was blood-stained, confirming the diagnosis of subarachnoid hæmorrhage. The erythrocyte count was 2,800,000, Hb 40%, leucocytes 6,800. The peripheral blood picture was normal. Urine analysis showed albuminuria; the sediment contained abundant erythrocytes and granular casts. Urine specific gravity was 1011. Blood urea was 90 mg./100 ml; urea clearance 40% of normal; glomerular filtrate 35 ml. It appeared that in the course of chronic nephritis a subarachnoid hæmorrhage had occurred.

The bleeding gums and skin hæmatomata drew our attention to the possibility of the existence of a haemorrhagic diathesis. Coagulation studies performed on November 1, 1955, gave the following results: bleeding time, 2½ minutes; coagulation time, 18 minutes (Lee and White method); retraction normal; platelet count, 220,000 (Fonio method); Rosenthal's heparin clotting-time, 120 minutes (normal 30 minutes); prothrombin time (Quick) 18 seconds (normal 12 seconds); antithrombin time (Winterstein) 25 seconds (normal 14 seconds). We found an abnormal result with the thromboplastin generation test of Biggs and Macfarlane and of Douglas (Fig. 1). During the performance of this test, the addition of normal A1 (OH)₃-treated plasma and normal serum or normal platelet suspension to the patient's blood did not produce any increase in the information of thromboplastin. When the patient's A1 (OH)₃-treated plasma or his serum was added to normal blood, the formation of thromboplastin was highly inhibited.

In *in vitro* experiments the addition of protamine sulphate and toluidine blue corrected the prolonged coagulation time and antithrombin time (1% toluidine blue solution or 1% protamine sulphate was added to the patient's blood or plasma in the ratio of 1 to 10). With the protamine titration method of Allen *et al.* (1949) clotting

FIG. 1.—November 1, 1955. 1=0.3 ml. of $\text{Al}(\text{OH})_3$ -treated patient's plasma (1 in 5), 0.3 ml. of patient's serum (1 in 10), and 0.3 ml. of patient's platelet suspension. 2=0.3 ml. of $\text{Al}(\text{OH})_3$ -treated patient's plasma (1 in 5), 0.3 ml. of normal serum (1 in 10), and 0.3 ml. of normal platelet suspension. 3=0.3 ml. of $\text{Al}(\text{OH})_3$ -treated normal plasma (1 in 5), 0.3 ml. of patient's serum (1 in 10), and 0.3 ml. of normal platelet suspension. 4=Normal reaction.

did not occur in the tube containing 0.2 mg. of protamine sulphate (normal clotting, 0.12–0.14 mg.). The administration of protamine sulphate intravenously temporarily corrected the signs of haemorrhagic diathesis.

All these experiments revealed the existence of a "heparin" or "heparin-like" circulating anticoagulant.

A thromboplastin generation test performed a week later (November 8) showed a slight change for the better (Fig. 2): although there was some clinical improvement, the haemorrhagic diathesis still existed. On November 15 the test showed that the formation of thromboplastin had become normal (Fig. 3), and the clinical findings of a haemorrhagic diathesis had disappeared. The patient was discharged at his own request on November 28, 1955.

FIG. 2.—Patient's reaction. November 8.

FIG. 3.—Patient's reaction. November 15.

Discussion

The defect in thromboplastin generation, the prolongation of the coagulation time, heparin clotting-time, prothrombin time, and antithrombin time, and *in vitro* correction of the abnormal findings by the addition of protamine sulphate and toluidine blue, and finally the result of the protamine titration method of Allen *et al.*, show that the circulating anticoagulant in this case is heparin or a heparin-like substance. The fact that one ampoule of protamine sulphate, when injected intravenously, temporarily corrected the prolonged coagulation time and prothrombin time confirmed these findings.

It is well known that heparin activates antithrombin and the clearing factor, and inhibits the conversion of prothrombin to thrombin. Douglas (1956) showed that heparin prevents the utilization of A.H.G. and labile factor, and thus prothrombin is not consumed.

In one of our previous publications (1957) we (Ulutin and Sestakof, 1957) recorded that heparin, when injected intravenously, inhibits the formation of thromboplastin in the thromboplastin generation test of Biggs and Douglas (1957).

Summary

A haemorrhagic diathesis manifesting itself by bleeding from the gums, subcutaneous haematomata, subarachnoid haemorrhage, and epistaxis developed in the course of chronic nephritis. The cause of this haemorrhagic diathesis was a heparin-like circulating anticoagulant. The literature concerning the existence of such an anticoagulant and hyperheparinaemia are briefly reviewed.

REFERENCES

- Alexander, B. (1955). *New Engl. J. Med.*, **252**, 432, 484, 526.
 Allen, J. G., Moulder, P. V., Elghammer, R. M., Grossman, B. J., McKeen, C. L., Sanderson, M., Egner, W., and Crosbie, J. M. (1949). *J. Lab. clin. Med.*, **34**, 473.
 Biggs, R., and Macfarlane, R. G. (1957). *Human Blood Coagulation and its Disorders*, 2nd ed. Blackwell, Oxford.
 Brünimann, R. (1954). *Acta Haemat. (Basel)*, **11**, 40.
 Conley, C. L., Rathbun, H. K., Morse, W. I., and Robinson, J. E. (1948). *Bull. Johns Hopk. Hosp.*, **83**, 288.
 Douglas, A. S. (1956). *J. clin. Invest.*, **35**, 533.
 Ferguson, J. H., Johnston, C. L., and Howell, D. A. (1957). *Proc. Soc. exp. Biol. (N.Y.)*, **95**, 567.
 Frank, E., and Peçikyan, R. (1954). *New Istanbul Contr. clin. Sci.*, **3**, 34.
 Hörder, M. H. (1955). *Acta haemat. (Basel)*, **13**, 235.
 Jürgens, J. (1955). *Ibid.*, **14**, 57.
 Ulutin, O. N., Karaca, M., and Gürson, C., *Türk Tip Cem. Mec.* In press.
 — and Sestakof, D. (1957). *Türk Tip Cem. Mec.*, **23**, 689.
 Wawner, R. H., Brannan, W. M., and Brinkhous, K. M. (1955). *Proc. Soc. exp. Biol. (N.Y.)*, **89**, 266.

IMMEDIATE SENSITIVITY REACTION TO POLIOMYELITIS VACCINE

BY

W. O. THOMSON, M.D., D.P.H., D.I.H., D.P.A.

AND

DUNCAN J. B. FLETCHER, M.B., Ch.B., D.P.H.

Department of Health and Welfare, Glasgow

Untoward reactions to poliomyelitis vaccination are uncommon, and it is generally agreed that poliomyelitis vaccine is exceedingly safe to use. Although the vaccine has been administered many thousands of times, both in this country and the U.S.A., reports of unpleasant sequelae have been few. Nevertheless, from the reports that are available, it is clear that allergic reactions may occur. Usually the skin is involved, but general systemic upsets have also been described. In most cases traces of penicillin in the vaccine have been suspected of being the sensitizing agent.

Most of the reported cases of allergy to poliomyelitis vaccine have occurred in the U.S.A. Zimmerman (1958) described the cases of six patients, all with a previous history of sensitivity to penicillin, who developed urticaria or bullous dermatitis a few days after vaccination. Each patient responded to penicillinase therapy and did not react when a penicillin-free vaccine was subsequently administered.