potassium in four out of five cases, and later those of sodium and chloride.

Possible mechanisms altering the diurnal tides are discussed.

I wish to thank Mr. J. L. MacNaughton for technical assistance.

REFERENCES

- Brown, E. B., Campbell, G. S., Elam, J. O., Gollan, F., Hemingway, A., and Vischer, M. B. (1949). *J. appl. Physiol.*, **1**, 848.
 Evans, B. M., Jones, N. C. H., Milne, M. D., and Steiner, S. (1954). *Clin. Sci.*, **13**, 305.
 Lewis, P. R., and Lobban, M. C. (1957a). *Quart. J. exp. Physiol.*, **42**, 356.
 — (1957b). *Ibid.*, **42**, 371.
 Longson, D., and Mills, J. N. (1953). *J. Physiol. (Lond.)*, **122**, 81.
 Love, A. H. G., Roddie, R. A., Rosensweig, J., and Shanks, R. G. (1957). *Clin. Sci.*, **16**, 281.
 Mills, J. N. (1953). *J. Physiol. (Lond.)*, **122**, 66.
 Norm, M. (1929). *Skand. Arch. Physiol.*, **55**, 184.
 Rosenbaum, J. D., Ferguson, B. C., Davis, R. K., and Rossmoisl, E. C. (1952). *J. clin. Invest.*, **31**, 507.
 Sanderson, P. H. (1952). *Biochem. J.*, **52**, 502.
 Stanbury, S. W., and Thomson, A. E. (1951). *Clin. Sci.*, **10**, 267.
 — (1952). *Ibid.*, **11**, 357.
 Wilkins, R. W., Bradley, S. E., and Friedland, C. K. (1950). *J. clin. Invest.*, **29**, 940.

TUBERCULOSIS CONTROL IN DENMARK

BY

H. FRANKS, M.B., B.S., D.P.H., B.Hyg.

Medical Officer of Health, Chigwell and Waltham Holy Cross Urban District Councils

During a recent visit to Denmark I attempted to assess the significance of the differences between the tuberculosis statistics in that country and those in England and Wales.

The decline of tuberculosis in Denmark has been much faster and has gone much further than in England and Wales. This could be due to differences in (a) population and environment, (b) notification criteria, and (c) the actual measures adopted for controlling the disease in the two countries.

General environmental conditions such as diet, housing, rest, and recreation, are equally good in both countries; the long dark winters of Denmark should outweigh on balance the adverse factor of atmospheric pollution in England.

There have been no tuberculous cattle in Denmark since 1951, and the three-yearly routine tuberculin tests performed on all cattle have revealed only an insignificant number of positive reactors infected from human or possibly avian sources.

Denmark is less industrialized than Britain, and has a population of under 4½m. Except in Copenhagen, there is less opportunity for an infected person to pass on his bacilli at work, at play, or when travelling. The Copenhagen incidence rates are actually almost twice as high as the rates for all Denmark and, as shown in the Table, are more comparable with those for the County of Essex.¹ The rates in England vary locally to an even greater extent,² the extremes in 1956 being Gateshead with 216 and West Suffolk with 22 per 100,000 of the population.

The keeping of a Central Tuberculosis Register in Denmark does much towards preventing duplication. Each notification is checked at the central register against a list of the tuberculous population, and also

New Cases of Respiratory Tuberculosis Notified Annually Per 100,000 of Population

Year	All Denmark	England and Wales	Copenhagen	Essex
1947 ..	85	100	146	98
1948 ..	80	101	154	93
1949 ..	68	102	116	88
1950 ..	58	96	98	87
1951 ..	56	97	82	85
1952 ..	48	95	74	82
1953 ..	40	93	64	72
1954 ..	35	84	66	62
1955 ..	28	76	56	49
1956 ..	25	70	47	49

against a separate list which is maintained of all names that have been discarded from the register on account of cure, emigration, or inability to examine for three successive years. A double-entry system is used to lessen the chance of duplication due to discrepancies in identification particulars.

Relapse is defined as recurrence after freedom from symptoms without treatment for three years or more, and 14% of positive-sputum diagnoses in Denmark in 1955 come into this category. The inadvertent inclusion of such cases in the British incidence figures as new cases could have a considerable effect on the statistics. As the proportion of notifications among the higher age groups increases, so will the chances of duplicate notification increase, since older people are more likely to have had a previous infection, and to have forgotten a notification of long ago, or to have some personal or family reasons for wishing to conceal it.

According to the Danish Tuberculosis Index, less than 3% of tuberculosis deaths had missed notification during life. In England and Wales, however, out of a total of 4,853 deaths from respiratory tuberculosis in 1956, no fewer than 1,244 (26%) are reported as not previously notified.² Might not a number of these cases have been notified in another area or under another (for example, maiden) name during life, thus appearing twice in the incidence statistics?

It would obviously be very difficult to maintain a central tuberculosis register for all England and Wales, because of the large population and incidence. Regionalized checking and recording, however, would be more likely to prevent duplication than our present system of accepting notifications from a very large number of local unrelated sources.

When comparing the total known cases of respiratory tuberculosis in the two countries (in 1955) it is found that in England and Wales they represent 691 per 100,000 of the population,³ but only 6.3% of these were known to be sputum positive at that time; in Denmark the total cases on the registers amounted to only 304.5 per 100,000 of the population, but 25.5% of these were known to be "open" cases, comprising 17.2% with positive sputa and 8.3% with positive stomach washings.

In Denmark about 90% of all notified cases of respiratory tuberculosis are confirmed at some stage by positive bacterial culture, compared with 41% in this country.³ The bacillus is diligently sought in every case, several sputa and stomach washings being cultured, if necessary. In Copenhagen 30% of the new cases are confirmed by culture of stomach washings, these being cases in which no sputum is obtainable. During 1956 a total of 50,000 sputa and 20,000 stomach washings were sent from all parts of Denmark to the Central Laboratory in Copenhagen for culture, and about 10% of these were positive.

Control of Contacts and Cases

In Copenhagen all patients referred by their doctors, or under surveillance as suspects, contacts, or arrested cases, are seen at the Central Tuberculosis Dispensary. Contacts are examined three times during the first year, twice during the second year, and are discharged three years after the last occurrence of open tuberculosis in the household or work place. Dispensary staff are examined every three months. A full-size picture is taken at the first visit, but subsequently, miniature (70 mm.) film is usually employed, and screening is reserved for difficult or urgent investigations in view of the potential risks of excessive irradiation.

The dispensary doctors are provided daily with details of the hospital bed-state, and have the right to admit immediately. The patient remains in the care of the hospital until treatment is completed, and then is returned to the dispensary for supervision, which continues throughout life.

Mantoux tests are done by the nurses, who also take blood for E.S.R. The tuberculosis nurses are all employed full time as such. For children under 12 the jelly patch test is used, followed by a Mantoux test if positive.

Defaulters are visited by the nurse if an open case is involved; otherwise a letter is sent, and repeated after a month, prior to a home visit being made.

When an infectious patient is removed to hospital or dies at home, the room, bedding, and clothes, are disinfected with formalin by the public health department.

Each patient receives a comprehensive 30-page booklet, *What Every Tuberculous Patient should Know*, and is trained in control measures whilst under treatment. If an open case persists in exposing others to undue risk of infection, and refuses admission to hospital, economic assistance from the State can be withheld. This is usually sufficient to induce full co-operation, but as a last resort the medical officer of health may recommend compulsory removal by the police to hospital or institution, and three or four patients are so dealt with each year in Copenhagen. The case does not go to court unless the patient (already in hospital) complains of unjust detention, and the M.O.H. must then be prepared to prove that a real danger to others actually existed. Husband and wife may not, however, be separated under the above provisions.

Centres for Treatment and Control

Sixty hospitals, sanatoria, etc., are available, with beds at the scale of 0.9 per 1,000 of the population. Most of them are administered by local authorities, but a few by the State through the agency of the National Association for the Fight Against Tuberculosis (N.A.F.T.), which corresponds broadly with our N.A.P.T. (now the Chest and Heart Association).

There are also 102 dispensaries, which are administered by local authorities, and in which 869,000 persons—that is, 20% of the whole population—were examined, and 1,336,430 consultations took place in 1955. During the same year 39,337 B.C.G. vaccinations were done; also 1,016 new cases of tuberculosis were discovered, and over 90% of their household contacts were examined. The dispensaries control the distribution of public and voluntary relief.⁴

In most of the 22 counties, the consultant in charge

at the main dispensary is the superintendent of the tuberculosis hospital, and he is also responsible for planning the programme of the mass radiography units throughout the county. This unified control ensures that the special needs of each locality are effectively met and co-ordinated, and eliminates all problems associated with the notification and control of cases discovered by mass radiography.

The fall in the number of cases has led to a great surplus of beds—especially children's beds—in spite of the growing tendency to hold on to non-tuberculous chest cases, which are officially the domain of the general physicians and the general hospitals. Two sanatoria have already been transferred to the Mental Deficiency Hospital Service. It seems probable that the eventual pattern will be a dispensary with a few tuberculosis beds based on each general hospital, and only one or two specializing hospitals retained to serve the whole country for the very chronic homeless and antisocial cases.

Mass Radiography

As in this country, there is a growing feeling in favour of more concentrated attention to selected areas and groups. The areas are, of course, those with a high incidence of tuberculosis; all school staff, Service men, and merchant seamen have annual mass radiography, and in the firms and other organizations which are now well under control after annual radiography for several years attention will in future be concentrated on ensuring x-ray examination of all new employees at the time of their appointment, with less frequent examination of the others. Annual examination of the older age groups is being encouraged more actively, also of persons in close contact with the general public—for example, employees in cafés, bars, cinemas, and theatres, bus conductors, hairdressers, dentists, etc. In England and Wales the standard mortality ratio of the above groups is high,⁵ so that in addition to having much opportunity to spread infection, they have a high capacity to do so. It is intended to advertise the reasons for selecting a particular area or group in the hope of stimulating response.

Up to now it has been the normal Danish practice to carry out a Mantoux test at the same time as mass radiography, but this is being discontinued in areas where most of the population have already become Mantoux-positive.

Approximately 30% of the population of Copenhagen are now under regular annual survey by virtue of their employment with firms and organizations (including the municipality) which have agreed to adopt this scheme as their general policy. It is popular with the workers, who recognize the protection it affords against risk of contact with an infected person.

Positive findings of active tuberculosis occurred in 0.06% of all persons partaking in mass radiography campaigns in Denmark compared with 0.22% in England and Wales; the figure for Essex was 0.14%. Of all newly notified cases of tuberculosis, 29% are discovered by M.M.R. in Denmark and 21% in England and Wales.

The Danish authorities are very actively dealing with the problem of inter-individual and intra-individual inconsistencies of interpreters of x-ray films. In a recent test series, Neilsen,⁶ using, with normal chests standard false or "phantom" infiltrates consisting on

aluminium foil x-rayed on 70 mm. film, found that the average score of 10 independent readers was 67% of the total number seen by one or more of them. The score of individual readers varied between 41 and 81%. Further studies are now proceeding on the cerebral and sensory factors associated with these subjective variations, and on training methods.

The newer mobile units are equipped with 70 mm. cameras, and all 35 mm. apparatus is gradually being replaced. Public sessions are mostly conducted from school premises and village halls, printed invitations having been sent through the post with an explanatory leaflet to every household in the area.

The films are taken to the Central Copenhagen Dispensary for developing, and are read by two doctors in separate rooms. They start at 8 a.m. daily and are not permitted to use the telephone or receive other messages or distractions until the work is completed. If either doctor queries the film, it is an invariable rule for a full-size picture to be taken.

Three or four days after the x-ray examination and Mantoux test the subjects attend again and are told the results. All negative reactors are offered (and accept) vaccination, which is done by the nurses, and a large film is immediately taken if indicated.

The response has been favourable, attendances of between 30 and 80% of those invited having been recorded in different surveys. Since 1946 about one-half of Denmark's population have been examined.

The National Tuberculosis Index is an office which was originally set up to study the future incidence of tuberculosis in classified groups of those who took part in the mass survey of Copenhagen between 1946 and 1948. It has since taken over the statistical analysis of all tuberculosis control measures throughout Denmark. An example of the great value of such an office is the finding that the incidence of tuberculous illness has been significantly higher among unvaccinated persons with a strong Mantoux reaction (12-24 mm.) during a subsequent period of four years' observation, than among all other groups, indicating that they should be frequently and regularly examined.

Prophylaxis

Since 1940 it has been the general policy in Denmark that negative reactors at all ages should be given suspended B.C.G. vaccine. No opportunity of finding non-reactors is missed: infants born in maternity hospitals in Copenhagen (until 1955) were vaccinated before they were 2 weeks old. Many other children are vaccinated at the time of the free medical examination available in the first or second year of life, by their own doctor, under the National Health Service; altogether about 25% of the children in Copenhagen are vaccinated before they enter school. On entry to school (which is at the age of 7 years in Denmark) the jelly patch test is an integral part of the routine medical inspection, and is repeated annually throughout school life in many places, followed by vaccination of non-reactors or revertors. The acceptance rate averages 90%, so that there are very few negative reactors among school children over the age of 8 years. In 1957, 86.6% of all school children in Copenhagen had been vaccinated, and only 8.2% were negative reactors—the rest having converted naturally. As already stated, vaccination is also performed in conjunction with mass radiography.

Up till 1953 B.C.G. was little used in England and Wales, except of contacts and other special groups. The most recent available estimate² of the total so far is 653,000. In Denmark, mass vaccination was introduced about 1936, and by the end of 1955 the total number of vaccinations recorded was 1,121,000,³ and this figure does not include those done privately and in some of the schools. In comparing these figures it must not be forgotten that the Danish population is only one-tenth of ours.

The recommendation to vaccinate all non-reactors has been clear, straightforward, and universal throughout Denmark, without qualifications regarding age or anything else. In many areas consent of parents to the vaccination of children in schools is assumed unless they object in writing as soon as they are informed of the proposed session.

So far as I could ascertain there have been no serious mishaps throughout this long and extensive use of the suspended vaccine. I was impressed by the great care with which it is prepared at the State Serum Institute. Nobody is allowed to enter the laboratory without first washing and changing (in an air-sealed anteroom) into sterile uniform, including cap and mask, from a sealed package. The vaccine is tested for sterility, viability, homogeneity, and pH value before issue, and subsequent confirmatory checks on the potency, safety, and allergizing capacity of each batch are done with guinea-pigs. Its reliability to produce allergy has been repeatedly confirmed, and is endorsed by the findings of the Medical Research Council in this country.

There are no signs of a change over to freeze-dried vaccine. It is held that the process of freeze-drying may kill up to 90% of the bacilli, and that faulty suspension, or subsequent exposure, by untrained field workers under adverse conditions, may do further damage. The suspended vaccine is therefore still preferred for general use because of its well-proved potency and safety, though the value of the dried vaccine as a reserve for emergencies is acknowledged.

As regards technique, the Tuberculosis Research Office of W.H.O. has shown⁷ that the size of the local reaction increases with the depth of the injection; also that a large local reaction is not necessary for a "good take" as measured by the strength of the tuberculin reaction afterwards. Great care is therefore taken to inject as superficially as possible. The indication given is a diameter of not less than 8 mm. for the weal, which would not be achieved with the standard doses of 0.1 ml. if injected into the deep layers of the skin or subcutaneously. The difficulty of making a very superficial injection without loss of time and leakage from the syringe is well understood; on the other hand, a crop of bad local reactions could make B.C.G. unpopular in a district for a long time.

Whether the size of the subsequent tuberculin reaction and its durability are a true measure of the protection afforded is still uncertain. The only reliable test of this must be the observation of disease incidence in vaccinated and control groups over a number of years. This is being studied by the Medical Research Council in this country,⁸ and the incidence rate of primary tuberculosis in the unvaccinated has been about fivefold that in the vaccinated group during the period (two and a half years) of observation so far. More information will be available when morbidity has been

followed up among a parallel series of natural reactors, induced (B.C.G.) reactors, and non-reactors found to be healthy and x-ray-negative during the mass radiography surveys in Copenhagen. It will be some years, however, before enough cases of tuberculosis will have occurred among them to enable any firm conclusions to be drawn from this investigation.

It might be appropriate at this stage to consider the arguments on which B.C.G. vaccination has so far been limited in this country to exposed groups and school leavers—namely, (1) that the risk of infection among children is now negligible; (2) that early identification of infected persons through routine Mantoux-testing (by permitting surveillance and prophylactic treatment with isoniazid) presents a more effective method of control than the extensive use of vaccination; and (3) that vaccination of infants and young children has often resulted in severe local reactions and lymphadenitis.

Despite the above arguments, the risk of children becoming infected rises rapidly at school entry, and the rate of natural Mantoux conversion in this country has been estimated at 4% per year of age.³ In every year since annual mass radiography of school staffs was established in Copenhagen (1949) one or more open cases of tuberculosis have been discovered, and in one year there were as many as seven; in addition, 30 to 40 persons with x-ray changes, requiring surveillance, are discovered among them each year. So long as infection of children can occur in school, or is being imported or met when travelling abroad, there is surely a strong case in favour of using a safe means of protection which is known to be 80% effective. Although most natural converters remain well, the incidence of tuberculous disease among them is nevertheless substantially higher than that among the B.C.G.-converted. Even assuming that reactors can be detected early by routine testing, and that isoniazid prophylaxis is safe and effective, there is no guarantee that the tablets would be swallowed regularly, or that control by this method would be any easier than by mass vaccination. Only in countries where the infection rate among children is known to be high might isoniazid prophylaxis be a suitable alternative to mass vaccination. The Danish authorities are in fact trying this out in Greenland, where over 6% of the population are tuberculous, and most children become tuberculin-positive before it is practicable to offer them vaccination.

The vaccination of newborn infants was suspended in Copenhagen in 1955 because of the frequency of complications. Since then, however, a few cases of tuberculous meningitis and miliary tuberculosis (which had become virtually extinct) have occurred each year, and views are divided on future policy. The Tuberculosis Research Office of W.H.O. has since shown⁴ that infants require only one-half to one-quarter the usual dose of B.C.G. to produce Mantoux conversion, and that such doses are much less likely to produce complications. There is little doubt that experience and skill in preventing small amounts of vaccine from getting under the very fine skin of an infant can also reduce this risk.

Mass Education

Propaganda material is used with discrimination in Denmark and is more or less limited to the actual occasions on which vaccination or mass radiography are

on offer, when it is delivered to each household by post. Nothing is included which might create fear of the consequences of infection. Response is good and suggests that the Danish population as a whole does not need to be constantly reminded that tuberculosis is an infectious disease. This is in contrast with recent findings in London by David and Chalke, where about 30% of people questioned at random did not think that tuberculosis was infectious.

Control of Cattle and Dairy Products

Tuberculosis has been entirely eliminated from cattle in Denmark by a policy of slaughtering all positive reactors and by strict movement control. Studies carried out during the campaign showed that of 106 tuberculous persons who had been working with markedly tuberculous herds, 59.4% were infected with the bovine type of bacillus, but it is reasonable to assume that this source of cattle infection will decrease with every year that goes by.¹⁰

The main features of the campaign were: (1) An adequate well-trained staff of veterinary practitioners to ensure biannual testing, examination, and certification of animals and to enlist the co-operation of the farmers. (2) Enlisting the support of the farmers' organizations on an area basis. (3) Adequate State compensation for the slaughter of positive reactors, with extra inducements for prompt action. (4) Compulsory tests and slaughter of reactors in districts with 90% free herds. (5) Medical examination of all cow-hands working with tuberculosis-free herds. (6) Progressive reduction of the selling price of milk produced on farms where segregation of reactors is not practised. (7) Pasteurization of all milk used for feeding stock.

In the beginning the owners of herds with only a small proportion of reactors often decided to get rid of them at once. In herds with more extensive infection, reactors were usually segregated as found, or an attempt was made in some instances to produce a new tuberculosis-free herd by isolating the calves from the adult herd as soon as they were born.

Towards the end of the campaign, progress was swift because the majority of farmers now had clean herds and put pressure on neighbours with infected animals which might reinfect their own; regulations obliged the owners of infected herds to erect an additional fence 2.5 metres inside their boundary fence to minimize the risk of spread to clean herds.

Conclusions

The incidence of tuberculosis has fallen faster and further in Denmark than in England and Wales. This appears to be due to a policy based on the belief that the disease can be entirely eradicated in the reasonably near future, and differing from our own procedures in the following main respects: (1) the maintenance of a Central Tuberculosis Register; (2) intensive examination of sputa and stomach washings in a specialized central laboratory; (3) intensive use of B.C.G. at all ages and in conjunction with mass radiography; (4) selective use of mass radiography, including annual x-ray examination of all staff employed in schools, public utilities, etc., as part of their terms of contract; (5) invitations to public mass radiography sessions to individual households by post; (6) wider use of powers for compulsory isolation; (7) unified control.

It would be clearly impracticable to maintain a central tuberculosis register and a central tuberculosis laboratory for the whole of England, but it is suggested that a centre incorporating these services could be established for the area of a regional hospital board, together with selective mass radiography and intensive vaccination under a unified control. Consideration should be given to annual x-ray examination as a condition of appointment of school staff and other special categories, and to greater use of existing powers under Section 172 of the Public Health Act, 1936, for the isolation of open cases.

My thanks are due to the Danish Government for a scholarship which enabled me to carry out these studies; to the Essex County Council and the Urban District Councils of Chigwell and Waltham Holy Cross, which, as my employing authorities, generously granted me special leave of absence; and to numerous colleagues, here and in Denmark, who kindly gave me much of their valuable time and facilities for study.

REFERENCES

- ¹ Annual Reports of County Medical Officer, Essex.
- ² Annual Reports of Chief Medical Officer, Ministry of Health.
- ³ Logan, W. P. D., and Benjamin, B. (1957). "Tuberculosis Statistics for England and Wales, 1938-55." *Studies on Medical and Population Subjects*, No. 10. H.M.S.O., London.
- ⁴ Lindhardt, M. (1957). "The Work of the Chest Clinics in Denmark." *Ibid.*, 1956-7.
- ⁵ Neilsen, H. (1957). *Testing and Teaching of Photofluorogram Interpreters*. Munksgaard, Copenhagen.
- ⁶ Medical Report of the National Health Service of Denmark, 1955.
- ⁷ Edwards, L. B., Palmer, C. E., and Magnus, K. (1953). *Wld Hlth Org. Monogr. Ser.*, No. 12.
- ⁸ Tuberculosis Vaccines Clinical Trials Committee (1956). *Brit. med. J.*, 1, 413.
- ⁹ Guld, J., Magnus, K., Tolderlund, K., Biering-Sørensen, K., and Edwards, P. O. (1955). *Brit. med. J.*, 2, 1048.
- ¹⁰ Neilsen, F. W. (1950). *Irish vet. J.*, 4, 178.

A HAZARD IN THE USE OF THE BEAVER PULMONARY VENTILATOR

BY

E. HARVEY FRANKS

Senior Registrar, Department of Anaesthetics,
Royal Infirmary, Cardiff

Recently there have been two potentially dangerous incidents connected with the use of the Beaver ventilator. This communication reports the details and suggests how such incidents may be prevented in the future.

Case Reports

Two patients were suffering from respiratory insufficiency and depression of the laryngeal reflexes. In each case tracheostomy was performed and a cuffed tracheostomy tube inserted. This was connected to a Beaver ventilator in the usual way via the inflating inspiratory-expiratory valve (Mushin, Rendell-Baker, and Thompson, 1959) supplied by the manufacturers.

Progress was satisfactory for several days. Then each patient suddenly became cyanosed immediately following a change of the inspiratory-expiratory valve for cleaning purposes. No cause was apparent, but fortunately the inflating valve was changed for still another. The cyanosis at once disappeared.

Examination of the Valves

In both cases the cyanosis followed immediately after changing the inspiratory-expiratory valve, and on each occasion changing the valve again cured the cyanosis. The suspect valves were carefully dismantled and examined. In both cases the edge of the rubber

FIG. 1.—Obsolete valve. Diaphragm with short studs.

FIG. 2.—Current valve. Diaphragm has long studs.

diaphragm was found to be crinkled and turned over upon itself instead of lying flat. In these circumstances, the flow of air from the ventilator passed straight to the atmosphere and the patient's lungs were not inflated.

With the expert assistance of Mr. E. K. Hillard, senior technician to this Department of Anaesthetics, an examination of several other inspiratory-expiratory valves was made. It was found that two types of valve were in use. Although outwardly nearly identical, the valves differ internally in one vital detail. In one type, now obsolete but still in circulation, the diaphragm has two large diameter short studs which fit correspondingly large shallow depressions in the valve casing. In the other, the diaphragm has smaller diameter long studs which fit correspondingly long grooves in the valve casing (Figs. 1 and 2).