the demonstration that a candidate is a 452 does not add any useful information regarding his abilities. But if his physique is outside the demonstrated distribution of swimmer-canoeists, it seems unlikely that he will be suitable for training. Experienced instructors probably use this knowledge intuitively, but a prospective study of trainees with progress reports might, it is suggested, prove a useful next step.

Summary

A study has been made of the physique of 160 men serving in the Royal Marines. Those in the more hazardous branches are shown to have a more muscular physique. This is considered to be in keeping with Sheldon's thesis that a relation exists between muscularity and somatotonic traits of temperament.

I wish to acknowledge my thanks to the Medical Director-General of the Royal Navy for permission to publish the above paper and to Dr. R. W. Parnell for his help and criticisms during its preparation.

REFERENCES

- Hill, A. Bradford (1955). *Principles of Medical Statistics*, 6th ed. Lancet Ltd., London.
Parnell, R. W. (1954). *Amer. J. phys. Anthropol.*, **12**, 209.
— (1957). *Brit. med. J.*, **1**, 1485.
Sheldon, W. H., and Stevens, S. S. (1942). *The Varieties of Temperament*. Harper, New York.
— and Tucker, W. B. (1940). *The Varieties of Human Physique*. Harper, New York.

REMISSION IN DIABETES

BY

ARNOLD BLOOM, M.D., M.R.C.P.

Physician, Whittington Hospital, London

The tendency to develop diabetes appears to be inherited as a mendelian recessive trait (Joslin, Root, White, and Marble, 1952), but the time of onset and the severity of symptoms vary considerably. In some diabetics, and particularly in the young, there appears to be an absolute failure of insulin production. Plasma studies in these cases show an absence of circulating insulin (Bornstein and Lawrence, 1951), while post-mortem studies on the pancreas fail to demonstrate an insulin content (Wrenshall, Hartroft, and Best, 1954). In diabetics of this type tolbutamide is ineffective, and permanent insulin is necessary to sustain life.

On the other hand, in other diabetics, particularly in obese patients, the circulating insulin appears to be normal, and at post-mortem examination the pancreas can be shown to yield insulin. In such cases the inherited defect may be no more than a functional disability of the pancreas to deal with abnormal stress situations. In potential diabetics this stress is often provided by overeating, but can also be brought about by emotional disturbances, by trauma, by infection, or by pregnancy, for example. Mild diabetes will often remit to an apparently normal state when the stress situation has been removed, as is common experience in obese diabetics following a restricted diet or when an infection has been overcome.

However, some cases of diabetes are neither so severe as to need immediate insulin nor so mild as to respond to simple dietary restriction, and these patients to-day are commonly treated with tolbutamide or similar hypoglycaemic agents. Nevertheless, the decision to put these moderate cases on insulin or tolbutamide is

necessarily an arbitrary one, depending on the severity of the diet and the length of observation. As a rule, weight and symptomatology are the deciding factors. The more overweight the patient is compared with normal for his height and age, the more restricted will the diet be and the longer the period of observation before recourse to adjuvant therapy. On the other hand, where the patient is not overweight, where symptoms are troublesome, and particularly when there is intercurrent infection, the sooner will insulin or tolbutamide be prescribed. The decision to put diabetics on insulin or tolbutamide is not irrevocable. Some patients placed on insulin show a decreasing need for it, so that eventually control is by diet alone. In recent trials of oral hypoglycaemic agents (Crowley, Wolff, and Bloom, 1957) it was found that many newly diagnosed diabetics, not responding to diet alone but responsive when tolbutamide was added, remained normoglycaemic and free from urinary sugar after a six-weeks course of this hypoglycaemic agent. Since then all new diabetics not controlled by diet but responsive to tolbutamide have been given a six-weeks course of these tablets, and the tablets were reintroduced only if relapse of the diabetic state occurred.

The purpose of this communication is to define the proportion of new diabetics (1) who have been controlled by dietary restriction alone from the start, (2) in whom dietary restriction was augmented initially by temporary insulin or tolbutamide, and (3) who need permanent insulin or tolbutamide as well as diet. The dangers of unnecessary prolongation of insulin therapy are stressed.

The patients under survey were those referred to the diabetic unit at Whittington Hospital largely during the period of 1952 to 1957, and all have been under observation for at least six months. In all, 1,125 patients with diabetes were referred during this period, but those who were under treatment when first seen are not included, which leaves 550 newly diagnosed cases for review. There were 344 females and 206 males; 287 (52%) were aged 40 to 65, 82 (15%) were younger than 40, and 181 (33%) were older than 65 at onset. The diagnosis was confirmed by a glucose-tolerance curve in all except severe and acidotic cases where the diagnosis needed no confirmation. 100 g. of glucose was administered to the patient in a fasting state, and venous blood sugars were estimated by Hagedorn and Jensen's method in the fasting state, half an hour, one hour, and two and a half hours after ingestion of glucose.

Initial Treatment

Initial treatment varied according to the type of diabetes present (see Diagram).

1. *Diet Alone*.—Mild and overweight diabetics were instructed concerning a 1,200- or 1,500-calorie diet according to the degree of obesity, and were instructed as out-patients to test the urine for sugar, initially three times daily. Control was regarded as satisfactory when the urine became sugar-free and blood sugars were at a normal or near normal range. As a rule the diet was increased up to 2,000 calories when weight had been lost. Diet alone was also tried in more severe diabetics, usually in the ward, except where insulin was indicated as stated below.

2. *Diet and Tolbutamide*.—Where diet alone failed to relieve symptoms and to control glycosuria and

hyperglycaemia as out-patients, cases were admitted to the wards and observed for a further period of five to seven days in the ward on a measured diet. Where hyperglycaemia persisted, tolbutamide was introduced. The length of out-patient observation varied according to the weight of the patient and degree of hyperglycaemia. In an obese mild diabetic on 1,200 or 1,500 calories, observation would extend for several weeks or months before admittance to the ward. On the other hand, more severe cases, not overweight and with heavy glycosuria, would be admitted to the ward as soon as possible. The diet in this group varied from 1,500 calories (approximately 150 g. of carbohydrate, 65 g. of protein, and 70 g. of fat) to 2,200 calories (approximately 232 g. of carbohydrate, 82 g. of protein, and 105

therapy in the shape of insulin or tolbutamide was necessary. It should be made clear that the number of patients treated with tolbutamide or with insulin bears no relation to the relative place of these two agents in therapy, since tolbutamide has been available for only the last few years of this survey.

Tolbutamide

Seventy-eight diabetics, having failed on simple dietary restriction, were treated successfully with tolbutamide and were maintained on this therapy for six weeks. Of these 78, 25 relapsed with glycosuria and hyperglycaemia after periods varying from a week to six months after the tablets were discontinued. In 18 of these patients tolbutamide was reintroduced and good control again obtained on a maintenance dose of 1 to 2 g. daily in divided doses. In the other seven control was effected with insulin. This means that, of 78 diabetics treated with a six-weeks course of tolbutamide, 53 have so far remained satisfactorily controlled on their dietary regime for periods of from four months to as long as 18 months after the tablets were discontinued. The following are three illustrative cases:

Case 1.—Man aged 57. Diagnosed October, 1956, with symptoms of thirst, polyuria, cramp in the legs, blurring of vision, pruritus penis, and white staining of clothes after micturition. Tolerance curve after 100 g. of glucose: fasting blood sugar, 200 mg./100 ml.; half-hour after, 298 mg.; 1 hour after, 355 mg.; 2½ hours after, 336 mg./100 ml. Height 5 ft. 9 in. (175 cm.). Clothed weight 11 st. 12 lb. (75.3 kg.). Observed for a week in ward on 1,800-calorie diet; blood sugars persistently raised, about 45 g. of sugar daily in urine. Tolbutamide introduced, with good results; blood sugars became normal, urine sugar-free. Tolbutamide maintained at 1 g. b.d. for six weeks and discontinued in December, 1956. Diet increased to 2,000 calories, urine remained sugar-free. Last visit July, 1958, weight 11 st. 3 lb. (71.2 kg.). Mid-afternoon blood sugar 154 mg./100 ml. Feels fit.

Case 2.—Woman aged 65. Diagnosed June, 1957. Thirst, polyuria, lassitude, paraesthesia, pruritus vulvae, and lost about a stone (6.4 kg.) in weight over previous year. Glucose-tolerance curve 129, 306, 350, 407 mg./100 ml. Height 5 ft. 5 in. (165 cm.). Clothed weight 10 st. 13 lb. (69.4 kg.). Observed in ward for one week on 1,800-calorie diet; blood sugars remained raised, daily sugar in urine about 16 g. Tolbutamide introduced successfully and maintained for six weeks. Since then sustained on diet only. June, 1958, weight 10 st. 1 lb. (68.5 kg.), urine sugar-free, mid-afternoon blood sugar 124 mg./100 ml.

Case 3.—Man aged 56. Diagnosed October, 1956. Thirst, polyuria, lassitude, paraesthesia, cramp in legs, change in vision, and loss of 2½ st. (17.5 kg.) in weight over the previous year and a half. Glucose-tolerance curve 197, 326, 424, 333 mg./100 ml. Height 5 ft. 9 in. (175 cm.). Clothed weight 13 st. 3 lb. (83.9 kg.). Observed in ward for nine days on a 1,500-calorie diet. Urine showed acetone (Rothera) and daily sugar about 40 g. Blood sugars remained elevated. Tolbutamide introduced successfully and continued until December, 1956. Since then controlled on 2,000-calorie diet only. June, 1958, weight 12 st. 8 lb. (79.8 kg.), urine sugar-free, blood sugar 111 mg./100 ml. Feels well.

Insulin

Insulin was deemed necessary in 318 newly diagnosed diabetics for the reasons given. Where there was no infection, patients usually spent about five days or so in the ward for the initiation of treatment and for instruction on diet, self-administration of insulin, and urine-testing. During subsequent supervision at the diabetic clinic, insulin dosage was reduced in patients

g. of fat). Where tolbutamide led to good control, cases were maintained on it for six weeks, mainly as out-patients, and it was then discontinued. Usually 1 g. was given twice a day initially, it later being reduced to 0.5 g. three times a day or twice a day. In patients in whom relapse occurred when tolbutamide was discontinued, the drug was usually reintroduced.

3. Diet and Insulin.—Insulin was used in the previous group of diabetics prior to the introduction of the sulphonylureas. Since then insulin has been used in diabetics in whom there was infection or gangrene, in diabetics who failed to respond to tablets, and in severe diabetics whose urine contained acetone. Diet in this group varied according to the patient's size, activity, and age, and lay between 1,500 calories and 2,800 calories (approximately 312 g. of carbohydrate, 102 g. of protein, and 127 g. of fat).

Later supervision at the clinic depended on the type of therapy employed. Under ordinary circumstances those who appeared to be adequately controlled on diet alone were seen two, three, or four times a year, whereas those on insulin were seen each month or each two months, depending on stability. Diabetics on tolbutamide were seen each week initially, and each week after discontinuation of the drug. This interval was lengthened if it became apparent that diet alone was offering good control.

Results

With these criteria, 184 out of 550 new diabetics were adequately controlled by diet from the start and have not relapsed during the period of observation, varying from a year to over six years. Urine has been sugar-free and blood sugars have been normoglycaemic or near normoglycaemic. In the remainder, adjuvant

complaining of hypoglycaemic symptoms or in those where urine tests were always sugar-free and where blood-sugar levels were normoglycaemic or hypoglycaemic. In this way insulin requirements of 154 patients were seen steadily to fall over periods varying from a few weeks to over two years and it was eventually discontinued. Of these 154 patients 62 gradually relapsed while on their dietary regime alone; insulin was restarted in 39, while tolbutamide was successfully introduced in 23. This means that, of 318 diabetics in whom insulin was deemed necessary as initial therapy, it was gradually withdrawn in 92 patients. These have subsequently been controlled on a simple dietary regime, as in the following examples.

Case A.—Woman aged 42. Diabetes diagnosed February, 1953, following a month's history of thirst, polyuria, and severe pruritus vulvae. Had one miscarriage, and one healthy baby, birth weight 8½ lb. (3,855 g.). Patient's mother has diabetes. Clothed weight 10 st. 2 lb. (64.4 kg.). Height 5 ft. 1 in. (155 cm.). Urine contained 2% sugar but no acetone. Glucose-tolerance curve 160, 170, 240, 300 mg./100 ml. Admitted to ward for a week and in view of symptoms placed on diet and given insulin. Discharged on mixture of 20 units soluble and 10 units zinc-protamine insulin. At subsequent attendances at the clinic, urine sugar-free and blood sugars in range 180 to 98 mg./100 ml. Insulin steadily reduced and finally discontinued in February, 1955, two years after it was started. Her weight was 8 st. 8 lb. (54.4 kg.). Since then seen about every four months at clinic; urine free from sugar and an afternoon blood sugar in March, 1958, was 157 mg./100 ml. Feels well in herself and keeps to diet: weight 8 st. 5 lb. (53.1 kg.).

Case B.—Man aged 45. Diabetes diagnosed March, 1951, following a three-months history of thirst, polyuria, lassitude, loss of a stone (6.4 kg.) in weight, and cramp in the legs at night. Father died of diabetes. Weight 12 st. 1 lb. (76.7 kg.), height 5 ft. 7½ in. (171.5 cm.). Urine contained sugar and acetone. Glucose-tolerance curve 158, 241, 270, 254 mg./100 ml. Admitted to ward for instruction on diet (2,000 calories) and urine-testing. After two days' persistent glycosuria, insulin started; good control as out-patient on 16 units soluble plus 8 units zinc-protamine insulin. At subsequent clinic attendances urine sugar-free and blood sugars in range 120 to 90 mg./100 ml. Insulin gradually reduced and finally discontinued in March, 1952, a year after it was started. Weight 12 st. 1 lb. (76.7 kg.). Has attended clinic about every four months since then, urine largely being free from sugar. March, 1958, weight 10 st. 10 lb. (69.9 kg.), afternoon blood sugar 134 mg./100 ml.

Further analysis of these patients in whom insulin was discontinued shows that initial insulin requirements varied from 20 to 46 units, the majority being controlled by 24 to 30 units. 40% have been under observation for more than four years, 16% for more than three years, 27% for more than two years, and 17% for more than a year since discontinuation of insulin. The majority are still under clinic care, but in some the period of observation was terminated by default of attendance or by death. While on insulin, 72% showed no marked change in weight, 18% gained more than half a stone (3.2 kg.), and 10% lost more than half a stone (3.2 kg.). After insulin was discontinued, 63% kept a steady weight, 23% lost more than half a stone (3.2 kg.), while 14% gained more than half a stone (3.2 kg.).

Dangers of Therapy Tolbutamide

Since its introduction, tolbutamide has been extensively used in the treatment of mild diabetics and

so far no serious toxic effects have been reported. Nevertheless, wide experience with the drug does not extend more than a few years and long-term toxic effects may yet appear.

Insulin

Apart from the well-recognized symptoms of acute hypoglycaemia, continued depression of the blood sugar can lead to insidious cerebral changes which may persist even after the blood sugar has returned to normal. These changes are particularly likely to occur in elderly patients and particularly when long-acting insulin is used. Jones (1947) has used the term "post-hypoglycaemic encephalopathy" to describe this condition, and two illustrative case histories follow.

Case I.—Woman aged 63. Diabetes diagnosed eight years previously and patient had self-administered a mixture of 24 units soluble and 24 units zinc-protamine insulin each morning. Had not attended clinic. Husband gave a history of repeated attacks of loss of consciousness, relieved by sugar, and bouts of kicking and screaming over the previous few months. For two days before admission she had been in a state of confusion and had had no insulin on day of admission. On examination was drowsy and disorientated, with spastic limbs and extensor plantar reflexes. Carotid pulses palpable and equal; no papilloedema. B.P. 195/100. Lipodystrophy at injection sites. Eczematous vulvo-vaginitis. Urine: no sugar, no albumin. Blood sugar: 124 mg./100 ml. C.S.F. normal. W.R. negative. Progress: 50 g. sugar intravenously led to definite improvement in her mental state, though this was not sustained. Blood and urinary sugar increased during day and insulin started again next day. Subsequently good control on 20 to 30 units insulin zinc suspension daily. Remained confused, incontinent, and spastic for about 12 days, then distinct improvement. Improvement maintained; left hospital on 20 units insulin zinc suspension subsequently reduced to 10 units in out-patient department. On this dosage urine largely sugar-free and blood sugars at normal or near normal. Mental state largely recovered; walked well, though plantar reflexes remained extensor. *Conclusion:* Post-hypoglycaemic encephalopathy induced by sustained overdosage of long-acting insulin, the cerebral damage persisting even though the blood sugars became normal or raised.

Case II.—Man aged 75. Diabetes diagnosed five years previously and zinc-protamine insulin 22 units daily self-administered since. Urine always blue or green when tested with "clinitest." Lived alone and seemed confused and incontinent for several days before admission, since when no insulin given. Had not attended clinic. On admission B.P. 200/90; conscious but irrational. Rigidity of arms, leg reflexes not exaggerated, plantar reflexes flexor. Urine free from sugar; blood sugar 86 mg./100 ml. Remained confused and restless for about 12 days. Urine sugar-free throughout, despite no insulin for first time in five years; blood sugar on two subsequent occasions 52 and 63 mg./100 ml. Once showed signs of hypoglycaemic coma but regained consciousness after intravenous glucose. After 12 days began to improve and finally left hospital in full possession of mental faculties. Did not require insulin. *Conclusion:* A mild diabetic in whom hypoglycaemic encephalopathy appears to have been induced by unnecessary insulin. Despite cessation of insulin, his mental confusion persisted for nearly two weeks.

Discussion

With few exceptions young underweight diabetics need continuous insulin therapy. Older, milder, and overweight diabetics respond to simple carbohydrate dietary restriction without adjuvant therapy. Between these two groups there are many diabetics whose cases

are neither so severe as to need immediate or permanent insulin nor so mild as to be satisfactorily controlled by dietary restriction alone.

Insulin is indicated in all young and severe diabetics, in diabetics where sepsis or gangrene exists, and, before the days of sulphonylurea therapy, in diabetics who continued to show glycosuria and hyperglycaemia despite adherence to a suitable diet. The initial or hospital dose of insulin in new diabetics is often higher than that subsequently necessary to control the diabetic state. In some diabetics insulin requirements steadily diminish until eventually control is obtained on dietary restriction alone. If these patients are maintained without adequate supervision on the dosage of insulin originally necessary for control, chronic hypoglycaemia may result. This can be particularly disastrous in elderly patients with a cerebral circulation impoverished by arteriosclerotic changes. In elderly patients relapse in the diabetic state can be less dangerous than the daily injection of insulin when the blood sugar is already below normal.

Since the advent of the sulphonylureas, tolbutamide has been used extensively in patients with a mild maturity-onset type of diabetes who have failed to respond to dietary restriction. In conducting trials of oral hypoglycaemic agents it was observed that many newly diagnosed diabetics responsive to tolbutamide did not relapse when therapy was discontinued after six weeks. This routine has been followed subsequently in new diabetics not responding to diet. The diet varied from 1,200 calories in the more obese patients to 2,200 calories in the underweight. The length of observation on diet alone before tablets varied from several months as out-patients in mild obese cases to five days in the ward for severe cases. Although it is always arguable that some of these patients might have eventually attained normoglycaemia after a more restricted diet and prolonged observation, in practice the disadvantages of a short course of tolbutamide are slight so long as proper emphasis is laid on the need for continued dietary restriction. There is evidence in experimental animals that prolonged hyperglycaemia, however produced, can itself cause permanent diabetes, but this can be averted if hyperglycaemia is not sustained (Dohan and Lukens, 1948). If this is true in man, the sooner the blood sugar is restored to normal the better. Patients who relapse when tolbutamide is withdrawn remain responsive to the drug when it is reintroduced, and thereafter treatment can be continuous.

Interval supervision continues to be necessary after insulin or tolbutamide has been discontinued. Good control of the diabetic state requires normoglycaemia or near normoglycaemia, and not merely freedom from symptoms. Hyperglycaemia may carry with it ultimate dangers of degenerative changes in the arteries, kidneys, eyes, and nerves, and these changes have occurred in apparently mild and symptom-free diabetics where hyperglycaemia was allowed to persist. Although it has been shown that remissions may last for many years, relapse may always occur, especially in the face of carelessness in diet or of infection.

Summary

In 550 newly diagnosed diabetics, 184 (33.4%) were adequately controlled by dietary restriction from the start, while a further 145 (26.3%) were ultimately controlled by dietary restriction alone after temporary

treatment with insulin or tolbutamide. The remaining 221 (40.2%) needed permanent insulin or tolbutamide.

The dangers of unnecessary prolongation of insulin therapy are exemplified by two cases of post-hyperglycaemic encephalopathy.

My thanks are due to Miss M. Crowley and her department for the biochemistry results quoted. I am most grateful to Miss G. Weeks for the care she has taken in going through the case records, and to Miss Mason for the preparation of the diagram.

REFERENCES

- Bornstein, J., and Lawrence, R. D. (1951). *Brit. med. J.*, **1**, 732.
Crowley, M. F., Wolff, F. W., and Bloom, A. (1957). *Ibid.*, **2**, 327.
Dohan, F. C., and Lukens, F. D. W. (1948). *Endocrinology*, **42**, 244.
Jones, G. M. (1947). *Amer. J. med. Sci.*, **213**, 206.
Joslin, E. P., Root, H. F., White, P., and Marble, A. (1952). *Treatment of Diabetes Mellitus*, 9th ed., p. 60. Lea and Febiger, Philadelphia.
Wrenshall, G. A., Hartroft, W. S., and Best, C. H. (1954). *Diabetes*, **3**, 444.

ATHEROSCLEROTIC BLOCKAGE IN THE LEFT SUBCLAVIAN ARTERY TREATED BY THROMBO-ENDARTERECTOMY

BY

H. S. SHUCKSMITH, T.D., B.Sc., M.B., F.R.C.S.
Surgeon, General Infirmary and St. James' Hospital, Leeds

Since Takayasu (1908) described blockage of the branches of the aortic arch in Japanese women the condition has been reported as "pulseless disease" (Shimizu and Sano, 1951; Caccamise and Whitman, 1952) and "reversed coarctation" (Frøvig, 1946). The simultaneous blockage of the branches of the aortic arch and of the aortic bifurcation reported by Martorell and Fabré Tersol (1944) is the first record of recognition of occlusion at these two levels in a single patient.

The early papers on Takayasu's syndrome give arteritis, syphilis, and trauma as possible origins of the disease, but later reports suggest that atheroma is the most frequent cause. If this be true, the condition comes into line with the usual pathogenesis of arterial occlusion in other areas. Since the introduction of the technique of aortography by percutaneous puncture of the femoral artery (Seldinger, 1953), demonstration of partial or complete occlusion of one of the branches of the aortic arch is possible before the complete syndrome is established (Wickbom, 1957; Towers *et al.*, 1958).

The basis of this report is atherosclerotic occlusion of the first part of the left subclavian artery in a patient which caused gangrene of the fingers of the left hand and in whom, after confirmation of the site of the block by aortography, thrombo-endarterectomy produced a dramatic improvement in the symptoms and signs.

Case Report

The patient, a woman aged 46, complained of pain in the palm and fingers of the left hand. Examination revealed small areas of fading ecchymoses in the left arm, early gangrenous changes in the left finger-tips (most pronounced in the middle finger), and wasting of the left forearm and small muscles of this hand.

She had had symptoms for 1½–2 years before gangrene of the finger-tips appeared. Small areas of ecchymoses were