BODY-BUILD OF MALE HOMOSEXUALS

BY

A. J. COPPEN, M.D., D.P.M.

Research Assistant, Institute of Psychiatry, London

Studies of the body-build of homosexual patients have produced conflicting reports. Weil (1924), in Germany, reported that homosexuals had a narrower biacromial diameter and a wider intertrochanteric diameter than a control group had. The latter unfortunately was obtained from a different part of the country and measured by a different observer. Wortis (1937) could find no abnormality in the body-build of homosexuals. Sheldon (1949) found there are no distinctive features of homosexual physique recognizable from a somatotyping photograph. He states, however, that the majority of homosexuals are recognizable by what he terms secondary gynandromorphy—that is, “the measures of the femininity of the face, hands, skin, and skin appendages; manner of movement and co-ordination; voice and facial behaviour; in short, the covert gynandromorphy that escapes the photograph.” This is in agreement with the common clinical impression that the effeminate appearance that many male homosexuals show is the result of mannerisms of gait and dress affected consciously or unconsciously by the subject rather than due to any abnormality of body-build.

The present investigation was undertaken to determine whether homosexuals have an abnormal discriminant androgyny score—an index of body-build—which, as is shown below, is related to the sexual development of the individual. The discriminant androgyny score (Tanner, 1951, 1955) of a patient is obtained by measuring the biacromial and bi-iliac diameters and substituting in the following formula:

$$3 \times \text{biacromial} - 1 \times \text{bi-iliac diameters (in cm.)}$$

The term androgyny may be defined as the presence of android features in a woman and gynaecoid features in a man. Tanner obtained a score of 90.1 with a standard deviation of 4.75 for men; the comparable figures for women were 78.9 and 4.57. This sex difference in physique arises at puberty, and before this age the androgyny score fails to discriminate between the sexes.

There is evidence that the androgyny score, or its most important component the biacromial diameter, is correlated with sexual characteristics and abnormalities. Ferriman *et al.* (1957) found that women with hirsuties had an increased urinary 17-ketosteroids output, a tendency to oligomenorrhoea, and an increased biacromial diameter. Raboch (1957) found a decreased biacromial diameter in hypogonadal men and in men with female sex chromatin. Lindegård (1956) showed that size of penis was correlated with the androgyny score. Patients suffering from pre-eclamptic toxæmia and frigidity have been found to have abnormally masculine androgyny scores (Coppén, 1958). Thus there is evidence that subjects who suffer from certain abnormalities related to sexual function will show abnormalities in their androgyny score. The hypothesis tested in the present investigation is that homosexuals have abnormally feminine androgyny scores.

A second androgyny score was used in the present series. This was based on the relative thickness of subcutaneous fat and bone of the calf obtained from a

soft-tissue x-ray film, and will be referred to as the fat-bone index of the calf (Reynolds, 1949; Tanner, 1955). This score is obtained by substituting in the following formula:

$$1 \times \text{bone diameter (mm.)} - 0.4 \times \text{fat diameter (mm.)}$$

The measurements are made at the maximum diameter of the calf. The index differentiates the sexes better than the discriminant androgyny score and misclassifies only 5% of a normal population of men and women (Tanner, 1955). This sex difference also arises at puberty.

The Investigation

Three groups of subjects were investigated: a group of homosexual patients, a control group, and a group of neurotic patients. The control group consisted of the members of a department of a large business organization who were attending for mass radiography. Over 90% of the department attended for the x-ray examination and all of these agreed to be examined for the purpose of the present investigation. The homosexual group consisted of 31 patients who had been either exclusively homosexual or predominantly homosexual, with only occasional heterosexual activity. The patients had attended the Maudsley Hospital primarily for homosexuality; the majority had been referred from the courts after they had been convicted of homosexual offences. Psychiatric symptoms—mainly depression or anxiety—were present to a varying extent in these patients but in none to a severe degree. However, in view of this another control group of 22 heterosexual neurotics was studied, as any differences found in the homosexuals may be related to the differences widely reported in psychiatric patients rather than to their specific sexual abnormalities.

All measurements were performed by myself, using a Martin anthropometer; the patients were measured standing, their shoulders relaxed. The x-ray films of the calf were taken as described by Tanner (1955).

The results are presented in Tables I-IV and the Chart.

TABLE I.—*Body Measurements in the Control, Homosexual, and Neurotic Groups; Comparison of Homosexual and Neurotic Groups With the Control Group*

Group	No.	Mean	S.D.	Compared with Control Group	
				t	P
Control	53	90.3	4.1		
Androgyny score ..		39.8	1.7		
Biacromial diameter (cm.) ..		29.3	1.3		
Bi-iliac diameter (cm.) ..		71.9	7.6		
Weight (kg.) ..		172.6	6.1		
Height (cm.) ..		41.5	10.5		
Age (years) ..	10	35.2	4.4		
Fat-bone index ..					
Homosexuals	31	88.5	5.6	1.68*	0.05
Androgyny score ..		39.1	2.0	1.68*	0.05
Biacromial diameter (cm.) ..		28.8	2.2	0.71	Not significant
Bi-iliac diameter (cm.) ..		73.0	10.8	0.54	
Weight (kg.) ..		172.5	7.9	0.07	
Height (cm.) ..		40.0	9.7	0.65	
Age (years) ..		34.6	3.1	0.47	
Fat-bone index ..					
Neurotic patients ..	22	88.4	4.8	1.74	Not significant
Androgyny score ..		38.9	1.9	2.26	0.05
Biacromial diameter (cm.) ..		28.3	1.0	3.17	0.01
Bi-iliac diameter (cm.) ..		66.7	9.8	2.50	0.02
Weight (kg.) ..		173.6	6.7	0.70	Not significant
Height (cm.) ..		35.1	11.8	2.37	0.05
Age (years) ..					

* One tailed test.

Correlation coefficient between androgyny score and fat-bone index 0.26—not significant.

TABLE II.—*Distribution of Subjects and Mean Androgyny Score According to Social Class*

Social Class	I	II	III	IV	V
Controls ..	(7) 92.6	(0)	(46) 90.1	(0)	(0)
Homosexuals ..	(0)	(8) 89.3	(15) 87.0	(7) 90.7	(1) 89.7

Number of patients in parentheses.

TABLE III.—*Results in Subjects Belonging to Social Class III*

	Mean	S.D.	Compared with Homosexuals	
			t	P
Controls (46):				
Androgyny score ..	90.1	3.2	2.78	0.01
Biacromial diameter (cm.) ..	39.7	1.8	2.17	0.05
Bi-iliac diameter (cm.) ..	29.5	1.6	1.71	Not significant
Age (years) ..	43.1	10.7	0.71	" "
Height (cm.) ..	172.2	5.6	0.36	" "
Weight (kg.) ..	71.5	8.6	0.26	" "
Homosexuals (15):				
Androgyny score ..	87.0	3.1		
Biacromial diameter (cm.) ..	38.5	2.26		
Bi-iliac diameter (cm.) ..	28.6	2.15		
Age (years) ..	40.8	11.7		
Height (cm.) ..	171.6	5.2		
Weight (kg.) ..	71.5	10.7		

TABLE IV.—*Homosexual Patients Divided Into Two Groups of Patients With and Without Other Psychiatric Symptoms*

	Without Other Psychiatric Symptoms (18)		With Other Psychiatric Symptoms (13)	
	Mean	S.D.	Mean	S.D.
Androgyny score ..	88.1	4.9	89.1	3.1
Biacromial diameter (cm.) ..	38.6	1.3	39.6	2.0
Bi-iliac diameter (cm.) ..	27.9	2.0	29.7	2.1
Age (years) ..	42.1	12.1	36.2	10.3
Height (cm.) ..	170.1	5.5	174.8	9.8
Weight (kg.) ..	69.8	10.1	76.9	10.7

Androgyny scores of the subjects in the three groups.

Discriminant Androgyny Score

Homosexuals differ from the controls in their biacromial diameter and androgyny score. There is no difference between the groups so far as the bi-iliac diameter, weight, height, and age are concerned. Because of the differences in body-build between subjects in different social classes that have been reported (Clements and Pickett, 1954; Board of Trade, 1957) both controls and homosexual patients were

divided into five social classes according to the classification of the Registrar-General (General Register Office, 1956). Both the control and the homosexual group have the greatest number of patients in social class III. Table III shows that when patients of this social class are compared with controls of the same class the difference between the group is increased. Moreover, if there are differences due to social class factors between the controls and the homosexual group one would expect to see this reflected in differences in stature between the groups, whereas the mean height of the two groups is almost identical. Indeed, Clements and Pickett (1954) have shown that differences in chest girth between social classes disappeared when standardized for a constant stature.

The homosexual group was divided into two groups, one consisting of patients who complained of some other psychiatric symptoms, the other of patients who appeared free of other symptoms. Table IV shows there was little difference between these two groups; indeed, the patients with no other symptoms differed from the controls more than did the group with other psychiatric symptoms. This suggests that the difference in body-build found in the homosexual group is not related to the presence of these other symptoms.

The neurotic group has an androgyny score and a biacromial diameter very similar to the homosexual group, and in addition has a decreased bi-iliac diameter.

Fat-bone Index

The fat-bone index in the patients and that in the controls are shown in Table I. There is no difference between the homosexuals and the controls. There is no correlation between the fat-bone index and the androgyny score.

Discussion

The results show that homosexuals have a decreased androgyny score and biacromial diameter compared with the control group. This difference, however, is not specific for homosexuality, as the neurotic patients in this study also differ from the controls to approximately the same extent as regards both androgyny and biacromial width. The androgyny score does not discriminate between homosexuals and controls better than does the biacromial diameter, though, as the Chart shows, three homosexual patients have very low androgyny scores, outside the range of the other two groups. It appears, therefore, that homosexuals are similar to people with other psychiatric disorders in having decreased breadth measurements (Rees and Eysenck, 1945), but that their sexual abnormality is not specifically related to these. Rather it seems that the homosexual is influenced by the similar (unknown) factors that produce the abnormalities in body-build found in other psychiatric patients.

The androgyny score is unrelated to the fat-bone index. As both these sexual differences arise at puberty it might have been expected that both would be dependent on the same factors and hence related. However, the androgyny score is dependent on skeletal development alone, whereas the fat-bone index is dependent on soft-tissue development as well as bone.

Summary

The discriminant androgyny score, a measure of physique related to sexual development, was obtained on a group of 31 male homosexuals. The scores of the

homosexuals were significantly lower than those of a control group. A group of heterosexual neurotics, however, showed a similar deviation from the control group. The androgyny score, however, did not differentiate the homosexuals from the controls any better than did the biacromial diameter, which is less in these patients. It was concluded that homosexuals have a body-build similar to that found in patients with other psychiatric disorders and that it could not be specifically related to their sexual abnormality.

I thank Dr. P. D. Scott for permission to examine his patients, Dr. R. D. Hoare and Miss A. Parker for taking the soft-tissue x-ray films, and Dr. H. S. Friend for arranging the examination of controls at Unilever House.

REFERENCES

- Board of Trade (1957). *Women's Measurements and Sizes*. H.M.S.O., London.
 Clements, E. M. B., and Pickett, K. G. (1954). *Brit. J. prev. soc. Med.*, **8**, 108.
 Coppen, A. J. (1958). *J. psychosom. Res.*, **2**, 241.
 Ferriman, P., Thomas, P. K., and Purdie, A. W. (1957). *Brit. med. J.*, **2**, 1410.
 General Register Office (1956). *Census 1951. Classification of Occupations*, p. 3. H.M.S.O., London.
 Lindgård, B. (1956). *Body-build, Body-function, and Personality*. Gleerup, Lund. [*Acta Univ. Lund. N.F. Avd. 2*, **52**, No. 4.]
 Raboch, J. (1957). *J. clin. Endocr.*, **17**, 1429.
 Rees, W. L., and Eysenck, H. J. (1945). *J. ment. Sci.*, **91**, 8.
 Reynolds, E. L. (1949). *Hum. Biol.*, **21**, 199.
 Sheldon, W. H. (1949). *Varieties of Delinquent Youth*. Harper, New York.
 Tanner, J. M. (1951). *Lancet*, **1**, 574.
 — (1955). *Growth at Adolescence*. Blackwell Scientific Publ., Oxford.
 Weil, A. (1924). *Arch. Frauenk. Eugen.*, **10**, 23.
 Wortis, J. (1937). *Amer. J. Psychiat.*, **93**, 1121.

DISTRIBUTION OF ANDROGYN IN MENTAL PATIENTS

BY

J. H. REY, M.D.

Physician, Bethlem Royal and the Maudsley Hospital, London

AND

A. J. COPPEN, M.D., D.P.M.

Research Assistant, Institute of Psychiatry, the Maudsley Hospital, London

The purpose of the present study is to investigate the physique of mental patients by using a measure of the relative maleness and femaleness of body build—the discriminant androgyny score (Tanner, 1951, 1955). Abnormalities in the psychosexual development of mental patients have been postulated and reported by many, and it is of interest to see if such abnormalities are reflected in the development of the corresponding physical endowment of these patients.

The androgyny score is a discriminant function from measurements of the bi-acromial and bi-iliac diameters of groups of normal men and women. The relative proportions of these two diameters have for many years been known to differ in males from those in females. The androgyny score enables the most effective differentiation to be made between the sexes using these measurements; the score is obtained as follows:

$$3 \times \text{bi-acromial} - 1 \times \text{bi-iliac diameter (cm.)}$$

Tanner found this score to have a normal distribution, with a mean and standard deviation of 90.1 and 4.7 for men and 78.9 and 4.6 for women. He found this score misclassified only 12% of the 409 men and women he examined.

The Investigation

Four groups of subjects of each sex were investigated. These were a control group, a group of schizophrenics, and a group with affective disorders; the last group was further divided into a group of patients suffering from a depressive illness and into a group suffering from various neurotic reactions. The male control group consisted of members of a large business organization. The female control group was obtained from an

FIG. 1.—Androgyny scores and bi-acromial diameters in female patients.

FIG. 2.—Androgyny scores and bi-acromial diameters in male patients.

antenatal clinic. None of the measurements used except weight are affected by pregnancy, and in each case the patient's earliest antenatal weight, recorded within the first months of pregnancy, was ascertained. The diagnostic criteria used followed those described by Mayer-Gross *et al.* (1954), with the exception that, for statistical purposes, depressive and neurotic patients were grouped together under the term "affective disorders."

The following measurements were made on each patient: weight, height, bi-acromial diameter, and