became almost normal. Section of this rubbery stuff showed dense mucus which appeared stratified and apparently consisted of a conglomeration of mucoid plugs from a number of bronchi. Entangled in the mucus were many necrotic cells, including some resembling epithelial cells. Culture of a laryngeal swab yielded *H. influenzae*, *Staphylococcus aureus*, and Friedländer's bacillus, but some intratracheal pus produced on culture only *Staphylococcus albus*.

Frequent aspirations were needed through the tracheotomy tube. The temperature remained at about 101° F. (38.3° C.) for the next few days, but the chest remained clear and the respiration rate was not raised. X-ray examination of the chest showed no abnormality. After two days she suddenly became distressed again and all the signs of anoxia returned. The night sister succeeded, after about 15 minutes with the electric suction machine, in bringing out a large blood-stained mixture of mucus and pseudomembrane which had apparently been obstructing the trachea lower down than the end of the tracheotomy tube. After five days in hospital the temperature had settled and phonation was normal. The tracheotomy tube was removed and she remained very well.

A specimen of serum was sent to the Virus Research Laboratory three days after the illness started and again after a fortnight. Complement fixation against influenza A (soluble antigen) showed a rise in titre from 12 to more than 64. Haemagglutination inhibition against influenza A/Singapore/1957 showed a rise in titre from 40 to 240.

Comment

It seemed likely that these patients would not have survived if tracheotomy had been withheld, and the surgeons who performed the operations were in agreement with this view.

Tracheotomy has almost certainly been indicated in cases of obstructive influenza in the past, but these two examples seemed worth recording because of the paucity of previous reports in proved cases of influenza.

Case 1 had subglottic oedema and probably oedema of the tracheal mucosa with copious outpouring of fluid into the trachea. Case 2 by contrast was obstructed by dessicated mucus which adhered to the trachea in thick layers.

I thank Dr. T. Colver for permission to publish these cases under his care, and also Dr. R. N. P. Sutton, of the Virus Research Laboratory, for his serological estimations.

REFERENCES

- Rawlins, G. A. (1957). *Lancet*, 2, 1169.
Roberts, G. B. S. (1957). *Ibid.*, 2, 944.
Stuart-Harris, C. H. (1953). *Influenza and Other Virus Infections of the Respiratory Tract*, pp. 73, 191. Arnold, London.

The value of inquiring into the patient's geographical history is stressed by Professor B. G. MAEGRAITH, Dean of the Liverpool School of Tropical Medicine, in the School's recently published annual report. "So far as I can see," he writes, "one of the bad gaps in the teaching of medical students in this country is the inadequate stress placed on the importance of a patient's geographical history. Failure to understand this may end in tragedy, as it did recently in the Midlands, and very nearly in our own clinic, where a European was admitted suffering from advanced malignant malaria which had been treated elsewhere for a week or more as influenza. In the latter case I found that the patient had told the doctor nothing of his recent visit to the tropics, and, I suspect, the doctor himself may not have thought of asking. One day I hope the importance of the geographical history in medicine will be emphasized at the right level—in undergraduate education."

A CASE OF EARLY YAWS

BY

JACK BARROW, M.B., B.S.

Medical Officer to the Department of Venereal Diseases

H. J. WALLACE, M.D., F.R.C.P.

Physician-in-charge, Department of Dermatology

AND

C. S. NICOL, M.D., M.R.C.P.

*Physician-in-charge, Department of Venereal Diseases
St. Thomas's Hospital, London*

With the immigration of so many West Indian negroes, the diagnosis of latent yaws, based on a history of the disease in childhood and positive serology, is being made more often. Occasional cases of clinical late yaws have also been seen in England (Laird, 1955), and one patient with early (secondary) yaws has been shown at the Section of Dermatology of the Royal Society of Medicine (Erskine, 1956), but clinical details of the case were not published. For these reasons it is felt that the following case may be of particular interest.

Case Report

On January 9, 1958, a negro boy, aged 10, from Dominica, was seen at St. John's Hospital for Diseases of the Skin, and was subsequently admitted to St. Thomas's Hospital.

He had a rash which his mother said had been present for about a month. It involved the scalp, trunk, and upper limbs. The scalp was almost completely covered by heavily crusted purulent lesions and there was a papular rash on the chest and back (see Fig.), these papular lesions also being partially crusted. In addition there were a few

Papular rash on back.

papules on the palms of the hands and one perianal condylomatous lesion but no involvement of mucous membranes. A few small glands were present in both posterior cervical triangles, but none elsewhere. Removal of the crusting from one of the papules on the trunk

revealed a raspberry-like lesion, and dark-ground examination of the exudate disclosed large numbers of treponemata—presumed to be *T. pertenue*.

Serological examination was carried out, and Reiter's protein complement-fixation test (R.P.C.F.) and Price's precipitation reaction (P.P.R.) were both positive. The boy's mother said that he had arrived in this country on November 25, 1957, coming to join his parents, who had been in England for four years. The voyage from Dominica had taken 15 days, and during that time a small ulcer had developed on the outer side of the right ankle. This ulcer had increased in size until treated by the family doctor with penicillin cream, when rapid healing had occurred.

Both parents gave a history of yaws in childhood. They had no signs of syphilis; their standard blood tests were negative, as was the treponemal Wassermann reaction, while the R.P.C.F. was positive in both. The boy had had no sex contact.

On this evidence a diagnosis of secondary yaws was made and a course of penicillin instituted—300,000 units daily for 10 days—and chlortetracycline ointment 1% was applied to the scalp, from which penicillin-resistant and chlortetracycline-sensitive *Staphylococcus aureus* had been cultured. Under this treatment all the lesions disappeared, leaving depigmented areas on the trunk.

A month later the serology was still positive, the P.P.R. only in neat serum. After a further month the P.P.R. had become negative, and the R.P.C.F. alone remained positive.

Summary

A boy aged 10 from Dominica developed an ulcer on his ankle during the voyage to this country. Some weeks later crusted papules appeared on the scalp and trunk, and *T. pertenue* was demonstrated in the serum from one of these lesions. Price's precipitation test and Reiter's protein complement-fixation test were positive in the blood. A diagnosis of early infectious yaws was made; the lesions responded quickly to treatment with penicillin.

REFERENCES

- Erskine, D. (1956). *Proc. roy. Soc. Med.*, **49**, 236.
Laird, S. M. (1955). *Brit. J. vener. Dis.*, **31**, 30.

LONG SURVIVAL IN MITRAL STENOSIS A CASE REPORT

BY

H. R. M. JOHNSON, M.A., M.B., B.Chir.

Junior Assistant Pathologist, University of Cambridge

Mitral stenosis is occasionally found, either clinically or at necropsy, in aged patients. Reports of such findings have been made in recent years (Cabot, 1926; Gelfman, 1943; Cookson, 1949; Howell, 1950; Monroe, 1951; Droller and Pemberton, 1953; Hebbert and Rankin, 1954; Bedford and Caird, 1956), and emphasize the good prognosis which such lesions may carry even though the degree of stenosis appears to be severe. It is possible that atheroma accounts for some of these cases, though in many a clear history of rheumatic fever is obtained.

The following case illustrates long survival in a patient with an anatomically severe degree of mitral stenosis who, furthermore, had borne five children.

Case Report

A housewife aged 69 had rheumatic fever in 1902 at the age of 14. She spent six weeks in bed and subsequently

was not allowed to play games. She married and bore five children, the first in 1910, when aged 22, and the last in 1925, at the age of 37. She remained well until February, 1954, when a left-sided hemiplegia developed from which a good recovery was made. When first seen at Addenbrooke's Hospital in December, 1954, she complained of breathlessness on exertion, headaches, and dizziness of ten months' duration.

On examination mitral systolic and mid-diastolic heart murmurs were heard and auricular fibrillation was present. Her blood pressure was 190/100. She had some residual weakness of her left hand. There were no signs of congestive failure. An electrocardiogram confirmed the auricular fibrillation, and cardioscopy showed a moderate degree of enlargement of the left auricle, right ventricle, and left ventricle with increase in hilar pulsation and pulmonary congestion; the aortic knuckle was small. Her blood haemoglobin was 15.2 g. per 100 ml. and her corrected sedimentation rate 10 mm. in one hour (Wintrobe). A diagnosis of mitral stenosis was made and she was treated with digitalis and a low-salt diet with restricted fluids.

She remained fairly well, but in October, 1956, was readmitted with symptoms and signs of pulmonary infarction. Anticoagulants and penicillin were given and she made a good recovery.

When seen again at the out-patient department in November, 1957, she had signs of congestive failure. These responded initially to diuretics, but she rapidly became worse and was readmitted to hospital, where she died 24 hours later.

Necropsy Findings

Cardiovascular System.—The pericardial sac was adherent over the right ventricle anteriorly and the left atrium posteriorly. Both ventricles were dilated and hypertrophied, particularly the right, the lateral wall of which was up to 1 cm. thick. The mitral valve was stenosed and would not admit the tip of one finger; the cusps were fused and calcified and the chordae tendineae thickened and shortened. The cusps of the aortic valve were not thickened and the commissures were normal; the cusps of the pulmonary and tricuspid valves were smooth and thin. The foramen ovale was closed. Ante-mortem thrombus filled the left atrium, partially blocking the inflow from the pulmonary veins, and extended into the left auricular appendage. Atheroma of the entire pulmonary arterial tree was conspicuous with recent ante-mortem thrombus in the lower main branch of the right pulmonary artery and the upper main branch of the left pulmonary artery. There was a moderate degree of atheroma of the coronary arteries, aorta, and great vessels. No ante-mortem thrombus was present elsewhere in the body.

Lungs.—Bilateral pleural effusions were found—600 ml. of straw-coloured fluid in the right pleural cavity and 250 ml. in the left. There was an area of recent infarction in the postero-inferior part of the left upper lobe, with occlusion by ante-mortem thrombus of the artery supplying it, and an area of old infarction with fibrosis in the right lower lobe.

Other Organs.—Old infarcts were present at the upper poles of both kidneys. There were five areas of softening in the brain; two in the thalamus on each side and one in the right hippocampal gyrus. The liver and spleen were severely congested. The peritoneal cavity contained 500 ml. of straw-coloured fluid.

Comment

Cabot (1926), Gelfman (1943), Baker and Musgrave (1947), Cookson (1949), Wood (1954, 1956), Olesen (1955), and Bourne (1957) have all emphasized the long survival of occasional cases of mitral stenosis without aortic lesions. This case illustrates that point and also the fact that pregnancy need not lead to any worsening of the condition.