REFERENCES

- Abrami, P., Bertrand-Fontaine, Mme, Lichtwitz, A., and Laudat M. (1931). *Bull. Soc. méd. Hôp. Paris*, **47**, 753.
 Bjerneboe, M., Brun, C., Gormsen, H., Iversen, P., and Raaschou, F. (1952). *Acta med. scand.*, **142**, Suppl. 266, p. 249.
 Galán, E., and Masó, C. (1957). *Pediatrics*, **20**, 610.
 Gedda, P. O. (1955). *Acta med. scand.*, **150**, 443.
 Goodman, H. C., and Baxter, J. H. (1957). *J. Amer. med. Ass.*, **165**, 1798.
 Grayzel, H. G., Grayzel, D. M., Heimer, R., and Saremsky, I. (1956). *Exp. Med. Surg.*, **14**, 332.
 Kennedy, W. R. (1935). *Canad. med. Ass. J.*, **33**, 385.
 Lange, K., Strang, R., Slobody, L. B., and Wenk, E. J. (1957). *A.M.A. Arch. intern. Med.*, **99**, 760.
 Latvalahti, J. (1953). *Acta endocr. (Kbh.)*, Suppl. 16.
 Lauson, H. D., Forman, C. W., McNamara, H., Mattar, G., and Barnett, H. L. (1954). *J. clin. Invest.*, **33**, 657.
 McCall, J. W., and Fisher, W. R. (1953). *Ann. Otol. (St. Louis)*, **62**, 316.
 Missen, G. A. K., and Taylor, J. D. (1956). *J. Path. Bact.*, **71**, 179.
 Paulsen, F. (1950). *Acta endocr. (Kbh.)*, **5**, 292.
 Peräsalo, O. (1948). *Ann. Chir. Gynaec. Fenn.*, **37**, Suppl. 2.
 — and Latvalahti, J. (1954). *Acta path. microbiol. scand.*, **34**, 208.
 Richter, G. W. (1954). *Amer. J. Path.*, **30**, 239.
 Rosenblatt, M. B. (1933). *Amer. J. med. Sci.*, **186**, 558.
 — (1936). *Arch. intern. Med.*, **57**, 562.
 Schneekloth, R. E., and Page, I. H. (1955). *Amer. J. med. Sci.*, **230**, 299.
 Sherlock, S. (1945). *Lancet*, **2**, 397.
 Shuttleworth, J. S., and Ross, H. (1956). *Ann. intern. Med.*, **45**, 23.
 Symmers, W. St. C. (1956). *J. clin. Path.*, **9**, 212.
 Trasoff, A., Schneeberg, N., and Scarf, M. (1944). *Arch. intern. Med.*, **74**, 4.
 Waldenström, H. (1928). *Acta chir. scand.*, **63**, 479.
 Walker, G. F. (1928). *Lancet*, **2**, 120.
 West, H. F., and Newns, G. R. (1952). *Ibid.*, **2**, 515.

INTESTINAL PERFORATION AND WIDESPREAD ARTERITIS IN RHEUMATOID ARTHRITIS DURING TREATMENT WITH CORTISONE

BY

R. A. PARKER,* M.D.*Visiting Assistant Professor, Department of Pathology*

AND

PHYLLIS M. THOMAS, M.B., B.Ch., B.Sc.*Beal Research Fellow, Department of Medicine**Stanford University School of Medicine, San Francisco,
U.S.A.*

Perforations of the small and large intestines in patients under treatment with cortisone have not often been reported, and the few cases recorded usually have had coexisting intestinal disease. Sloan, Briggs, and Halsted (1951) report one case of intestinal perforation, Tulin, Kern, and Almy (1952) three cases, Pessel, Beairsto, Edwards, Rathmell, and Wise (1953) one case, all in ulcerative colitis. Levin, Rivo, Scott, Figueroa, Fred, and Barrett (1953) describe a case of perforation in rheumatoid arthritis with widespread arteritis; Sauer, Dearing, and Wollaeger (1953) one case in ulcerative colitis and two in regional enteritis; Bradley (1954) one case of rheumatoid arthritis without arteritis; Palmer, Mason, and Adams (1955) two cases of diverticulitis; and Gumpel, Lewis, and Thomson (1957) one case of ulcerative colitis.

In the present paper three cases of intestinal perforation are described—one case with perforations in the jejunum and duodenum, one case in the caecum, and another in the sigmoid colon. All three were cases

of rheumatoid arthritis under treatment with cortisone, and at necropsy each showed a widespread arteritis.

Case 1

A 54-year-old man was admitted to hospital in May, 1956, for seven weeks for progressive stiffening of joints. In 1948 he developed pulmonary tuberculosis and was treated in a sanatorium until 1952 with resulting arrest of the disease. In January, 1956, he developed pain in the fingers and rheumatoid arthritis was diagnosed. In May, 1956, he was treated with prednisolone, 30 mg. a day. He was readmitted in December, 1956, and on an attempt to reduce the drug the arthritis became worse, so the prednisolone was discontinued. No lupus erythematosus cells were found. He was readmitted in August, 1957, because he had become bedridden, and was found to have the typical deformities of rheumatoid arthritis, with contraction of the hands, elbows, and knees, and many subcutaneous nodules. He was started on cortisone, 150 mg. a day, which was decreased to 70 mg. a day on September 1. On September 7 his third left finger became gangrenous. The cortisone was decreased to 60 mg. a day on September 10. On September 23 he developed severe abdominal pain, and on laparotomy was found to have a recent perforation 4 cm. in diameter of the sigmoid colon. His course was downhill, and after repair of the perforation he died on October 2.

At necropsy generalized peritonitis was found, with 150 ml. of thick yellow fluid in the abdominal cavity. The lungs contained numerous areas of fibrosis. The heart weighed 310 g., no pericarditis being present. The liver was pale and the kidneys appeared normal. The other organs showed no relevant changes. Microscopy revealed an acute arteritis in about half the small arteries in the heart, pancreas, small and large intestines, adrenals, and psoas muscle. The arteritis consisted (Fig. 1) of a deeply staining eosinophilic material (fibrinoid necrosis) extending through part or all of the artery wall and infiltration of its wall with inflammatory cells, mostly polymorphonuclear leucocytes. The heart, pancreas, small and large intestines, adrenals, psoas muscle, kidneys, liver, testes, and oesophagus showed a few small arteries (Fig. 2) with a grossly thickened fibrous intima and partial disruption of the internal elastic lamina; these vessels were thought to be a healed stage of the arteritis. The adrenals were depleted of fat.

Case 2

A 39-year-old man was admitted to hospital with perforation of the caecum in August, 1957. In 1942 he developed pain and stiffness in the joints of his hands and feet. A diagnosis of rheumatoid arthritis was made, and this pursued a chronic course. He was treated with cortisone in 1949 and was given several courses over the next few years, but details and dosage are not known. In 1955 he became short of breath, and was found to be anaemic, with an enlarged spleen. The spleen was removed and a diagnosis of Felty's syndrome was made. Cortisone was again given, and in August, 1957, he developed severe abdominal pain. Laparotomy was performed and a perforation of the caecum with surrounding abscess formation was found. He developed generalized peritonitis, and four days later a resection of part of his small intestine and caecum was performed. After this he was given cortisone, 50 mg. daily. A faecal fistula and subphrenic abscess supervened and he died one month later.

At necropsy there was generalized chronic peritonitis with an abscess measuring 10 cm. in diameter on the posterior aspect of the liver. The heart weighed 340 g. and there was a moderately acute pericarditis. The adrenals were atrophic. The other organs showed no relevant changes. Microscopy revealed an acute arteritis in the small and large intestines, testes, pancreas, kidneys, liver, and adrenals. A healed arteritis was seen in the heart, small and large intestines, pancreas, and kidneys. The adrenals were depleted of fat and atrophic, and about one-quarter of each was necrotic.

*Now at Department of Pathology, University of Cambridge.

Case 3

A 45-year-old man was admitted to hospital on January 21, 1953, for stiffness of finger-joints and redness of knee-joints. In 1936 he developed stiffness of his finger-joints, which was diagnosed as rheumatoid arthritis, and in 1946 he had a duodenal ulcer, both continuing with remissions. On January 12, 1953, he had a haemorrhoidectomy, and on the 17th his joints became stiffer and red. On admission he was found to have redness of both knee-joints, elbows, and second and third finger-joints, and he was placed on cortisone 150 mg. a day. His joints improved and the cortisone was stopped on January 28; however, moderate albuminuria was detected at this time. On February 3 he was put on corticotrophin 20 mg. a day, which was raised to 40 mg. on the 5th. He became unconscious for 12 hours on February 10. On February 15 he developed abdominal pain. Laparotomy revealed a duodenal perforation, which was repaired. He died the next day. At necropsy generalized peritonitis was found with 3,000 ml. of dark brown fluid in the abdominal cavity. A sutured chronic duodenal ulcer and a recent perforation 0.5 cm. across, 60 cm. from the start of the jejunum, was found. The heart weighed 500 g., with left ventricular hypertrophy and pericarditis. The kidneys were deep red in colour, weighed 300 g. each, and were otherwise normal. The other organs showed no relevant changes. Microscopy revealed an acute arteritis in the large and small intestines, testes, adrenals, and kidneys. An area of recent necrosis 2 mm. across, extending throughout the cortex, was found in the right adrenal. Fat was depleted in the adrenals.

Discussion

Three cases of rheumatoid arthritis have been described, each ending fatally with intestinal perforation and a widespread acute arteritis with fibrinoid necrosis. Many arteries also showed marked intimal thickening, which was thought to be a healed stage of the acute arteritis. The intestinal perforation is not thought to be caused by the arteritis, as gastric and duodenal perforations seen during cortisone therapy are usually without an arteritis, and Bradley's (1954) case of perforation of the large intestine occurred without an arteritis in a patient with rheumatoid arthritis on cortisone. However, it is impossible to exclude entirely the arteritis as a cause, because it was found in the intestine in every one of the three present cases.

The causation of the arteritis is also controversial, as arterial lesions have been described as occurring in rheumatoid arthritis for many years. Kemper, Baggenstoss, and Slocumb (1957), reviewing the arterial lesions in rheumatoid arthritis, state that the following types have been described; perivascular accumulation of

cells, vascular infiltration without severe necrosis, and acute arteritis with necrosis of the vessel wall limited to one or two organs—namely, the joints, heart, or subcutaneous tissue. Cruickshank (1954), in a study of 72 necropsy cases, found 11 with a subacute arteritis, sometimes with necrosis; the heart was affected in nine, muscle in three, nerve in five, and trachea, oesophagus, and pancreas in one. A few cases of generalized acute arteritis have been described in which cortisone was not given. Nyström (1953) records one case and Ball (1954) three cases. Therefore it appears that acute arteritis occurs infrequently in rheumatoid arthritis, most often being localized but very occasionally generalized.

However, since the advent of cortisone treatment of rheumatoid arthritis many more cases of generalized acute arteritis have been described; West and Newns (1953) record one case, Levin *et al.* (1953) two cases, Ball (1954) one case, Ogryzlo (1953) an unspecified number, Weinberger (1953) an unspecified number, Slocumb (1953) one case, Finck (1955) one case, and Kemper *et al.* (1957) four cases. Therefore it would seem that widespread acute arteritis can occur in rheumatoid arthritis, but it occurs more often in patients on cortisone. As widespread acute arteritis is uncommon in other diseases treated with cortisone, it appears that patients with rheumatoid arthritis are unduly sensitive to the administration of cortisone. It seems that patients with rheumatoid arthritis have a very slight tendency to develop widespread acute arterial lesions, a tendency which is accentuated by treatment with cortisone.

A possibility to be considered is that the present cases are not of rheumatoid arthritis, but are instances of a disease in which arterial lesions are well recognized and with which arthritis sometimes occurs. The two main possibilities are polyarteritis nodosa and systemic lupus erythematosus; for Friedman, Schwartz, Trubek, and Steinbrocker (1953) have described the occurrence of arthritis in these diseases. However, in the present cases, on careful consideration of their histories and findings, no other diagnosis than rheumatoid arthritis can be made, and the diagnosis of polyarteritis nodosa and disseminated lupus erythematosus is not tenable.

We conclude that the intestinal perforation and arteritis in the present cases are probably separate abnormal reactions to cortisone treatment, and that cases of rheumatoid arthritis are more likely to react abnormally to cortisone therapy than many other diseases.

Summary

Three cases of intestinal perforation and widespread arteritis in rheumatoid arthritis during treatment with cortisone are described. Most of the few cases of intestinal perforation previously described in patients under treatment with cortisone have had intestinal disease.

Widespread acute arteritis is rarely found in rheumatoid arthritis, but since the advent of cortisone therapy such cases appear to have increased. It is suggested that cases of rheumatoid arthritis on cortisone have a greater liability than other diseases to develop an acute arteritis.

We consider that the intestinal perforation and arteritis in the present

FIG. 1

FIG. 2

FIG. 1.—Case 1. Small artery showing acute arteritis with cellular infiltration in its wall. (H. and E. $\times 75$.) FIG. 2.—Case 1. Small artery with marked intimal thickening. (H. and E. $\times 75$.)

cases are probably separate abnormal reactions to cortisone which are more likely to occur in cases of rheumatoid arthritis than in other diseases.

We thank Drs. R. Davison, D. M. Gallagher, and G. B. Robson for permission to use the clinical histories, and Mr. P. Tracy for the photomicrographs.

REFERENCES

- Ball, J. (1954). *Ann. rheum. Dis.*, **13**, 277.
 Bradley, R. L. (1954). *W. Va. med. J.*, **50**, 148.
 Cruickshank, B. (1954). *Ann. rheum. Dis.*, **13**, 136.
 Finck, P. A. (1955). *A.M.A. Arch. Path.*, **60**, 374.
 Friedman, H. H., Schwartz, S., Trubek, M., and Steinbrocker, O. (1953). *Ann. intern. Med.*, **38**, 732.
 Gumpel, R. C., Lewis, E. C., and Thomson, R. V. (1957). *A.M.A. Arch. intern. Med.*, **99**, 660.
 Kemper, J. W., Baggenstoss, A. H., and Slocumb, C. H. (1957). *Ann. intern. Med.*, **46**, 831.
 Levin, M. H., Rivo, J. B., Scott, W., Figueroa, W. G., Fred, L., and Barrett, T. F. (1953). *Amer. J. Med.*, **14**, 265.
 Nyström, G. (1953). *Ann. Med. intern. Fenn.*, **42**, 52.
 Ogryzlo, M. A. (1953). *Ann. rheum. Dis.*, **12**, 323.
 Palmer, T. H., Mason, P. J. H., and Adams, A. C. (1955). *J. Maine med. Ass.*, **46**, 349.
 Pessel, J. F., Beairisto, E. B., Edwards, M. H., Rathmell, T. K., and Wise, J. S. (1953). *Amer. J. dig. Dis.*, **20**, 318.
 Sauer, W. G., Dearing, W. H., and Wollaeger, E. E. (1953). *Proc. Mayo Clin.*, **28**, 641.
 Sloan, S., Briggs, J. D., and Halsted, J. A. (1951). *Gastroenterology*, **18**, 438.
 Slocumb, C. H. (1953). *Ann. rheum. Dis.*, **12**, 323.
 Tulin, M., Kern, F., and Almy, T. P. (1952). *J. Amer. med. Ass.*, **150**, 559.
 Weinberger, H. J. (1953). *Ann. rheum. Dis.*, **12**, 323.
 West, H. F., and Newns, G. R. (1953). *Lancet*, **2**, 1123.

A CONTROLLED TRIAL OF VARIOUS FORMS OF PHYSIOTHERAPY IN ARTHRITIS

BY

D. E. HAMILTON, M.C.S.P.

E. G. L. BYWATERS, M.B., F.R.C.P.

AND

N. W. PLEASE, B.Sc.

From Hammersmith Hospital and the Postgraduate Medical School

In the field of physical medicine, therapeutic measures are critically assessed in comparatively few centres in this country or elsewhere. Much physiotherapy seems to be determined by an authoritarian approach or by superficial observation based on general impressions—for example, that if the patient does well the treatment must therefore be effective. The methods of quantitative assessment seem to be unknown in many non-teaching hospital departments. In so far as a physiotherapeutic department often contains and demands a great deal of expensive equipment, takes up considerable space, and personnel and patient's time, particularly in the fields of orthopaedics and rheumatism, we have felt that a controlled trial of various forms of physiotherapy would not only be useful in itself but be of educational value within the department.

A number of studies have compared the short-term or immediate physiological effects of various forms of physical treatment, but we wished to ascertain the relative value of various forms of physiotherapy as used in the ordinary way in the department over periods of several weeks in the care of patients with chronic arthritis. We were aware of the difficulties in running

a long-term trial, but we did not then realize how formidable they were. It still seems to us, however, that a trial of a realistic nature such as this, conforming to the conditions under which such patients are normally treated, gives information which is of much greater value to the clinician, to the patient, and to the public, than data, more easily arrived at, relating to the immediate physiological effects such as have been published not infrequently in the past.

The questions which the trial was designed to answer were: (1) Which of four forms of physiotherapy (short-wave diathermy, infra-red radiation, faradic stimulation, or paraffin-wax baths) produces the greatest functional improvement during one month's treatment? (2) Is there any significant difference between these treatments and the treatment given to patients on the same basic regime without the additional form of physiotherapy?

Plan of Trial

The trial was set up to study out-patient treatment of individuals with either rheumatoid arthritis involving the hands or knees, or with degenerative joint disease involving the knees. These joints were chosen because they are the easiest on which to assess function accurately. If other joints of these patients were also affected, they were to be treated but not assessed. The criteria for diagnosis were those operative in the rheumatism unit—that is, for rheumatoid arthritis a polyarthritis of unknown aetiology with pain, swelling, and limitation of joints and a raised sedimentation rate; for degenerative joint disease we accepted clinical and radiological evidence with a normal sedimentation rate. It was planned to select 100 patients consecutively referred to the physiotherapy department for treatment and meeting these criteria.

Instead of comparing four separate groups of patients treated in different ways, we treated each patient with each of the four treatments, so that each patient served to some extent as a control on himself. Because, however, it is known that the order of treatments is important and that in general the first treatment of whatever character given to any patient is usually more effective than the second, the order of treatment was allocated at random to each patient. Each patient therefore received one month of treatment A, one month of treatment B, one of C, and one of D, separated each from the other by one week's rest, but randomized in regard to sequence. For each of the three diagnostic categories (rheumatoid arthritis, hands; rheumatoid arthritis, knees; or degenerative joint disease, knees), a series of envelopes were prepared listing the order of these four treatments and sealed. When, therefore, a patient was selected for the trial and met the criteria, the next envelope in the appropriate one of the three categories was opened; this was intended to make sure that over the group as a whole no particular treatment had a greater chance of being first or last than the other treatments. Thus the patients were selected at random to a particular treatment order, but each patient received all four treatments.

The treatments selected for comparison consisted of: (1) Short-wave diathermy given for 20 minutes three times a week. (2) Infra-red radiation for 20 minutes three times a week. (3) In patients with knee involvement, faradism to the quadriceps 20 minutes three times a week, or, for patients with hand involvement, wax treatment 20 minutes three times a