- Frommeyer, W. B., jun., Epstein, R. D., and Taylor, F. H. L. (1950). *Blood*, **5**, 401.
 Hardisty, R. M. (1954). *J. clin. Path.*, **7**, 26.
 Hougie, C. (1953). *Ibid.*, **6**, 30.
 — (1955). *Brit. med. Bull.*, **11**, 16.
 — and Fearnley, M. E. (1954). *Acta haemat. (Basel)*, **12**, 1.
 Joules, H., and MacFarlane, R. G. (1938). *Lancet*, **1**, 715.
 Lozner, E. L., Jolliffe, L. S., and Taylor, F. H. L. (1940). *Amer. J. med. Sci.*, **199**, 318.
 Pons, E. R., jun., and de Torregrosa, M. V. (1952). *Blood*, **7**, 20.
 Richards, M. D., and Spaet, T. H. (1956). *Ibid.*, **11**, 473.
 Soulier, J. P., and Larrieu, M. J. (1953). *J. Lab. clin. Med.*, **41**, 849.
 Van Crefeld, S., Hoorweg, P. G., and Paulssen, M. M. P. (1953). *Blood*, **8**, 125.
 Verstraete, M., and Vandenbroucke, J. (1956). *J. Lab. clin. Med.*, **48**, 673.

CLINICAL METHOD OF ASSESSING TONUS AND VOLUNTARY MOVEMENT IN HEMIPLEGIA*

BY

J. BREWSTER BRENNAN, M.D., D.Phys.Med.
*Department of Physical Medicine, St. James' Hospital,
 London*

During the trial of a new method of spasmolysis in a series of hemiplegic patients difficulty was experienced in finding a method of clinical examination capable of providing an accurate record of the response to treatment. It was desirable to assess its effect upon both tonic spasm and voluntary movement, improvement in the latter being the ultimate aim. Some treatises, such as that by Kendall and Kendall (1949), give excellent guidance on examining individual muscles showing flaccid paresis, but no complete system of examining muscle groups presenting motor deficit and hypertonus has been found, though several workers have made valuable contributions (Teitelbaum and Vynner, 1949; Rinzler *et al.*, 1951; Twitchell, 1951; Hinckley, 1954).

Many difficulties and pitfalls are encountered in elaborating a satisfactory method. Foremost is the influence exerted by certain intrinsic and extrinsic factors upon the tonus and activity of spastic muscle. Among these are temperature, emotion, attitudinal reflexes, and synergy movement. A method of examination which aims to detect spontaneous change, or change in response to treatment, must have a rigid standardization of all such factors during each examination, otherwise the clinical findings may be quite misleading. When this has been done, some means of measuring change in the degree of tonus and in the range and power of voluntary movement is required. Electromyography or mechanical apparatus has usually been used (Berger and Schwartz, 1948; La Joie and Gersten, 1952; Basmajian and Szatmari, 1955), but it appears unsatisfactory that trials of new therapeutic agents of such increasing importance as the spasmolysants should be restricted to the few institutions where such instruments are available. Furthermore, neither apparatus is adequate for examining both of the above modalities.

The following clinical method was devised to study the hypertonic musculature of patients with residual hemiplegia, but it was also found applicable to most other upper motor neurone diseases. It is best described

in two sections: the avoidance of errors which may arise through inadequate control of influencing factors and the method of examination. It is reported in the hope that it may provide a basis for further work.

Control of Factors Influencing Tonus and Activity in Skeletal Muscle

Temperature.—It is thought that external heat facilitates and that cold depresses voluntary muscle action (Schneider and Karpovich, 1948). Again, warming hypertonic muscle reduces the intensity of tonus and cooling increases it. Therefore, to avoid the influence of wide variations in temperature at different examinations all were conducted in a room which was comfortably warm—that is, between 70 and 80° F. (21 and 26.5° C.). More careful control of temperature does not appear to be required, since La Joie and Gersten (1952), studying the effect of heat on hypertonus, found that it was lessened only when the limb was thoroughly warmed by close-range heating.

Emotion.—Bucy (1949) noted that tonic muscle spasm was intensified by emotional stress. In my study it was found that excitement or worry could influence the examination findings to an appreciable extent. Certain steps were therefore taken to avoid strain. These were: exclusion from the examining room of all persons other than the patient and doctor; the patient was settled comfortably; examination was conducted quietly and methodically.

Attitudinal Reflexes.—Several workers have shown that these reflexes exert significant effects upon tone and movement in human hemiplegic patients (Yamshon, Machek, and Covalt, 1949; Twitchell, 1951; La Joie and Gersten, 1952). Their influence cannot be eliminated, but it can be standardized so that the findings are affected to equal degree at successive examinations. The following method was used: the upper limb was examined with the patient sitting upright and looking straight ahead at eye level; the lower limb was examined with the patient lying on the unaffected side, his head supported on a pillow so that it lay in line with the body. These positions were also convenient for conducting the examination.

Flexion and Extension Synergies.—If voluntary movement is not confined to the joint controlled by the muscle groups being tested the influence of synergy movements (Twitchell, 1951) may falsify the findings. The part of the limb (or body) proximal to the joint whose movement was being examined was therefore held immobile.

Volition.—It was observed that voluntary contraction of a hypertonic muscle group often led to a temporary increase in the tone of that group. Tonus was therefore assessed after active movement.

Rate of Passive Movement.—One of the methods used in estimating the intensity of tone was measuring the degree of resistance offered to passive movement. It has been noted (Twitchell, 1951) that the amount of resistance varies directly with the speed at which a hypertonic muscle group is stretched. An attempt was made to standardize this factor by moving the joint through its full range in the space of one second. It was not always possible to do so where hypertonus was pronounced, but only by attempting always to carry the movement through in that time could change in resistance be fairly assessed at different examinations.

*The work presented is part of the material submitted for the M.D. examination of Glasgow University.

Method of Examination

Tonus

Change in the intensity of tonus was measured in three ways, each of which was complementary to the others.

"Range of Normal Tonus."—When any joint in an affected limb was moved passively from one extreme of its range to the opposite it was found that the early section of the total arc in either direction was free of resistance. In the series of residual hemiplegic patients studied the lengths of these arcs, under controlled conditions, remained constant in the absence of treatment. When, with spasmolytic therapy, tonus as measured by other means became less their lengths increased. This phenomenon was also noted by Teitelbaum and Vyner (1949) and Twitchell (1951). The length of the free arc was termed the "range of normal tonus" of the muscle group under observation, and its value was found to be a most reliable index of the degree of spasm. A goniometer was used to measure it.

Degree of Tonus.—In addition to measuring the range of normal tonus, the degree of resistance offered to passive movement through the remainder of the arc was measured. Symbols were used to record it: normal tonus=A, mild=B, moderate=C, marked=D, intense=E. Intermediate values were indicated by coupling letters. It was found that these values altered towards normality at the same time as the range of normal tonus increased. It may be worth recording that clasp-knife rigidity presented in only 70% of the muscle groups observed, the others offering a steady resistance throughout. In order to include both types of response in a standard method the degree of tonus was measured as the amount of resistance encountered in the first 10 to 15 degrees of the resisted arc.

Postural Position of Limb.—This method of measurement was applicable only to the upper limb. Before treatment began all patients held the arm adducted at the central and flexed at the peripheral joints. As tonus decreased a more normal resting posture was assumed. The angle of flexion at the elbow, wrist, and fingers became less acute and the amount of change compared with their original positions was recorded at each examination.

Voluntary Movement

Range.—The range of movement was measured with a goniometer.

Power.—The force of contraction was recorded in the manner recommended by the Medical Research Council. No movement=0, flicker of movement=1, movement with the force of gravity eliminated=2, movement against gravity=3, movement against gravity and resistance=4, normal power=5. Intermediate values were recorded by using the lesser figure followed by a plus sign. It was not uncommonly found that power varied in different sections of the arc through which active movement could be made. For example, a patient might be able to extend his elbow from the position of full flexion through an arc of 45 degrees with a power of 4+ but could only extend with a lessened force, perhaps 3+, through the remainder of the active range. Where such a feature could be measured a record of it was made. As might be expected, the locus at which this lessening of power

became evident was usually close to that which marked the end of the "range of normal tonus."

Examination Procedure

To illustrate how the above system worked in practice the procedure for examining the flexors and extensors of the elbow joint are described. The patient was allowed to sit in a comfortably warm room for some 30 minutes before examination. Clothes above the waist were then removed, and he was seated on an upright chair and asked to look straight ahead. When he was comfortably settled and relaxed the angle of postural flexion at his elbow-joint was measured. Then, with the upper arm held still he was instructed to flex his elbow as far as possible and the range of flexion was measured. Next he extended it to the limit obtainable and the range of extension was noted. These movements were then repeated against resistance and the power gauged. Tonus was then measured, the examiner placing the elbow in the fully flexed position and attempting to put it through its full range of extension in the course of one second. This period of time was measured by saying the words "one million" *sotto voce*. When spoken deliberately they occupy one second. Within a few degrees of resistance becoming evident movement was stopped, the "range of normal flexor tonus" measured, and the degree of resistance recorded. This manoeuvre was repeated three times to obtain an average figure. Tonus in the extensors was measured by reversing the procedure, beginning with the elbow in the fully extended position.

Summary

Examining hypertonic muscle groups presents difficulties which are not encountered in the examination of normal or flaccidly paralysed muscles. Chief of these is the influence upon the clinical findings of certain factors such as temperature, emotion, attitudinal reflexes, and synergy movements. Unless the effect of each of these is strictly and uniformly standardized the examination findings will not be a true reflection of change in tonus or active movement.

A method of clinical examination is described which is thought capable of providing accurate information regarding change, spontaneous or as a response to treatment, in the tonus and contractility of muscle groups presenting tonic spasm and motor deficit.

I thank Dr. P. Bauwens, physician-in-charge of the departments of physical medicine and electrodiagnosis, St. Thomas's Hospital, for his most helpful criticism of the work.

REFERENCES

- Basmajian, J. V., and Szatmari, A. (1955). *A.M.A. Arch. Neurol. Psychiat.*, **73**, 224.
- Berger, F. M., and Schwartz, R. P. (1948). *J. Amer. med. Ass.*, **137**, 772.
- Bucy, P. C. (1949). *The Precentral Motor Cortex*, 2nd ed. Univ. of Illinois Press, Urbana.
- Hinckley, K. F. (1954). *Phys. Ther. Rev.*, **34**, 520.
- Kendall, H. O., and Kendall, F. P. (1949). *Muscles: Testing and Function*. Williams and Wilkins, Baltimore.
- La Joie, W. J., and Gersten, J. W. (1952). *Arch. phys. Med.*, **33**, 595.
- Rinzler, S. H., Brown, H., and Benton, J. G. (1951). *Amer. Heart J.*, **42**, 710.
- Schneider, E. C., and Karpovich, P. V. (1948). *Physiology of Muscular Activity*, 3rd ed. Saunders, Philadelphia.
- Teitelbaum, H. A., and Vyner, H. L. (1949). *Arch. Neurol. Psychiat. (Chicago)*, **62**, 92.
- Twitchell, T. E. (1951). *Brain*, **74**, 443.
- Yamshon, L. J., Machek, O., and Covalt, D. A. (1949). *Arch. phys. Med.*, **30**, 706.