fibrillation (Bourne and Williams, 1948b; Lock and Greiss, 1955). This may hold true also if, after an initial pudendal block with procaine-hyalase-adrenaline mixture, it becomes necessary to administer general anaesthesia in an emergency. Plain procaine was therefore considered to be a safer agent to use in the cases above. An effective pudendal block, achieved in this way, assures one of the co-operation of the mother, who in most instances is thrilled to have actively participated in the birth of her child and is glad to be able to see and hear her baby cry for the very first time.

Summary and Conclusions

A technique for pudendal nerve block anaesthesia for routine use is described.

Cases delivered under pudendal block and those under general anaesthesia are recorded and analysed.

An unconventional use of pethidine hydrochloride and self-administered trichlorethylene as an adjunct to pudendal block is described and discussed.

A critical study of local analgesic agents commonly used for this purpose has been made.

It is concluded that 40 ml. of 1% procaine hydrochloride (plain), injected according to the technique described above, produces a satisfactory pudendal nerve block and perineal analgesia; also that the perineal analgesia so developed, when combined with the use of pethidine and self-administered trilethal analgesia, permits most of the operative vaginal deliveries to be carried out without general anaesthesia.

ADDENDUM.—Since the preparations for this paper were completed, 40 ml. of 1% lignocaine hydrochloride has been substituted in place of procaine. It is perhaps a very premature observation at this stage, but it appears that so far it has lived up to its reputation, and lignocaine, when used with proper caution, may prove to be better than procaine. It is more potent and produces a more prompt and extensive analgesia. It is as safe as procaine if the strength of 1% is not exceeded. While 2% procaine may in some instances be used for nerve blocks, lignocaine in this strength should on no account be used for pudendal block.

I am grateful to Mr. J. N. I. Emblin for his constant guidance, both in the clinical work and in the preparation of this paper, and also for his permission to publish. I am also thankful to Drs. S. K. Ghai, P. K. Irani, and H. Watson for their co-operation and participation; to Miss S. M. Green for making available the obstetric records; and to Sister L. Townsend and her staff for their much-appreciated help in the labour wards.

REFERENCES

- Bourne, A. W., and Williams, L. H. (1948a). *Recent Advances in Obstetrics and Gynaecology*, 7th ed., p. 89. Churchill, London.
- (1948b). *Ibid.*, 7th ed., p. 106.
- (1958). *Ibid.*, 9th ed., p. 187.
- Holland, Sir E., and Bourne, A. (1955). *British Obstetrics and Gynaecological Practice: Obstetrics*, p. 1045. Heinemann, London.
- Kobak, A. J., Evans, E. F., and Johnson, G. R. (1956). *Amer. J. Obstet. Gynec.*, **71**, 981.
- Leshock, J. C. (1957). *Ibid.*, **74**, 1128.
- Lock, F. R., and Greiss, F. C., jun. (1955). *Ibid.*, **70**, 861.
- Micks, R. H. (1957). *The Essentials of Materia Medica, Pharmacology and Therapeutics*, 7th ed., p. 31. Churchill, London.
- Taylor, E. S., Fumetti, H. H. von, Essig, L. L., Goodman, S. N., and Walker, L. C. (1955). *Amer. J. Obstet. Gynec.*, **69**, 348.
- Woolsey, E. L. (1955). *Ibid.*, **69**, 131.

MANIFESTATIONS AND AETIOLOGY OF HYPERTENSION IN THE COLOURED AND BANTU

BY

B. N. FRASER, M.B., M.R.C.P.Ed.

Coronation Hospital, Johannesburg

Hypertension is known to be common in the Bantu in South Africa. A survey was made among both urban and country Bantu by Ordman (1948), who found a significant proportion with a systolic blood pressure of over 150 mm. Hg and a diastolic over 90 in all age groups, though there was a sharp drop in incidence of high blood pressure in males over the age of 60.

Comstock (1957) carried out a survey among both Whites and Negroes in the Southern United States, and his figures showed that the average blood-pressure readings, both systolic and diastolic, are higher in the American adult Negro than his white counterpart. Lennard and Glock (1957), in an appraisal of the distribution of hypertension among Negroes and Whites in the United States, stated that all studies support the conclusion that Negroes more often have essential hypertension, hypertensive heart disease, hypertensive cardiovascular disease or related disorders, than Whites.

Reports from other parts of Africa other than South Africa have revealed variable results. Gelfand (1957) stated that in Rhodesia hypertension, both essential and secondary to chronic nephritis, is common, but he was unable to decide whether it was as frequent as, less frequent, or more frequent than in the white population. He felt comparison between the two races was difficult, since the Europeans' expectation of life is on the whole much higher.

Donnison's (1929) investigation of the blood pressure of males among tribes in the highlands of Kenya showed that up to the fourth decade the average blood pressure was similar to that of European males, but that thereafter there was a tendency to lower pressures. Williams (1944) stated that the majority of cases seen by him in Uganda were secondary to nephritis and urethral stricture. He considered essential hypertension to be rare.

It was decided to investigate hypertension among Bantu and Coloured in-patients as regards incidence, aetiology, morbidity, and mortality. Cases admitted to one of the two medical units at Coronation Hospital during the period January 1, 1955, to December 31, 1956, were used for this purpose. However, owing to the infrequency with which permission was obtained for post-mortem examination, a longer period was taken for investigating necropsy material as regards the incidence of underlying renal disease. The figures of 140 mm. Hg systolic and 90 mm. Hg diastolic were taken as the dividing line between raised and normal blood pressure. This was perhaps not as desirable as relating the blood-pressure reading of each case to a figure taken from a table showing the average blood pressure for that age and sex group, as has been suggested by Pickering (1955). However, no such tables have been compiled for the South African Bantu and Coloured. Comstock's (1957) survey in America has shown that in all age groups the average blood pressure is higher in the American Negro, and the same might well apply to the South African Bantu. Also in many of the Bantu

cases the exact age is uncertain, though this would not necessarily make much difference to a sex-and-age adjusted score. Indian and Chinese cases were excluded from the survey as there were too few to merit inclusion.

Age of Presentation

Table I shows the number of Bantu and Coloured patients admitted during the period of study, with the numbers and percentages of cases presenting with persistent benign hypertension, malignant hypertension, and non-persistent hypertension. It will be seen from the percentage figures that the peak age of presentation of patients with benign hypertension, in both sexes, is over 60, and that there is a tendency for female cases to show themselves at an earlier age than males. Although it is striking that many of the cases present at

TABLE I.—All Bantu and Coloured Cases (January 1, 1955, to December 31, 1956). Deaths to December 31, 1957

Age	No. of Cases	Cases with Persistent Benign Hypertension		Cases with Malignant Hypertension		Cases with Non-Persistent Hypertension		Deaths with Hypertension
		No.	%	No.	%	No.	%	
Males								
Under 20 ..	92	—	—	—	—	—	—	—
20-29 ..	236	2	0.8	1	0.4	2	0.8	—
30-39 ..	331	16	4.8	1	0.3	4	1.2	5
40-49 ..	252	15	6	3	1.2	2	0.8	6
50-59 ..	201	25	12.4	0	—	7	3.5	8
60 and over	172	34	19.8	1	0.6	6	3.5	15
Total ..	1,284	92	7.2	6	0.5	21	1.6	34
Females								
Under 20 ..	94	2	2.1	—	—	—	—	1
20-29 ..	220	5	2.3	—	—	3	1.4	1
30-39 ..	195	21	10.8	3	1.5	6	3.1	5
40-49 ..	174	27	15.5	6	3.4	10	5.7	6
50-59 ..	110	17	15.6	2	1.9	2	1.9	7
60 and over	115	27	23.5	2	1.7	12	10.4	12
Total ..	908	99	10.9	13	1.4	33	3.6	32

an early age, there is no doubt that the shorter longevity of the non-European as compared with the European contributes to the impression that hypertension occurs at an earlier age in the former than in the latter. It will be noted that there are fewer total admissions in the 50-59 and the 60-and-over age groups than in the younger age groups. If more non-Europeans lived to a later age, then more would present in this period of life.

The peak age of presentation of symptoms in Europeans is reported by Søbye (1948) to be 60-64 in men and 65-69 in women with benign hypertension. It is apparent that the European and non-European are alike in presenting late in life. Søbye gives the age group 50-54 as being the main one for presentation of symptoms in both European men and women with malignant hypertension. Our figures are too small for comparison, though malignant hypertension would appear from our few cases to occur at an earlier age in the non-European than benign hypertension.

Presenting Symptoms and Cause of Death

In Table II are shown the main presenting features of the cases with persistent hypertension. It will be seen that the main causes of death are cardiac failure, cerebral accidents, and uraemia in about equal proportions. This contrasts with the findings in European cases, where the figures given by various authors vary but are approximately 50% heart disease,

TABLE II.—Main Presenting Symptoms and Causes of Death in Cases with Persistent Hypertension

	Males		Females		Total	
	No.	Deaths	No.	Deaths	No.	Deaths
Congestive cardiac failure ..	43	12	47	4	90	16
Cerebrovascular symptoms	26	9	30	14	56	23
Renal symptoms and uraemia	17	11	16	8	33	19
Diabetes	3	—	8	—	11	—
Epistaxis	5	—	7	—	12	—

25% cerebrovascular accidents, and 10% uraemia. The lower incidence of deaths due to cardiac failure in the non-European is probably due to the infrequency of coronary artery disease. Coronary artery disease and thrombosis is almost unknown in the Bantu. The incidence is higher in Coloureds, though not nearly as high as in Europeans. Only one of our cases—a Coloured female—had definite myocardial infarction. The higher incidence of uraemia is attributed to the prevalence of primary renal disease.

Sex Incidence and Mortality

Excluding readmissions, 264 cases with hypertension were admitted to the unit during the years 1955 and 1956—in all, 12% of the total number of Bantu and Coloured patients admitted during this period. Of these, 66 patients died before the end of 1957, accounting for just over 25% of all deaths (see Table III). Although far fewer Bantu females were admitted than males, there were slightly more who had hypertension. There are two reasons for the greater number of admissions of Bantu males. A census carried out in 1950 in three areas adjacent to the hospital, from which large numbers of patients are drawn, showed a total population of 66,577 Bantu. Over the age of 14, 54.5% of this population were males and 45.5% females. In addition, the number of medical beds available in the hospital for adult Bantu female patients is half that available for males. Consequently a tighter control is exercised over female medical admissions. The Coloured population in the area of the hospital is

TABLE III.—All Cases, January 1, 1955 to December 31, 1956. (Deaths to December 31, 1957)

	African Males		African Females		Coloured Males		Coloured Females		Total	
	No.	Deaths	No.	Deaths	No.	Deaths	No.	Deaths	No.	Deaths
Cases without hypertension ..	1,019	93	622	65	146	18	131	17	1,928	193 (10%)
Cases with hypertension ..	94	25	105	22	25	9	40	10	264	66 (25%)
All cases ..									2,192	259 (12%)

thought to be more or less equal as regards the sexes and an equal number of medical beds is available for male and female patients, so that the number of admissions of both sexes is about equal. However, far more Coloured females were admitted with hypertension than Coloured males. Although the mortality rate of the non-European female is slightly lower than that of the male, the greater incidence results in at least as many deaths, and the conspicuously better prognosis found in European women with hypertension as compared with European men is certainly not evident in the Bantu and Coloured.

In 54 of the cases the raised blood pressure was not sustained, dropping to normal levels within a period of

a few days to two weeks. The death rate was lower in this group than in the group with persistent hypertension, and it approximated the rate for cases without hypertension. The more severe grades of hypertension were, as to be expected, associated with a higher mortality rate (see Table IV).

TABLE IV.—*Cases of Hypertension According to Their Degree of Severity*

Degree of Severity	No. of Cases	Deaths
Non-persistent	54	6 (11%)
Mild	74	15 (20%)
Moderate	79	17 (22%)
Severe	30	13 (43%)
Gross	8	5 (62%)
Malignant	19	10 (53%)

Mild—General trend of diastolic pressure between 90 and 110 mm. Hg. Moderate—Between 110 and 125 mm. Hg. Severe—Between 125 and 140 mm. Hg. Gross—Over 140 mm. Hg. Malignant—Hypertension + papilloedema.

Aetiology

Apart from the very rare case in which the patient died within a few hours of admission, microscopical examination of the urine was carried out, often on several occasions, as well as frequent testing for albumin and specific gravity. Biochemical examination of the blood—urea, chloride, sodium, potassium, and carbon dioxide combining power—was done as a routine. Where considered necessary, estimation of serum cholesterol and the protein fractions was carried out, as well as any other laboratory investigation that was indicated. In many cases, especially where a kidney lesion was suspected, intravenous pyelography and cystoscopy with retrograde pyelography was undertaken. It is often very difficult to judge whether a case is primarily renal or not. Bechgaard (1946) has pointed out that this is particularly difficult in elderly European women where hypertension, nephrosclerosis, and cystitis often occur together, as elevation of the blood pressure, pyuria, and some derangement of renal function in these patients could easily lead to a false diagnosis of a primary chronic pyelonephritis.

In the non-European group studied, contrast radiography and operative procedures demonstrated two cases of bilateral hydronephrosis, two of renal stone, and one each of double ureter, bilateral bifid pelves, and unilateral hypoplastic kidney. In all, of the 264 cases 70 were diagnosed on radiological, operative, necropsy, and clinical grounds as being primarily renal in origin. This impression that there was a high incidence of underlying renal disease was strengthened by necropsy results. Permission for necropsy was unfortunately obtained in only a minority of patients who were admitted to the unit between January 1, 1951, and December 31, 1956, and who died before January 1, 1958—47 cases in all. Twenty-one cases showed underlying renal disease other than nephrosclerosis. Two cases showed prostatic hypertrophy with bilateral hydronephrosis, and there were two with chronic glomerulonephritis. There was one case of diabetic glomerulosclerosis, and another diabetic revealed glomerulosclerosis and necrotizing papillitis. The remaining 15 cases showed chronic pyelonephritis, in two the process was focal.

Necropsy figures given by other authors have shown a high incidence of underlying renal disease in cases

of hypertension in the non-European in Johannesburg. Becker (1943), Uys (1953), and Levin (1957) quote figures compiled from the necropsy material of the pathology department of the University of the Witwatersrand Medical School. Becker stated that of 247 necropsies done on Coloureds and Bantus with hypertension between 1924 and 1938, 40% were nephrogenous. Uys reviewed 3,707 consecutive Bantu necropsies from 1940 to 1951; of these, 800 had renal disease. There were 76 cases of hypertension nephrosclerosis (23 of these, malignant nephrosclerosis) and 71 cases showed hypertension with underlying kidney disease. Of these 71 cases, 42 showed glomerulonephritis, 18 pyelonephritis, and 8 hydronephrosis. Levin, in a more recent review, stated that of 398 consecutive necropsies in the non-European, 54 showed evidence of hypertension, of which 34 were essential and 18 secondary to chronic pyelonephritis. In none of our cases was an endocrine cause found, though appropriate clinical and laboratory tests were instituted if any indication existed; nor was any found at necropsy. Coarctation of the aorta was not present in any case.

Electrocardiographic Changes

Electrocardiography was performed in the majority of cases. Table V shows the changes found. It will be noted that, even in the milder group, the electrocardiogram often showed changes suggesting left ventricular preponderance. Gillanders (1951) found that the electrocardiogram often showed left ventricular hypertrophy and a horizontal position of the heart in

TABLE V.—*Electrocardiographic Changes*

Group	Total	L.A.D.	V.	H.	L.V.H. and L.V.S.	L.V.H.	L.V.S.	L.B.B.B.	R.B.B.B.	Ab-normal Rhythm
Non-persistent ..	52	35	10	7	11	3	8	3	—	9
Mild ..	69	37	16	16	9	8	9	2	—	10
Moderate to malignant ..	124	89	15	20	38	8	37	4	1	14

L.A.D.—Horizontal heart with left axis deviation in standard leads.

V.—Vertical heart.

H.—Horizontal heart with no left axis deviation in standard leads.

L.V.H.—Left ventricular hypertrophy; where one or more of the following features present: R in lead V_1 or V_2 greater than 25 mm.; S in lead V_1 + R in lead V_2 greater than 35 mm.; R in lead aVL 13 mm. or more tall where lead aVL showed a left ventricular pattern; R in lead aVF 20 mm. or more tall where lead aVF showed a left ventricular pattern.

L.V.S.—Left ventricular strain: T-wave inversion $\pm$ depressed S.T. segments in one or more left ventricular leads.

L.B.B.B.—Left bundle branch block.

R.B.B.B.—Right bundle branch block.

Abnormal rhythm—Auricular extrasystoles 3, ventricular extrasystoles 21, auricular fibrillation 5, auricular flutter 1, auricular flutter-fibrillation 1, transient auriculo-ventricular dissociation 1, paroxysmal ventricular tachycardia 1.

cases of malnutrition heart disease, and this may account in part for the changes found in the cases with mild or non-persistent hypertension, where the raised blood pressure may have played only a minor part in the electrocardiographic changes, especially as many of these cases were in cardiac failure. In Europeans electrocardiographic changes are not a feature of hypertonia (uneventful blood pressure).

Summary

A study among Coloured and Bantu in-patients reveals that hypertension and its complications are a common cause of admission and that these cases carry a high mortality. Females are more often affected than males. Renal disease is a common underlying cause. Compared with the European, hypertension presents at a similar age, but is more serious in the female, and the cause of death is more evenly distributed between heart failure, cerebrovascular accidents, and uraemia.

Endocrine causes are rare. Electrocardiographic changes are found commonly even among the milder cases.

My thanks are due to Dr. G. D. Elliott, medical superintendent of Coronation Hospital, for permission to publish, and to Dr. S. Grieve, senior physician, for the use of clinical material in his wards and for much encouragement and advice. I would also like to thank Professor B. J. P. Becker, of the pathology department of the University of the Witwatersrand Medical School, for permission to publish necropsy findings; and Dr. M. E. Davenport, of the x-ray department of Coronation Hospital, Dr. R. Cassel and Miss J. Horn, of the South African Institute for Medical Research, and Mr. C. D. Kisner, who performed the urological investigations.

REFERENCES

- Bechgaard, P. (1946). *Acta med. scand.*, Suppl. 172.
 Becker, B. J. P. (1943). Cardiovascular Disease in the Bantu and Coloured Races of South Africa. M.D. Thesis, University of the Witwatersrand.
 Comstock, G. W. (1957). *Amer. J. Hyg.*, 65, 271.
 Donnison, C. P. (1929). *Lancet*, 1, 6.
 Gelfand, M. (1957). *The Sick African*, 3rd ed. Juta, Johannesburg.
 Gillanders, A. D. (1951). *Brit. Heart J.*, 13, 177.
 Lennard, H. L., and Glock, C. Y. (1957). *J. chron. Dis.*, 5, 186.
 Levin, N. W. (1957). *Leech (Johannesburg)*, 27, 29.
 Ordman, B. (1948). *Clin. Proc.*, 7, 183.
 Pickering, G. W. (1955). *High Blood Pressure*. Churchill, London.
 Sjøbye, P. (1948). *Heredity in Essential Hypertension and Nephrosclerosis*. Munksgaard, Copenhagen.
 Uys, C. J. (1953). Renal Disease in the Bantu. M.D. Thesis, University of the Witwatersrand.
 Williams, A. W. (1944). *E. Afr. med. J.*, 21, 328.

TREATMENT WITH CORTISONE OF HAEMORRHAGIC DISEASE DUE TO A CIRCULATING ANTICOAGULANT

BY

W. M. R. HENDERSON, M.D.

Pathologist, Pembury Hospital, Tunbridge Wells, Kent

Reports of haemophilia-like disease due to circulating anticoagulants are still comparatively rare, though they have become more frequent since the introduction of the thromboplastin generation and other sensitive tests which help to distinguish these cases from true haemophilia or Christmas disease. Two cases have, however, been seen in this hospital within a period of three years. The first, in an old lady with rheumatoid arthritis, has already been reported by Hardisty (1954).

Reviewing the literature since Lozner *et al.* (1940) first reported a case, Verstraete and Vandenbroucke (1956) were able to find 61 cases. Of these, 20 were in patients with haemophilia or Christmas disease who had developed the anticoagulant following blood transfusions; 16 were in women within one year of a pregnancy; one was in a newborn infant; eight were in females unrelated to pregnancy; and 16 in non-haemophilic males. Most of the latter cases were associated with one or other of a variety of constitutional diseases, a list of which is given by Hougie (1955), but in a few no associated disease could be found (Joules and MacFarlane, 1938; Pons and de Torregrosa, 1952; Hougie, 1953; and others). The case reported here comes into the latter group.

Case History

A gardener aged 56 was sent into hospital by his doctor with a history of having spontaneously started passing blood-stained urine five weeks previously. At times it had been profuse, and small clots had been present. There was no frequency or dysuria. In the last two weeks he had also noticed the appearance of spontaneous bruising of his skin and had bled more readily after shaving. He also had palpitations on exercise and cramp-like substernal pain.

In the past he had had "black measles" during the 1914-1918 war and also nephritis, for which he was in hospital three months. He had never had any bleeding episodes or bruising previously. There was no family history of any bleeding disease.

On admission on April 8, 1955, he was very pale and slightly jaundiced. There were large bruises on both his arms. Petechiae were not seen in the skin. His pulse was 100 and blood pressure 120/70. A systolic murmur was present at the apex and there was slight oedema of the right ankle. Neither the liver nor the spleen was palpable, but the right kidney could be felt though it was not tender.

The urine on admission contained fresh blood, and microscopy showed red cells + + + + and occasional leucocytes. Culture grew *Escherichia coli*.

Cystoscopy revealed spurts of blood coming from the right ureteric orifice. Retrograde pyelography showed an irregular caliceal pattern on the right, with the ureter dilated and tortuous; it was supposed that this was due either to a new growth or to the presence of blood clots. Preliminary laboratory investigation showed: Hb 56% (8.3 g./100 ml.); red blood cells 2,990,000/c.mm.; C.I. 0.94; white cells 10,400 (neutrophils 79%, monocytes 10%, lymphocytes 1%); platelets 425,000/c.mm.; serum bilirubin 1.15 mg./100 ml.; blood urea 70 mg./100 ml. The blood group was A rhesus positive. It was noticed that the blood sent to the laboratory for grouping and cross-matching had not clotted after four hours, so further investigations were carried out.

Laboratory Investigations.—E.S.R. (Westergren) 62 mm. at first hour, 100 mm. at second hour. W.R. and Kahn tests were negative. Examination for cold agglutinins was negative. Agglutination against *Salmonella* and *Brucella* organisms was negative. Serum proteins 6.08 g./100 ml. (albumin 3.44 g., globulin 2.64 g.). A/G ratio 1.3:1. Electrophoresis of plasma showed a slight increase in the α_2 globulin, but was otherwise normal. Fibrinogen 650 mg./100 ml. Thymol turbidity and alkaline phosphatase were normal. Blood culture grew *E. coli* from the bile salt broth. The direct Coombs test was negative.

Investigation of Coagulation.—Clotting-time (Lee and White), 70 minutes. Prothrombin time, 15 seconds (control 15 seconds). Bleeding-time (Duke) normal. Clotting-time of recalcified plasma 430 seconds (control 160 seconds). Mixing-tests with patient's plasma and normal plasma showed that the patient's plasma, when mixed with normal plasma, had the power of prolonging the clotting-time of the latter (Table I). A modified prothrombin consumption test was abnormal.

TABLE I.—Clotting-times of Mixture of a Normal and Patient's Plasma on Admission

Source of Plasma	Quantities of Plasma (ml.) Used				
	0.1	0.1	0.1	0.05	0.01
Patient	—	—	—	—	—
Normal	0.1	0.1	0.1	0.1	0.1
Saline	0.1	0.1	—	0.05	0.09
M/40 CaCl ₂	0.2	0.2	0.2	0.2	0.2
Clotting-time in seconds	430	160	255	170	155

The results suggested the presence of a circulating anticoagulant in his blood, and on April 15 samples of plasma and serum were sent to Dr. W. R. Pitney, of the department of haematology, Postgraduate Medical School, Hammer-smith, who kindly performed a thromboplastin generation