Drug Treatment of Disease

MYASTHENIA GRAVIS

BY

J. W. ALDREN TURNER, D.M., F.R.C.P.*Neurologist, St. Bartholomew's Hospital*

Effective drug treatment of myasthenia gravis was introduced in 1934 when Dr. Mary Walker discovered that an injection of physostigmine would relieve the muscular weakness of the disease. In the following year she found that neostigmine had a similar action, and as it is more effective and less apt to cause side-effects it is now the standard drug used in treatment. In myasthenia gravis there are no recognized histological abnormalities in the central nervous system, the peripheral nerves, or the skeletal muscles, with the possible exception of slight changes at the motor end plates. The disorder is thought to be due to a disturbance of conduction at the neuromuscular junction, the exact nature of this being still undecided. Neostigmine is an inhibitor of the enzyme choline esterase, which inactivates acetylcholine, and it is partly owing to this action that the drug improves muscular power in myasthenia gravis, but there is also some evidence that it exerts a direct action on voluntary muscle fibres.

Neostigmine

Neostigmine can be given by the mouth as neostigmine bromide, each tablet containing 15 mg., or by subcutaneous injection as neostigmine methylsulphate in doses of 0.5 to 2.5 mg.; 15 mg. by the mouth is approximately equivalent to 0.5 mg. by injection.

Oral Treatment with Neostigmine.—Most patients on neostigmine receive it by mouth. The dosage varies considerably in accordance with the severity of the disease, and often it may have to be varied from day to day, this depending on the patient's activities. Oral neostigmine increases power in about thirty minutes, and the effect usually lasts from three to four hours. Doses therefore have to be given at regular intervals of about three hours, though some patients find it more satisfactory to have two-hourly or even hourly doses. Many myasthenic patients have difficulty in swallowing and sometimes in chewing, and for this reason doses are best given about half an hour before meals; even for patients without bulbar disease this is the best procedure, because neostigmine has a greater effect when taken before meals. If it causes nausea, some milk or a biscuit can be taken at the same time. Because of the short action of the drug a dose should be taken last thing at night and immediately on waking (when muscle weakness is often marked), and in severe cases a dose will have to be taken during the night.

In starting treatment of a patient it is usual to begin with a small dose such as 15 mg. three- or four-hourly during the day, and by trial adjust the amount used and the timing of the drug to get the maximum benefit without causing serious side-effects. Co-operation from the patient is most important in achieving this. The total dosage varies greatly, so much as 1,000–1,200 mg. in 24 hours being needed for short periods in some patients.

Neostigmine by Injection.—An injection of neostigmine is of great value as a diagnostic test in

myasthenia gravis, 1.5 mg. together with atropine 1/100 gr. (0.65 mg.) being given subcutaneously. In myasthenia there is considerable and sometimes dramatic improvement in power. The effects last for about three hours. This marked increase in power after a neostigmine injection does not occur in other conditions causing weakness of muscle, though slight improvement may be seen in a variety of diseases. In some cases of myasthenia gravis, particularly those of long standing, the weakness may not be completely relieved; this is especially true of the external ocular muscles, a fact which must be realized in assessing the results of a diagnostic injection of neostigmine. In normal people, and those with other diseases, 1.5 mg. of neostigmine by injection may cause widespread muscle fasciculation, which does not occur in patients with myasthenia.

Injections of neostigmine are not needed in the treatment of most myasthenic patients, but they may be useful before an exceptional increase of activity or during an acute infection, such as pneumonia, which often aggravates myasthenia. In a myasthenic crisis or immediately after thymectomy, neostigmine should be given by injection, and to secure a continuous effect it can be given in a slow intravenous drip.

Side-effects of Neostigmine.—Neostigmine may produce side-effects. By its muscarinic action it causes contraction of the smooth muscles of the alimentary tract, resulting in colicky abdominal action, pain, nausea, vomiting, and diarrhoea, and by its similar action on glands produces salivation, lacrimation, sweating, and increase of bronchial secretions. In some myasthenic patients, particularly in the early stages of treatment, abdominal pain, nausea, and diarrhoea occur, and these troublesome symptoms can usually be controlled by the oral administration of tinct. belladonnae, 10–15 minims (0.6–0.9 ml.) or ext. belladonnae sic., $\frac{1}{4}$ gr. (32 mg.) three times a day, or in my experience preferably by propantheline bromide, 15 mg. two or three times a day. When neostigmine is given by injection it is advisable to give atropine also in a dose of 1/100 gr. (0.65 mg.), the frequency of injections being adjusted to prevent the abdominal symptoms.

A more serious effect which occasionally occurs in myasthenia when large doses of neostigmine are given is due to the nicotinic action of the drug, which may cause widespread fasciculation and increased weakness of voluntary muscles. This condition, sometimes called a cholinergic crisis (to distinguish it from a myasthenic crisis), can usually be recognized because the increased weakness comes on within 15 to 45 minutes of administration of neostigmine and is accompanied by muscle fasciculation, salivation, increase of bronchial secretions, small pupils, and frequently coldness and clamminess.

Some workers have found the "tensilon" test valuable in distinguishing between a cholinergic and a myasthenic crisis. Tensilon (edrophonium chloride) is a

very short-acting anticholinesterase drug; when it is given intravenously, myasthenic weakness improves in 30–40 seconds and the increase in strength lasts about five minutes. However, when it is injected into a patient in a cholinergic crisis there is temporary aggravation of the muscle weakness. This test probably has some value in differentiating the two types of crisis, but it may be difficult to evaluate the result owing to the very short action of the drug.

If a cholinergic crisis (due to overdosage of neostigmine) occurs, administration of the drug must be stopped temporarily, atropine should be given in a dose of 1/60 gr. (1 mg.) intramuscularly, and this dose repeated hourly until signs of atropinization occur. Measures must also be taken to treat weakness of bulbar and respiratory muscles, such as tracheal intubation, tracheotomy with positive-pressure respiration, or assisted breathing in a respirator, depending on the distribution of the muscle weakness.

Pyridostigmine

One of the main disadvantages of neostigmine is its short action, and pyridostigmine ("mestinon"), a pyridine analogue of neostigmine, was introduced in 1954 as it was said to have a rather longer action and was less apt to cause side-effects on the alimentary canal than neostigmine. A 60-mg. tablet of pyridostigmine by mouth corresponds to 15 mg. of neostigmine. The total dosage and timing of their administration is worked out as described for neostigmine. In general, pyridostigmine produces side-effects less frequently and is the more useful drug for the control of myasthenia gravis, but individual patients vary in their response and some prefer to continue with neostigmine. In others a combination of the two drugs gives the best results, neostigmine being used during the day and pyridostigmine for the last dose at night, as its rather longer action may obviate the need for a dose during the night in the more severe cases.

Other Drugs

Organic Phosphate Compounds.—During the past twelve years several long-acting choline esterase inhibitors have been tried in the treatment of myasthenia gravis; the best known are di-isopropyl fluorophosphate (D.F.P.), tetraethylpyrophosphate (T.E.P.T.), and octamethyl pyrophosphoramide (O.M.P.A.). These drugs were found to be uncertain in their action, and toxic effects were frequent and difficult to control. There does not seem to be any place for them in the routine treatment of the disease.

Ephedrine has some effect in increasing muscle power in myasthenia, but the mechanism of this is uncertain. It can be given in a dose of $\frac{1}{2}$ gr. (25 mg.) two or three times a day in addition to neostigmine or pyridostigmine. In my experience it has proved most useful in patients who wake during the night with choking and respiratory difficulty, and it seems likely here that its main value is from its bronchodilator effect.

Drugs to be Avoided

Quinine aggravates myasthenia, an effect opposite to that seen in myotonia, where the delayed relaxation of muscle is often relieved by quinine and aggravated by neostigmine. Myasthenic patients are very sensitive to curare, and this drug should not be administered in operations on patients with the disease unless this danger

is realized and adequate provision made for assisted respiration until the effects of the curare have worn off. A.C.T.H. and cortisone have been given in the treatment of myasthenia, but they often exacerbate the disease after a few days; a remission sometimes occurs after the drug has been stopped. In view of this, great care should be taken in using these drugs for a coincidental disease in a myasthenic patient.

Treatment Apart from Drugs

About 12% of myasthenic patients have a neoplasm of the thymus. It can usually be shown radiologically by screening of the chest, antero-posterior and lateral pictures, and if necessary tomography. These patients should be given deep x-ray therapy to the tumour, and in some cases operative removal of the thymoma should be carried out at a later date. During deep x-ray treatment the necessary dosage of neostigmine may vary considerably from day to day and careful supervision of this is needed. The question of operative removal of the thymus in myasthenia patients who have not got a radiologically demonstrable thymoma has been discussed recently in a leading article in this *Journal* (1959). In general the operation should be carried out in patients under the age of 45–50 if the disease is becoming generalized, or progressively getting worse, and if the patient is needing increasing doses of neostigmine or pyridostigmine and notable remissions are not occurring. In exacerbations of the disease, when there is bulbar palsy or paralysis of the respiratory muscles, the patient needs treatment by methods similar to those used in poliomyelitis or acute infective polyneuritis—namely, postural drainage, physiotherapy to the chest, and in some cases tracheotomy and positive-pressure respiration.

REFERENCE

Brit. med. J., 1959, 1, 288.

REACTIONS OF CANCER PATIENTS ON BEING TOLD THEIR DIAGNOSIS

BY

JEAN AITKEN-SWAN, A.M.I.A., A.H.A.

AND

E. C. EASSON, M.D., F.F.R., D.M.R.T.

*Christie Hospital and Holt Radium Institute,
Manchester*

"I asked the doctor—I looked him straight in the face and I asked 'Is it cancer?' and he said, 'Yes.' Well, I'd rather know the truth—it's better than imagining all the time. I asked my own doctor in the first place, 'was it . . . ?' and I didn't get to say the word, and he said 'We don't know,' so naturally I thought it must be. If I'm ill I'd rather know what I'm suffering from, because you don't die any sooner for knowing about it."

Doctors and journalists sometimes discuss what the patient with serious disease should be told about his illness. The patient's own views are seldom heard. It is generally assumed that patients with cancer, even those with localized and curable conditions, prefer not to be told the diagnosis, and that silence or reassurance is the best solution to the patient's (and doctor's) anxiety. This policy has unfortunate repercussions for cancer education. It means that the only cancer patients