reading, which has proved to be impracticable in conventional mobile mass radiography, would present relatively little difficulty at static units with 100-mm. cut films. It is obvious that even a large number of these static units would be unable to deal with the periodical routine x-ray examination of all men in the critical age groups, but general practitioners could help by sending for chest x-ray examination all men over 35, irrespective of the reasons for visiting their surgeries.

In the meantime, it would be wrong in our opinion to engage in any form of propaganda that suggests that at present routine mass radiography is a certain means of detecting "early cancer," and attention should not be deflected from the much more important preventive aspects of the present situation.

Summary

A series of 238 cases of cancer of the lung in men found by mass radiography in the Midlands are reviewed. Comparisons are made between cases found by conventional routine mass radiography and patients referred to the units by general practitioners.

The 12-months-survival rates irrespective of treatment and the resection rates were higher for routine survey cases than for general-practitioner referrals. A larger proportion of routine survey cases had a lobectomy than general-practitioner patients. Various aspects and reasons for these differences are discussed.

Symptomatic cases were referred by general practitioners generally without undue delay, and the majority of resections were carried out within the first two months after x-ray examination.

The very small number of truly asymptomatic cases and the absence of small peripheral shadows in this series are discussed in their relation to future mass radiography policy.

The installation of more 100-mm. x-ray units to deal with a larger number of general-practitioner referrals, particularly of men in the critical age groups, is suggested.

ADDENDUM: Second-year Survival Rates.—Since this paper was written a second-year follow-up report has been obtained on all but three patients (two general-practitioner referrals and one routine survey case) of the 238 in the series. Assuming all three had died, there were 38 general-practitioner referrals and 12 routine survey cases alive two years after the date of x-ray examination. Hence the second-year survival rates regardless of treatment for the two survey groups were respectively: G.P. referrals, 20%; routine survey cases, 25%. The rate for all patients was 21%. Of the 57 general-practitioner cases and 21 routine survey patients resected, 25 and 8 respectively were alive two years after the date of operation. Hence the second-year survival rates for the two groups of patients were respectively 44% and 38%, and for both groups combined 42%.

We are indebted to the chest physicians and thoracic surgeons in the region for their full co-operation in providing information of patients referred to them. We are grateful to Miss J. M. Levi, of the Birmingham Regional Cancer Registry, for obtaining follow-up details of our patients; to Mrs. M. Sheldon and Miss A. Holdsworth for clerical assistance, and to Miss J. Yates for help with processing of the material.

REFERENCES

- Belcher, J. R. (1956). *Lancet*, **1**, 349.
 Bignall, J. R. (1958a). *Carcinoma of the Lung*, p. 155. Livingstone, Edinburgh.
 — (1958b). *Ibid.*, p. 170.
 — (1958c). *Ibid.*, p. 168.
 — (1958d). *Ibid.*, p. 175.
 — (1958e). *Ibid.*, p. 154.
 — (1958f). *Ibid.*, p. 204.
 — and Moon, A. J. (1955). *Thorax*, **10**, 183.
 Boucot, K. R., and Sokoloff, M. J. (1954). *Amer. Rev. Tuberc.*, **69**, 164.
 Brooks, W. D. W., Davidson, M., Thomas, C. Price, Robson, K., and Smithers, D. W. (1951). *Thorax*, **6**, 1.
 Cleland, W. P. (1958a). In Bignall's *Carcinoma of the Lung*, p. 220. Livingstone, Edinburgh.
 — (1958b). *Ibid.*, p. 227.
 Cross, K. W., McDowell, L. A., and Posner, E. (1958). *Brit. med. J.*, **1**, 862.
 D'Abreu, A. L., and Taylor, A. B. (1957). *Lancet*, **1**, 684.
 Ehler, A., Stranahan, A., and Olson, K. B. (1954). *New Engl. J. Med.*, **251**, 207.
 Ellman, P. (1952). *Essentials in Diseases of the Chest for Students and Practitioners*, p. 46. Oxford Univ. Press, London.
 Gifford, J. H., and Waddington, J. K. B. (1957). *Brit. med. J.*, **1**, 723.
 Guiss, L. W. (1952). *Cancer*, **5**, 1035.
 Hinson, K. F. W. (1958). In Bignall's *Carcinoma of the Lung*, p. 119.
 Kerley, P. J. (1958). In *Carcinoma of the Lung*, p. 21. N.A.P.T., London.
 Lutwyche, V. U. (1956). *Brit. J. Tuberc.*, **50**, 294.
 McDowell, L. A. (1958). *Lancet*, **1**, 632.
 McNulty, J. M. (1954). *New Engl. J. Med.*, **250**, 14.
 Ministry of Health (1956). *Report for the Year Ended 31st December, 1955*, Part II. H.M.S.O., London.
 — (1957a). *Report for the Year Ended 31st December, 1956*, Part II. H.M.S.O., London.
 — (1957b). *Memorandum H.M. (57)*, 94.
 Nicholson, W. F., Fox, M., and Bryce, A. G. (1957). *Lancet*, **1**, 296.
 Overholt, R. H., Bougas, J. A., and Woods, F. M. (1955). *New Engl. J. Med.*, **252**, 429.
 — and Schmidt, I. C. (1949). *J. Amer. med. Ass.*, **141**, 817.
 Rigler, L. G., O'Loughlin, B. J., and Tucker, R. C. (1953). *Dis. Chest*, **23**, 50.
 Smith, R. Abbey (1957). *Thorax*, **12**, 79.
 Taylor, A. B., and Waterhouse, J. A. H. (1950). *Ibid.*, **5**, 257.
 Thomas, C. Price (1952). *Ann. roy. Coll. Surg. Engl.*, **11**, 205.
 Thompson, V. C. (1957). *Lancet*, **1**, 425.
 United Birmingham Hospitals (1954). *Annual Cancer Report*, **2**, 32.

KALA-AZAR OF UNUSUAL DURATION, ASSOCIATED WITH AGAMMAGLOBULINAEMIA

BY

MARY INGLE WRIGHT,* M.R.C.S., D.C.P.

Formerly of the Royal Hants County Hospital, Winchester

The case here reported is of extraordinary duration in that over ten years must have elapsed between the last possible date of infection with *Leishmania donovani* and ultimate diagnosis. It is of further interest because the serum gamma-globulin was abnormally low, and because the patient finally developed secondary amyloidosis.

Case Report

An unmarried Pole aged 31 was admitted to the Royal Hampshire County Hospital, in the care of Dr. W. D. Brinton, in November, 1956, with the following history. He was born in 1925, and had lived and worked on the family farm near Cracow until taken to Austria in 1941, where he worked on farms near Klagenfurt for nearly four years. In 1945 he went to Italy, where he lived for 18 months in camps in the Bologna region; he mentioned Monte Prandone and San Giuliano particularly, and Budrio, a town of some size, ten miles (16 km.) east of Bologna. In the autumn

*Now at the Department of Pathology, Royal Free Hospital, London.

of 1946 he came to England, and has remained here ever since, doing heavy manual work. He had not regarded himself as ill at any time, nor any member of his family as particularly susceptible to infectious diseases. He had, however, been in hospital twice during his ten years in this country.

In July, 1950, he was admitted with delayed healing following a dental extraction. Evening pyrexia, granulopenia, and enlargement of the liver to three fingerbreadths below the costal margin were noted. Jaundice developed and lasted one week. The laboratory investigations supported a diagnosis of infective hepatitis. The W.R. was negative. In September, 1950, he looked well, and the granulopenia and liver-function tests were improving. He returned to work, his weight being 9 st. 5 lb. (59.4 kg.).

His next admission to hospital was in December, 1955. He complained of excessive sweating, chiefly at night, and anorexia for the previous month, following a cough and epistaxis. A few superficial lymph nodes were palpable, as was the spleen, but the liver was not enlarged. The blood pressure was 140/70. All blood cultures, skin tests, agglutinations, and W.R. were negative, but a diagnosis of brucellosis was made, and was supported by an apparent response to chlortetracycline (total dosage 16 g.); the temperature fell to normal, and the spleen diminished in size. After one month in hospital he was discharged; he weighed (as before) 9 st. 5 lb. (59.4 kg.), and returned to heavy work.

In November, 1956, he was admitted as an emergency case, having fainted at work. He said that increasing ill-health and weakness had followed a sore throat two months previously. He had also suffered a moderate laceration of the left hypothenar eminence, which had been sutured, and for which he had been given prophylactic antitetanus serum ten days previously. On examination he was found to be emaciated and febrile (100° F.—37.8° C.), with a moderately enlarged soft spleen, and the edge of the liver was only just above the level of the umbilicus. Blood pressure was 110/70, pulse 90, and weight 7 st. 11 lb. (49.4 kg.). His face, however, looked "healthy" and was well coloured. Although muscular wasting was present in all groups, it was most pronounced in the upper arms. Alopecia was noted in patches on the head and legs. Leucopenia was more pronounced than formerly. Blood cultures, W.R., and agglutinations were again negative.

Kala-azar was specifically discounted at this time because of the length of time since possible infection and also a negative formol-gel test, and a diagnosis of histoplasmosis or toxoplasmosis was thought more probable. Sternal marrow smears showed an active normoblastic erythropoiesis; the granular series was chiefly represented by rather fragile myelocytes, many of which were apparently disrupted in spreading, and few mature forms. Few monocytes, lymphocytes, or plasma cells were seen, and all were mature forms. Platelet production appeared normal. Numerous parasites were seen, mostly intracellular in sections; few were so in the smears, possibly due to the trauma of spreading. The organisms were of about the same size as platelets, and presented a cigar-shaped appearance in the thicker parts of the smears, somewhat similar to *Toxoplasma gondii*. The characteristic appearances of *L. donovani*, with trophonucleus and kinetoplast, were present, however, and were identified by Dr. H. T. Findlay. Cultures were unfortunately overgrown by contaminants. The formol-gel test was still negative, as was the urea-stibamine test. Total protein in the serum was 4.1 g./100 ml. Paper electrophoresis showed a slight diffuse increase in the alpha₁ position and some decrease in albumin. The peripheral blood showed a reticulocytosis of 7% and haemoglobin of 68% (10.2 g./100 ml.).

Treatment was instituted with 15 daily intramuscular injections of 200 mg. of pentamidine isethionate (4:4'-diamidinodiphenoxypentane di-β-hydroxyethane-sulphonate). This caused considerable local inflammation

and pain, and almost certainly prolonged the pyrexia, but the spleen began to diminish in a few days. Owing to the apparent lack of effect of the drug on the temperature, the patient was transferred temporarily to the Hospital for Tropical Diseases, in the care of Professor A. W. Woodruff. While there the patient became afebrile, and absence of gamma-globulin was noted on paper electrophoresis. No further anti-leishmanial treatment was given, and the patient returned to the Royal Hampshire County Hospital.

Four weeks after the end of the pentamidine course he had gained 15 lb. (6.8 kg.). The white cell count was increasing; all tests for increased haemolysis were negative. The spleen was not palpable, but the liver was still enlarged to four fingerbreadths below the costal margin. Proteinuria was noted for the first time, but there were no casts or cells in the urinary deposit. Blood pressure was 140/80. The Congo-red test showed only 17% retention of dye in the plasma at one hour, when 60% or more might be expected in a healthy individual. The patient's serum agglutinated A and B red cells—which were agglutinable by control sera to titres of 1/128 and 1/256—only to titres of 1/8 and 1/4 respectively (patient's red cells were group O). It was not considered necessary to investigate adrenal cortical function at this time, as there had been no generalized increase in pigmentation, the blood pressure had returned to the 1955 level, and the serum chloride was within normal limits (104 mEq/l.; 608 mg./100 ml.). The 24-hour output of neutral 17-ketosteroids was 4.9 mg. (normal for an adult male, 9–14 mg.).

During the next four months the patient's weight increased to 10 st. 4 lb. (65 kg.), and he returned to work. By the end of May, 1957—that is, five months after the end of the pentamidine course—it was clear that some of the earlier improvement was not maintained, and one month later fever had returned and the patient's weight had fallen to 9 st. (57 kg.). The liver and spleen did not appear to have changed in size during the relapse. It was not thought advisable to repeat the Congo-red test in view of the potential danger (MacCuish, 1957). A second course of pentamidine was therefore given, in the same dosage as before, but intravenously. No unpleasant effects followed administration and he improved clinically for a few months, but proteinuria continued, and as the liver and spleen remained large and firm it was thought that amyloid disease had developed. This was not proved until March, 1958, when the patient was readmitted to the Hospital for Tropical Diseases, in the care of Professor A. W. Woodruff.

Discussion

It is often said that kala-azar should not be diagnosed in an illness that has lasted longer than 18 months. But unusually long incubation periods are reported by Sweeney, Friedlander, and Queen (1945), Norman (1946), Stone, Tool, and Pugsley (1952), and Jopling (1955); all these cases became ill two to three years after possible infection, and Mu and Huie (1929) refer to durations of one month to four years. Napier (1922) mentions four cases of more than five years' duration. The only recorded case (Castorina, 1924) of even longer duration is that of an Italian child of 7, who died a proved case with a compatible history since the age of 6 months. Castorina also mentions another boy of 11 in whom the disease had lasted seven years, but in this latter case the presence of the parasite was never proved.

The above patient's spleen was not palpable in 1950. His fever has never shown the classical double daily rise, nor has the blood pressure been recorded below 110/70. There has been progressive anaemia, presumably haemolytic in type; for before treatment in December, 1956, there was a reticulocytosis of 7%, with rapid improvement after pentamidine. Ascites was never present. Anorexia was present only during pyrexia.

Epistaxes occurred in 1955. Diarrhoea was troublesome in December, 1956, prior to treatment; no other specific cause was found. An irritating cough with little sputum was complained of in 1955 and 1956; Napier (1951) states that such a cough may be attributed to vagal stimulation by an enlarged spleen. The nervous system was free from abnormal signs, though the selective wasting of the muscle groups of the upper arm was at first suspected as evidence of motor neurone disease. Patchy alopecia has been observed in all admissions, but no undue pigmentation.

The low output of urinary 17-ketosteroids indicates hypogonadism, which is a feature of the disease (Napier, 1951) in man, and in the experimental animal, the Australian marsupial the possum (Bolliger and Backhouse, 1948). The question of amyloidosis was raised because of the extraordinary length of illness and the fact that it is a feature of the disease proved by biopsy in experimentally infected hamsters (Gellhorn, van Dyke, Pyles, and Tupikova, 1946). These animals do not develop hyperglobulinaemia (Stein and Wertheimer, 1942). Sweeney *et al.* (1945) record 20% retention of Congo red at one hour in their case, which, however, also had proteinuria. 17% and 20% retention is strongly suggestive of amyloid disease.

Clearly, infection with *L. donovani* was possible only during the Italian interlude, and reference to the *World Atlas of Epidemic Diseases* (1952) confirms that the area in which the patient lived for 18 months is one in which the prevalence of visceral leishmaniasis is suspected. It is south of the 10°C. mean annual isotherm, and two *Phlebotomus* species are found there, *P. papatasi* and *P. major*. The accepted vector for *L. donovani* in the western Mediterranean is *P. perniciosus*, which has been identified further north, near Lake Como, and further south, near Pescara, and in Rome and Naples. In the eastern Mediterranean the vector is *P. major*, which has been reported at Ferrara, some 20 miles (32 km.) north of Budrio in the Po Valley. The patient had been nearer the eastern Italian seaboard, 40 miles (64 km.) away, where visceral leishmaniasis is known, on train journeys. A case has been reported (Fusaroli, 1953) at Forlì, only 35 miles (56 km.) south-west of Budrio, in a district where it was not previously recognized.

Part of the interest of this case lies in the low total globulin, especially the gamma fraction. Although the formol-gel test is usually positive in kala-azar, it is not a specific test for the disease, but depends on the very high globulin found in the patient's serum. De Vries (1939) has shown that it is positive only when the serum globulin is not less than 3.7 g./100 ml. In the present case the initial globulin level was only 1.7 g./100 ml., and in previous admissions had not exceeded 2.5 g./100 ml. Actual gamma-globulin estimations, by the method of Wolfson, Cohn, Calvary, and Ichiba (1948), were unfortunately not performed until eight weeks after the end of the pentamidine course, giving a result of 0.4 g./100 ml. (normal 0.8–1.2 g.). Serum from a congenital agammaglobulinaemic, from an affected family, gave a result of 0.12 g./100 ml. by this method. During the ensuing four months the gamma-globulin rose to 0.9 g./100 ml.—that is, to within the accepted limits of normal: and at the same time there was some improvement in protein levels, with an unexplained increase in fibrinogen above normal limits. The erythrocyte sedimentation rate (Wintrobe) rose, in spite of an increased packed cell volume.

Cases of kala-azar with a negative formol-gel test fall into two classes: those in which the characteristic protein changes of falling albumin and rising globulin have not yet developed, changes which usually take four or five months; and cases in which such changes never develop. The latter class is held to carry a poor prognosis; Lloyd, Napier, and Paul (1929) found a high relapse rate in this group. The case reported by Chemnitz and Kirsch (1948) had a negative formol-gel throughout two years of illness, during which he had seven courses of anti-leishmanials, and finally died a few weeks after developing herpes zoster: one month before death the total globulin was 1.7 g./100 ml., of which the euglobulin was 0.1 g.

In view of the fluctuation of serum gamma-globulin levels, some just within normal limits, it seems unlikely that this is a case of congenital agammaglobulinaemia; nor is the personal or family history at all suggestive of it. It seems that this is the acquired type, which may be idiopathic or apparently secondary to other conditions, such as disease of the reticulo-endothelial system (Arends, Coonrad, and Rundles, 1954; Zinneman, Hall, and Heller, 1954; Seltzer, Baron, and Toporek, 1955; Good and Mazzitello, 1956), nephrosis leading to hypoproteinaemia, and inadequate protein intake (Krebs, 1946). These cases do not always show an undue proneness to infection (Martin, 1954) or have an absolute absence of gamma-globulin, but it is assumed that an underlying defect in antibody production exists.

Corkill (1949) has suggested that in kala-azar there is a critical point of hypoproteinaemia "below which the body fails to synthesize gamma-globulin in sufficient quantity or kind to meet specific antibody demands, and at such point any factor tending to break down protein or limit its resynthesis would seem thereby to cause activation of a latent infection, or a relapse, or a shortened incubation period." It is possible that in a case of ten years' duration some such balance was present, and permitted only gradual deterioration of the general clinical condition. There seems to be no convincing intercurrent precipitating factor for any of the exacerbations, and the urine did not contain protein before 1957.

It is of course possible that this prolonged mild infection of the reticulo-endothelial system might in some way have affected normal gamma-globulin production. It is harder to conceive of low immune bodies being responsible for an unusually long and symptomless disease.

Summary

A case of kala-azar in an adult Polish male is reported.

Ten years must have elapsed between the patient's leaving an endemic area and the diagnosis of his disease.

The patient was found to have very low serum gamma-globulin, and it is suggested that this might be secondary to the prolonged disease of the reticulo-endothelial system; secondary amyloidosis developed.

I thank Dr. W. D. Brinton, Dr. C. H. Wrigley, Dr. H. T. Findlay, and Professor A. W. Woodruff for permission to publish this case, and for advice and encouragement.

REFERENCES

- Arends, T., Coonrad, E. V., and Rundles, R. W. (1954). *Amer. J. Med.*, **16**, 833.
- Bolliger, A., and Backhouse, T. C. (1948). *Trans. roy. Soc. trop. Med. Hyg.*, **41**, 797.

- Castorina, G. (1924). *Pediatrics (Napoli)*, **32**, 174.
 Chemnitz, H., and Kirsch, E. (1948). *Z. ges. inn. Med.*, **3**, 336.
 Corkill, N. L. (1949). *Ann. trop. Med. Parasit.*, **43**, 261.
 De Vries, A. (1939). *Acta med. scand.*, **99**, 425.
 Fusaroli, A. (1953). *Riv. ital. Igiene*, **13**, 6.
 Gellhorn, A., van Dyke, H. B., Pyles, W. J., and Tupikova, N. A. (1946). *Proc. Soc. exp. Biol. (N.Y.)*, **61**, 25.
 Good, R. A., and Mazzitello, W. F. (1956). *Dis. Chest*, **29**, 9.
 Jopling, W. H. (1955). *Brit. med. J.*, **2**, 1013.
 Krebs, E. G. (1946). *J. Lab. clin. Med.*, **31**, 85.
 Lloyd, R. B., Napier, L. E., and Paul, S. N. (1929). *Indian J. med. Res.*, **16**, 1065.
 — and Paul, S. N. (1928). *Ibid.*, **16**, 203, 529.
 MacCuish, R. K. (1957). *Brit. med. J.*, **1**, 1403.
 Martin, N. H. (1954). *Lancet*, **2**, 1094.
 Mu, Jui-wu, and Huie, D. (1929). *Nat. med. J. China*, **15**, 436.
 Napier, L. E. (1922). *Indian med. Gaz.*, **57**, 406.
 — (1951). *The British Encyclopaedia of Medical Practice*, 2nd ed., **7**, 295.
 Norman, A. P. (1946). *Lancet*, **2**, 437.
 Seltzer, G., Baron, S., and Toporek, M. (1955). *New Engl. J. Med.*, **252**, 252.
 Stein, L., and Wertheimer, E. (1942). *Ann. trop. Med. Parasit.*, **36**, 17.
 Stone, H. H., Tool, C. D., and Pugsley, W. S. (1952). *Ann. intern. Med.*, **36**, 686.
 Sweeney, J. S., Friedlander, R. D., and Queen, F. B. (1945). *J. Amer. med. Ass.*, **128**, 1020.
 Wolfson, W. O., Cohn, C., Calvary, E., and Ichiba, F. (1948). *Amer. J. clin. Path.*, **18**, 723.
World-Atlas of Epidemic Diseases, Part I, Map 31. Falk, Hamburg, 1952.
 Zinneman, H. H., Hall, W. H., and Heller, B. I. (1954). *J. Amer. med. Ass.*, **156**, 1390.

ELECTROPHORETIC DIFFERENTIAL SERUM PROTEIN PATTERN IN KALA-AZAR*

BY

AJAI SHANKER, M.D.

Lecturer in Medicine, Gandhi Medical College, Bhopal, India

Ever since Tiselius first introduced the technique of electrophoresis in 1937, much work on protein analysis in various diseases has been undertaken—well summed up by Miller (1955) and Sherlock (1955). Kala-azar, which is widespread in India, has, however, had only a sparse and casual mention in the vast literature on the subject. The *British Medical Journal* (1957) referred to the extensive work of Feinstein and Petersdorf (1956) on the diagnostic importance of gross, moderate, and mild hyperglobulinaemia, but failed to specify kala-azar even though this disease involves a large proportion of the world's population. This failure is not surprising when it is realized that there are only six references to serum protein electrophoretic pattern in kala-azar—namely, those of Cooper *et al.* (1946), Benhamou *et al.* (1949), Le Bourdelles *et al.* (1949), Sen Gupta *et al.* (1953), Agarwal (1957), and Kunwar *et al.* (1957). The work of Cooper *et al.* was based on a study of the sera of one active case and one convalescent case of kala-azar. Benhamou *et al.* also studied two cases, Sen Gupta *et al.* studied the sera of 9 cases, and my colleagues and I (Kunwar *et al.*, 1957) published records of 19 cases.

In addition to hypo-albuminaemia and hypergamma-globulinaemia, Benhamou *et al.* (1949) reported the presence in the sera of his patients of a Factor X fraction with more rapid mobility than the albumin fraction. Other workers, however, failed to confirm the presence of such a fraction. Both Cooper *et al.* and Benhamou *et al.* identified in their cases the presence in strength of a γ -globulin of slower mobility than that of normal

γ -globulin. Sen Gupta *et al.* also found this slow-moving γ -globulin in the sera of their 9 cases. This slow-moving fraction, akin to the slow-moving γ_2 -globulin found in the sera of rabbits hyperimmunized against type I pneumococci, was held to be characteristic of kala-azar cases and had not been described in other important conditions associated with hyperglobulinaemia. Sen Gupta *et al.* also found a correlation between the complement-fixation test for kala-azar and the presence of this fraction.

Because of sparse and conflicting reports my colleagues and I (Kunwar *et al.*, 1957) considered it worth while to publish a series of 19 cases of kala-azar in which a reassessment of serum was done after the patients had been cured. Our series was larger than any hitherto published, and our findings were at variance in some respects with those of other workers. It would be interesting to compare the findings with those of other workers. I shall attempt to do that.

Our work was based on an analysis of 19 cases with a definite diagnosis of kala-azar. In each case spleen or sternal puncture was positive for Leishman-Donovan bodies. The initial assessment consisted of the clinical picture, Napier's (1921) aldehyde test, and Chopra's (1936) urea-stibamine test; cephalin-cholesterol flocculation, and thymol turbidity tests; estimation of total serum proteins; and paper electrophoretic analysis of serum protein by Durham's hanging strip electrophoretic apparatus, using Whatman No. 3 mm. filter-paper and sodium barbitone buffer at a pH of 8.6 ($U=0.075$), run for 18 hours at 6 mA. (110 volts). The strips were left in a solution of 0.01% bromophenol blue for 18 hours and were scanned by means of a photovolt scanner. Total and differential white-cell counts were done. An assessment of serum proteins was again made after a full course of urea-stibamine therapy when sternal punctures had become negative for Leishman-Donovan bodies.

Our findings (see Table I) and those of other workers, are given below.

Total Serum Proteins

Our 19 cases showed a hyperproteinaemia (average 7.58 g./100 ml.†) before the start of therapy (Kunwar *et al.*, 1957). Only two cases had an initial total serum protein level of less than 7.4 g./100 ml. Normal averages with the same technique in a separate study (Misra, Shanker, and Bajpai, unpublished) were found to be 7.07 g. It is shown below that this rise in total serum protein occurred in spite of marked hypo-albuminaemia and was mainly due to a rise in γ -globulin. However, Agarwal (1957) failed to elicit any rise in the total serum protein.

Serum Albumin

Hypo-albuminaemia had been reported by Chakravarty *et al.* (1949). All of our cases showed a marked fall in the serum-albumin level (average† 30.14%; range, 20.16%–36.22%). Sen Gupta *et al.* and Agarwal also found a generalized fall of serum albumin, the average value in Agarwal's cases being 25.36%. We found no oedema in spite of hypo-albuminaemia, and the presumptive conclusion is that hypo-albuminaemia is due to hepatic damage.

Alpha-globulins (α_1 and α_2).—These fractions showed only a slight rise of doubtful significance. Normal

†These average figures represent only 14 cases which were followed up.

*Work on the subject was done at the department of medicine, King George's Medical College, Lucknow, in collaboration with Dr. K. B. Kunwar and Dr. J. S. Bajpai.