REFERENCES

- Ansell, G. B., and Richter, D. (1954). *Biochim. biophys. Acta*, **13**, 87, 92.
- Block, R. J. (1937). *J. biol. Chem.*, **120**, 467.
- Block, W. (1954). *Hoppe-Seyler's Z. physiol. Chem.*, **296**, 108.
- Brattgård, S. O. (1952). *Acta radiol (Stockh.)*, Suppl. 96.
- Chapman, L. F., and Wolff, H. G. (1958). *Trans. Ass. Amer. Physns.*, **71**, 210.
- Clarke, D. D., Neidle, A., Sarkar, N. K., and Waelsch, H. (1957). *Arch. Biochem.*, **71**, 277.
- Clouet, D. H., and Richter, D. (1959). *J. Neurochem.*, **3**, 219.
- Cohn, P., Gaitonde, M. K., and Richter, D. (1954). *J. Physiol.*, **126**, 7 P.
- (1959). *J. Neurochem.*, in press.
- Flexner, L. B. (1951-2). *Harvey Lectures*, **47**, 156.
- Folch, J., and Lees, M. (1957). *J. biol. Chem.*, **191**, 807.
- and Uzman, L. L. (1948). *Fed. Proc.*, **7**, 155.
- Gaitonde, M. K., and Richter, D. (1956). *Proc. roy. Soc. (B)*, **145**, 83.
- Gersh, I., and Bodian, D. (1943). *J. cell. comp. Physiol.*, **21**, 253.
- Greenberg, D. M., Friedberg, F., Schulman, M. P., and Winnick, T. (1948). *Cold Spring Harbor Symp. quant. Biol.*, **13**, 113.
- Hamburgh, M., and Flexner, L. B. (1957). *J. Neurochem.*, **1**, 279.
- Heald, P. J. (1958). *Biochem. J.*, **68**, 580.
- Karcher, D., van Sande, M., and Lowenthal, A. (1959). *J. Neurochem.*, **4**, in press.
- Kiyota, K. (1957). *Ibid.*, **1**, 301.
- Koenig, H. (1959). *Ibid.*, **4**, in press.
- LeBaron, F. N. and Folch, J. (1956). *Ibid.*, **1**, 101.
- Mullins, L. J. (1953). *Amer. J. Physiol.*, **175**, 358.
- Porter, H. (1959). In *Chemical Pathology of the Brain*, edited by J. Folch. Pergamon Press, London. In press.
- and Folch, J. (1957). *J. Neurochem.*, **1**, 260.
- Reiss, J. M., Reiss, M., and Wyatt, A. (1956). *Proc. Soc. exp. Biol. N.Y.*, **93**, 19.
- Richter, D. (1957). *Schizophrenia: Somatic Aspects*. Pergamon Press, London.
- Robertson, D. M. (1957). *J. Neurochem.*, **1**, 358.
- (1959). *Ibid.* In press.
- Rose, W. C., Johnson, J. E., and Haines, W. J. (1950). *J. biol. Chem.*, **182**, 541.
- Schmitt, F. O. (1957). In *Metabolism of the Nervous System*, edited by D. Richter. Pergamon Press, London.
- Ungar, G., Aschheim, E., Psychoyos, S., and Romano, D. V. (1957). *J. gen. Physiol.*, **40**, 635.
- Vrba, R. (1956). *J. Neurochem.*, **1**, 12.
- Waelsch, H. (1957). In *Metabolism of the Nervous System*, edited by D. Richter, p. 431. Pergamon Press, London.
- Winzler, R. J., Moldave, K., Rafelson, M. E., and Pearson, H. E. (1952). *J. biol. Chem.*, **199**, 485.

TRIAMCINOLONE IN THE TREATMENT OF RHEUMATOID ARTHRITIS

BY

MALCOLM THOMPSON, M.D., M.R.C.P.

Consultant Physician, Department of Physical Medicine,
Royal Victoria Infirmary, Newcastle upon Tyne

Triamcinolone is a synthetic steroid, being the 9 α -fluoro-16 α -hydroxy analogue of prednisolone (see Formula). It has been shown to possess powerful anti-inflammatory properties with minimal mineralo-corticoid activity. Bernstein *et al.* (1956) reported that, by rat-liver glycogen assay, triamcinolone was 18 to 36 times as potent as hydrocortisone, and in the adrenalectomized rat it did not produce any sodium retention. Hellman *et al.* (1957) described its clinical and metabolic effects, and demonstrated that it had excellent anti-inflammatory properties with, in ordinary therapeutic dosage, little or no effect upon nitrogen, calcium, potassium, or sodium balance. Assessment of the anti-inflammatory effects of triamcinolone suggested that it was more powerful than prednisolone in an approximate ratio of 4:3 when administered orally.

Sherwood and Cooke (1957) described the results of treatment of 26 patients suffering from allergic conditions, chiefly asthma or dermatitis, and thought that an equal weight of triamcinolone was about twice as potent as prednisone or prednisolone. Rein *et al.*

(1957) also reported upon the use of triamcinolone in dermatological cases, and concluded that an equivalent therapeutic effect was produced by a dosage between one-half and two-thirds of that of prednisolone. Shelley *et al.* (1958) considered that triamcinolone had a significant suppressive effect on psoriasis in 60% of their 60 cases, but that the relapse which invariably followed cessation of treatment limited its value.

Hartung (1958) reported favourably on the use of triamcinolone in 67 patients suffering from rheumatoid arthritis, and stated that it possessed several advantages over other steroids, including fewer gastro-intestinal symptoms. Dubois (1958) treated 31 cases of disseminated lupus erythematosus with triamcinolone, and his experience suggested that this new steroid did not precipitate gastro-duodenal symptoms or lesions. Hart *et al.* (1958) did not find any evidence that triamcinolone either aggravated or reduced gastro-duodenal side-effects in their series of 25 rheumatoid patients, all of whom had received previous treatment with prednisolone.

Since the demonstration by Bunim *et al.* (1955) that the Δ -1- or predni- analogues possessed four or five times the anti-inflammatory activity of cortisone and hydrocortisone, with much fewer sodium-retaining effects, there have been further attempts to find even more powerful steroid compounds. The mere anti-inflammatory potency of a steroid compound is not, of itself, important in assessing the value of the drug as a therapeutic agent, as a less potent steroid could be administered in larger dosage provided this did not produce undesirable side-effects. It is largely on the basis of side-effects, therefore, that a new steroid is to be evaluated. The lessening or elimination of mineralo-corticoid effects, especially hypertension and hypopotaemia, are important considerations, but other side-effects to be considered are peptic ulceration, mental disturbance, osteoporosis, lowered glucose tolerance, purpura, telangiectasia, and subcutaneous haemorrhages, facial mooning and flushing, and acne and hirsuties.

The purposes of the present investigation were to determine (a) whether triamcinolone was as effective or better than prednisolone when administered in a dosage not exceeding four-fifths of the prednisolone dosage; (b) whether triamcinolone warranted a place for selected, or indeed all, patients for whom systemic

steroid therapy was considered advisable; (c) whether, on the basis of a short-term trial, side-effects appeared to be qualitatively or quantitatively different from those seen with prednisone or cortisone; and (d) whether triamcinolone could be used as an immediate substitute for prednisolone or cortisone in the event of side-effects arising with the latter drugs.

Method of the Trial

Of the 16 patients concerned, 15 were suffering from rheumatoid arthritis and one from ankylosing spondylitis with predominantly peripheral arthritis. All received treatment successively with prednisolone and triamcinolone, the ratio of dosage being 5:4 by weight. The daily maintenance dosage of prednisolone ranged from 7.5 to 20 mg. The duration of treatment with triamcinolone varied from 2 to 26 weeks, and 11 of the patients were then given a further course of treatment with prednisolone, allowing a second comparison to be made. The patients had become accustomed to receiving steroid tablets of different sizes from different manufacturers, and they were not informed that any change in treatment had been made, so that any bias in favour of a new treatment was avoided.

Response to treatment was judged by the patient's own assessment of pain and the duration of morning stiffness, in addition to the physician's assessments of functional capacity and disease activity. Alterations in the number of joints affected were noted, and blood pressure, weight, and strength of grip were recorded. Grip was tested at weekly intervals by recording the height in millimetres of mercury reached by squeezing a rubber bag 6 by 3 in. (15 by 7.5 cm.) attached to a sphygmomanometer. Three measurements of grip were made for each hand, and the average was calculated. Then the average for both hands was calculated. Serial recordings of capillary resistance, measured by the apparatus and method described by Potter and Wigzell (1957), in their first survey of capillary resistance in rheumatoid arthritis, were made in seven patients. Serial measurements of the erythrocyte sedimentation rate (Westergren method) were done at weekly or fortnightly intervals. The lowest reading obtained while the patient was receiving prednisolone was compared with the average reading recorded while the patient was receiving triamcinolone.

Results

The results are presented in Table I. Of the 16 patients, six preferred triamcinolone, five preferred prednisolone, and five had no preference. These preferences were based on the patient's own assessments of pain and stiffness, plus sense of well-being. In respect of pain, six patients preferred triamcinolone, four felt pains to be increased while on triamcinolone, and six noticed no difference. With regard to severity and duration of morning stiffness, seven patients considered this to be less while they were receiving triamcinolone, two felt it to be worse, and in seven there was no difference. Functional capacity improved significantly in three patients while receiving triamcinolone; in the remainder it was unchanged. Disease activity was significantly reduced in two patients while receiving triamcinolone, and in the remainder it was unchanged.

Measurements of the grip, by the method described above, were recorded in all cases. In view of ordinary variations in this measurement, an overall average

TABLE I.—Results of Treatment with Triamcinolone Compared with Those of Prednisolone

Case No.	Sex	Joint Pains	Stiffness	Functional Capacity	Disease Activity	Grip	E.S.R.		Patient's Preference
							Pred.	Triam.	
1	F	Less	Less	0	0	0	38	33	Triam.
2	M	"	"	0	0	Imp.	42	26	"
3	F	"	"	0	0	0	12	10	"
4	F	"	"	0	0	0	28	25	None
5	F	0	Less	0	0	0	4	13	"
6	F	0	0	0	0	0	20	15	"
7	F	0	0	0	0	0	6	6	"
8	F	Inc.	Inc.	0	0	0	11	14	Pred.
9	M	Less	Less	Imp.	Less	Imp.	27	12	Triam.
10	F	0	0	0	0	0	16	12	None
11	F	Less	Less	Imp.	0	Imp.	5	5	Triam.
12	F	0	0	0	0	0	28	17	Pred.
13	F	Inc.	0	0	0	0	13	12	"
14	F	0	0	0	0	0	39	45	"
15	M	"	Inc.	0	0	0	15	11	"
16	M	Less	Less	Imp.	Less	Imp.	33	15	Triam.

improvement of at least 20 mm. of mercury was considered necessary to constitute significant improvement. This level was achieved in four patients during triamcinolone therapy, there being no significant increase or decrease in the remaining 12.

Comparisons of the E.S.R. during the two treatments showed it to be lower during triamcinolone therapy in 11 instances, equal in two, and lower during prednisolone therapy in three. In only two patients, however (Cases 9 and 16), was the difference at all striking, and both of these were in favour of triamcinolone. The average lowest E.S.R. recorded, for the group as a whole, during prednisolone therapy was 21 mm., compared with an average of 16.6 mm. during triamcinolone therapy.

Serial measurements of the capillary resistance in seven patients did not show any important difference, the high levels characteristic of patients receiving systemic steroid therapy being found during both treatments.

Side-Effects

The side-effects are recorded in Table II. All but two patients showed *loss of weight* while receiving triamcinolone. Of the five patients who lost 4 lb. or more, three had developed oedema of the ankles during previous steroid therapy. No significant changes in the levels of plasma sodium, potassium, or chloride were seen in any of these patients, and in no instance was there any complaint of weakness or muscular fatigue. (Case 10 was on a 1,000-calorie diet during the period of the trial and no comparison of weight was possible.) The last, and representative, *blood-pressure* readings during prednisolone therapy are compared with those obtained at the end of the period of triamcinolone therapy. Nine patients had a diastolic pressure of 90 mm. or more during prednisolone therapy; in six of them there had been no evidence of hypertension prior to steroid therapy (cortisone or prednisolone); in three instances a pre-existing hypertension had increased during steroid therapy. In all nine patients there was a fall in blood pressure during treatment with triamcinolone. This was especially so in two known hypertensive patients (Cases 2 and 3), who had shown B.P. levels of over 220 systolic and 120 diastolic on occasion during prednisolone therapy.

In respect of *dyspepsia*, it is seen that four patients who complained of dyspepsia during prednisolone therapy retained these symptoms during triamcinolone therapy and in three instances they were more severe. In addition a further four patients complained of dyspepsia for the first time during triamcinolone

therapy, and one of these had haematemesis following vomiting attacks extending over three days.

Two patients had developed *purpura* and *multiple subcutaneous ecchymoses* during prednisolone therapy, and these lesions regressed when triamcinolone was substituted. One of these (Case 3) subsequently exhibited three small areas of apparently spontaneous subcutaneous bruising, but it was at this time that her triamcinolone was discontinued, so that further observation of the progress of the lesions was not possible. *Acneiform skin lesions*, noted in two patients, remained unchanged during treatment with triamcinolone. *Facial mooning*, evident in three patients, and *hirsuties*, troublesome in one, remained unaltered during both treatments.

Two patients were noted to be *euphoric* and restless while taking prednisolone. Both were calmer and less loquacious while taking triamcinolone, and one complained of drowsiness.

One patient had developed glycosuria with a diabetic type of blood-sugar curve during prednisolone therapy. These features remained unchanged when she was given triamcinolone, a repeat blood-sugar curve after 18 weeks of triamcinolone therapy being virtually unchanged. *Flushing and sweating* were symptoms reported by two patients during triamcinolone therapy and not observed with prednisolone.

Additional Cases

In addition to the patients concerned in the trial, a further four patients have been treated with triamcinolone. Three of these were women suffering from rheumatoid arthritis, and the fourth was a man suffering from ankylosing spondylitis. The dosage was 4 mg. four times daily initially, reducing to 12 mg. total daily dose. Two of the female patients have had continuous treatment for 13 weeks without any evidence of side-effects, and one of them, who had pre-existing hypertension, has shown consistently lower blood-pressure readings since she started treatment.

The third female patient has also obtained sustained benefit during the same period, the only complication being the appearance of a crop of spontaneous ecchymoses over the ankles and shins 10 days after the starting of treatment. In the absence of any evidence of haemorrhagic complications it was decided to continue treatment, under close observation, with the same dosage of triamcinolone, 12 mg. daily. The lesions

faded during the ensuing week and no new lesions appeared until 10 weeks later, when similar small crops of lesions occurred over the forearms and shins.

The fourth patient, a man aged 27, had ankylosing spondylitis with severe peripheral arthritis. He had taken large quantities of A.P.C. tablets, up to 20 daily, and calcium-aspirin tablets, with only slight dyspepsia. A barium-meal examination did not reveal any evidence of a peptic ulcer, and triamcinolone therapy was started on June 2, with a maintenance dosage of 16 mg. daily. He responded well, and within a week his E.S.R. fell from 112 to 31 mm. in one hour. After brief premonitory symptoms of nausea on July 2 he had a profuse haematemesis and was admitted to hospital immediately in a state of severe shock. He recovered with transfusions of 4 pints (2 litres) of blood, and steroid therapy was maintained, initially with intravenous hydrocortisone and subsequently with intramuscular cortisone. He has since been treated with injections of zinc-protamine corticotrophin and has remained well on a maintenance dosage of 30 units every third day, without any recurrence of gastric symptoms.

Discussion

Although the place of a new steroid preparation in the treatment of a chronic disease can be fully assessed only on long-term therapy, short-term studies do provide some useful information. It is evident that, in answer to the first question raised in the introduction, triamcinolone is as effective as prednisolone in a dosage ratio of 4:5. When any advantage occurred in changing from one drug to the other the difference was small. The findings of Sherwood and Cooke (1957) in the treatment of allergic disorders, and of Rein *et al.* (1957) with regard to skin diseases, that triamcinolone could be as effective as prednisolone in half to two-thirds the dosage, were not substantiated in this series of patients treated for rheumatoid arthritis. The maintenance dosage was found to be between 6 and 16 mg. daily according to the severity of the case.

In this trial triamcinolone did not exercise any adverse influence upon hypertension. Where hypertension had been induced (six patients) or increased (three patients) by previous steroid therapy, consistent improvement occurred on transfer to triamcinolone. Only long-term studies will determine whether this effect will be maintained or whether it may be associated with other, possibly undesirable, effects—

TABLE II.—*Side-effects of Triamcinolone Compared with those of Prednisolone*

Case No.	Sex and Age	Duration of Treatment (Weeks)	Weight Loss		Blood Pressure		Side-effects	
			lb.	g.	Pred.	Triam.	Prednisolone	Triamcinolone
1	F 51	11	1½	680	160/92	132/80	Moon face. Hirsuties. Hypertension. Mild dyspepsia	Moon face. Hirsuties. Increased dyspepsia
2	M 53	4	2½	1,134	192/116	164/106	Increased hypertension	Decreased hypertension
3	F 49	12	5½	2,495	218/104	178/90	Increased hypertension. Bruising	Less bruising. Less hypertension. Haematemesis
4	F 37	14	4½	2,040	142/92	134/82	Hypertension. Bruising and purpura	Mild dyspepsia
5	F 58	14	1½	680	160/90	134/84	Moon face. Hypertension	Moon face
6	F 67	3	4	1,815	154/94	132/82	Hypertension. Euphoria + +.	Less oedema of ankles.
							Oedema of ankles	Less euphoria
7	F 62	6	Nil		174/100	154/86	Euphoria. Hypertension	Flushing and sweating.
8	F 60	2	1½	680	115/64	114/66	Mild dyspepsia	Slight drowsiness
9	M 49	3	4	1,815	128/82	124/80	Acne	Mild dyspepsia. Flushing and sweating
10	F 53	20	On diet		140/86	122/80	Glycosuria	Acne
11	F 46	15	3	1,360	140/84	134/72	Glycosuria	Glycosuria
12	F 50	2	Nil		108/69	108/66	Moon face	Moon face
13	F 71	3	1½	680	184/104	168/86	Nil	Nil
14	F 47	9	3	1,360	184/108	143/84	Mild dyspepsia. Hypertension	Dyspepsia
15	M 40	2	6	2,720	128/86	138/86	Acne	Acne. Anorexia and nausea. Dyspepsia
16	M 57	5	1	455	128/84	120/72	Nil	Mild dyspepsia

for example, sodium depletion. Loss of weight was seen in most of the patients transferred from prednisolone to triamcinolone; and oedema, when present, was diminished. No weight loss was seen in the four patients who had not received previous steroid therapy, but such weight loss has been recorded (Hellman *et al.*, 1957). At present, therefore, triamcinolone would appear to be the compound of choice when steroid therapy is indicated for patients who have hypertension or evidence of sodium retention, or who have developed these features as a result of previous steroid therapy.

Apart from hypertension and oedema, the other side-effects seen during triamcinolone therapy were very similar to those of prednisolone. Dyspepsia was more evident with triamcinolone, this finding being at variance with the experience of Hartung (1958) and Dubois (1958). The fact that, in this series, 2 out of 20 patients treated suffered haematemeses is an additional warning that triamcinolone can cause and aggravate gastro-duodenal lesions. One of these patients had previously had two years of continuous steroid therapy without dyspepsia.

In respect of other side-effects the only striking difference observed was in one patient (Case 4) who developed multiple subcutaneous ecchymoses during prednisolone therapy, which cleared rapidly on transfer to triamcinolone and have not recurred during 14 weeks of treatment. Another patient (Case 2) showed similar lesions with prednisolone which also cleared on transfer to triamcinolone but recurred slightly after 12 weeks of treatment, when triamcinolone was discontinued because of a haematemesis. Skin lesions of this type occurring during steroid therapy have been studied by Nashelsky and Smyth (1958) and appear to be particularly common with the "predni-" groups of steroids, where there is a double bond in the 1, 2 position of the steroid nucleus. Their occurrence during treatment with another prednisteroid, 6 α -methyl-prednisolone, has been described by Boland and Liddle (1957). It is clear that such lesions occur also during triamcinolone therapy, as noted in two patients in the present series, but their appearance and severity in individual patients did not parallel their severity during prednisolone therapy. The clinical significance of these ecchymoses is as yet unknown; they are not apparently associated with any haemorrhagic tendency or visceral lesions, and investigations of capillary resistance have indicated raised levels in adjacent skin areas. In the event of such lesions being severe and unsightly, a change from prednisolone to triamcinolone would be indicated.

In respect of other complications—for example, glycosuria, moon face, hirsuties, facial flushing, sweating, and acne—no important difference was observed between the two groups. No observations in respect of osteoporosis could be made in this trial. In no instance did any unusual or new arthritic lesions develop, as seen by Wells (1958) during the treatment of certain non-articular diseases with triamcinolone.

When the more serious complications of steroid therapy develop there may be an urgent need to change the steroid preparation, followed when necessary by gradual withdrawal of steroid therapy altogether. On the results of this trial and others, triamcinolone would appear to be a suitable steroid compound when steroid therapy was indicated for (a) hypertensive patients or those who have developed hypertension with another steroid; (b) patients with oedema and evidence of salt

retention; (c) patients who develop purpuric or ecchymotic lesions during treatment with other steroids; and (d) it may be of value for patients who have become restless or unduly euphoric on other steroids.

Modifications of the chemical structure of cortisone, including methyl-, hydroxyl-, and halogen substitution, and dehydrogenation, have produced more potent steroid compounds, with fewer undesirable side-effects. To some extent the pharmacological properties of a new steroid can be predicted from its chemical formula. It is possible that further compounds, representing almost pure anti-inflammatory substances, may become available. Certainly a steroid free from gastro-duodenal side-effects would be a valuable drug in present circumstances. However, the long-term consequences of anti-inflammatory therapy in rheumatoid arthritis are not yet known, and the place of steroid therapy in this disease is still in question. Despite the introduction of stronger and safer steroids, the maxim that "the absence of contraindications does not mean that the drug is indicated" is to be kept in mind.

Summary

The treatment of 15 patients suffering from rheumatoid arthritis and one suffering from ankylosing spondylitis was changed from prednisolone to triamcinolone for a period of 2 to 20 weeks. In addition four other patients were treated with triamcinolone. Triamcinolone was found to be as effective as prednisolone in relieving arthritic symptoms in a dosage ratio of 4 : 5. The average daily maintenance dosage of triamcinolone ranged from 6 to 16 mg., according to the severity of the disease.

Triamcinolone was found to be useful in hypertensive patients when hypertension had been induced or increased by previous steroid therapy. Weight loss and reduction of oedema were noted when patients receiving prednisolone were transferred to triamcinolone. Gastro-duodenal side-effects were more frequent and severe with triamcinolone than with prednisolone, and in the total series of 20 patients two had haematemeses while taking triamcinolone. Subcutaneous ecchymoses and other side-effects of prednisolone were also encountered with triamcinolone, but the pattern of involvement was not identical.

The value and place of triamcinolone in the treatment of rheumatoid arthritis are discussed.

I wish to acknowledge the help of Mrs. J. Dobson and Mrs. M. Downey in the preparation of this paper.

REFERENCES

- Bernstein, S., Lenhard, R. H., Allen, W. S., Heller, M., Littell, R., Stolar, S. M., Feldman, L. I., Blank, H. H. (1956). *J. Amer. chem. Soc.*, **78**, 5693.
- Boland, E. W., and Liddle, G. W. (1957). *Ann. rheum. Dis.*, **16**, 297.
- Bunim, J. J., Pechet, M. M., and Bollet, A. J. (1955). *J. Amer. med. Ass.*, **157**, 311.
- Dubois, E. L. (1958). *Ibid.*, **167**, 1590.
- Hart, F. D., Golding, J. R., and Burley, D. (1958). *Lancet*, **2**, 495.
- Hartung, E. F. (1958). *J. Amer. med. Ass.*, **167**, 973.
- Hellman, L., Zumoff, B., Schwartz, M. K., Gallagher, T. F., Bernstein, C. A., and Freyberg, R. H. (1957). *Ann. rheum. Dis.*, **16**, 141. (Abstract.)
- Nashelsky, G. M., and Smyth, C. J. (1958). *Ibid.*, **17**, 76.
- Potter, J. L., and Wiggall, F. W. (1957). *Ibid.*, **16**, 357.
- Rein, C. R., Fleischmajer, R., and Rosenthal, A. L. (1957). *J. Amer. med. Ass.*, **165**, 1821.
- Shelley, W. B., Harun, J. S., and Pillsbury, D. M. (1958). *Ibid.*, **167**, 959.
- Sherwood, H., and Cooke, R. A. (1957). *J. Allergy*, **28**, 97.
- Wells, R. (1958). *Lancet*, **2**, 498.