active stage again at the time of this review; and 170 (68.5%) were no longer active. We have no information about the last 11.

Conclusions and Summary

As there was enough evidence available by 1949 that B.C.G. was of value in the prevention of tuberculosis, it would have been ethically indefensible to withhold B.C.G. vaccination from any of our contact babies and so study comparable series of unvaccinated controls. Consequently, there is no absolute evidence that B.C.G. vaccination prevented tuberculosis in any. Nevertheless, the fact that from such a particularly unfavourably situated group of infants none died of tuberculosis, and only two developed symptoms which could possibly be attributed to tuberculous infection, suggests that the vaccinations were of value. It is more definite and satisfactory to record the prolonged effectiveness of the vaccination as judged by the persistence of the positive tuberculin reaction in 98% of the cases for five years or more.

267 children aged 2 years or less who were in close household contact with cases of tuberculosis and who remained tuberculin negative six weeks after segregation from contact were vaccinated with B.C.G. between October, 1949, and October, 1952. Of these, 261 were traced five years or more after vaccination. There were no deaths from tuberculosis; only two had minor clinical symptoms associated with tuberculous lung lesions which healed by calcification, and one other had a calcified lung lesion. There were only four who reverted and became tuberculin negative. In contrast, of the "index" cases 39 (15.7%) died of their tuberculosis and 28 (11.3%) still had active disease at the time of this review.

As there were no comparable controls it cannot be shown conclusively that B.C.G. vaccination prevented tuberculosis in this group of infants. Nevertheless the results are highly suggestive that this, in fact, was the case.

It is a pleasure to acknowledge and thank the help given by the chest physicians of Sheffield and many other areas; Dr. Llywellyn Roberts, the Medical Officer of Health of Sheffield, and his health visitors; and Dr. Marion C. Taylor, principal school medical officer of Sheffield, without whose help it would have been impossible to secure such a full follow-up; Professor R. S. Illingworth for his criticism; and the Tuberculosis Research Fund of the University of Sheffield for its grant to one of us (P. C. M.).

REFERENCES

- Aronson, J. D., Aronson, C. F., and Taylor, H. C. (1958). *A.M.A. Arch. intern. Med.*, **101**, 881.
 Briggs, B., Illingworth, R. S., and Lorber, J. (1955). *Lancet*, **1**, 263.
 Hyge, T. V. (1947). *Acta tuberc. scand.*, **21**, 1.
 Medical Research Council (1956). *Brit. med. J.*, **1**, 413.

The second edition of Colonel Sir R. N. Chopra's *Indigenous Drugs of India* has been largely rewritten by the original author in collaboration with I. C. Chopra, K. L. Handa, and L. D. Kapur. In its preparation account has been taken of the considerable accumulation of relevant literature since the appearance of the first edition in 1933. This is a very useful reference book, carefully documented and fully indexed. It is published by Dhur and Sons Ltd., 15, Bankim Chatterjee Street, Calcutta, 12 (pp. 816+xxxii; 1958. £5 5s.).

"B.C.G. SARCOIDOSIS"

BY

PHILIP ELLMAN, M.D., F.R.C.P.

Consultant Physician in Diseases of the Chest, North-east Metropolitan Hospital Board, and the Chest Unit, Plaistow Hospital, London

AND

LESLIE G. ANDREWS, M.B., D.C.H.

Registrar, Chest Unit, Plaistow Hospital, London

Since the earliest major contributions on sarcoidosis were made (Hutchinson, 1875-8; Besnier, 1889; Boeck, 1899, 1905; Schaumann, 1914; Jüngling, 1919) the aetiology has been much debated. Hutchinson (1898), in his study of the disease, thought the cause "may not improbably be a tuberculous affection and one of the lupus family." A large school of thought believes sarcoidosis to be a group of granulomatous diseases of known distinct aetiology—for example, tuberculosis, histoplasmosis, brucellosis, berylliosis—or to be idiopathic. Scadding (1950, 1956b) recommended the use of the terms "tuberculous sarcoid," "histoplasma sarcoid," "beryllium sarcoid," etc., whilst idiopathic cases belong to the tuberculous group passing through an unusual phase of tuberculosis. In latter years cases of sarcoidosis occurring in patients previously given B.C.G. vaccination have been recorded in the literature (Larsen, 1950; Löfgren and Lundbäck, 1952; Törnell, 1954; Pfisterer *et al.*, 1954; Ganguin, 1956; Birkhäuser, 1957; Fried and Genz, 1958). The term "B.C.G. sarcoidosis" is being applied to these cases. We record here a further case of histologically proved sarcoidosis occurring in a girl given B.C.G. vaccination seven years previously.

Case Report

In December, 1950, when the patient was 14 years old, she was vaccinated with Danish B.C.G. as a Mantoux-negative school-leaver. She was subsequently followed up under the Medical Research Council Tuberculosis Vaccines Clinical Trial by skin-sensitivity testing and chest x-ray examinations (see Table).

Results of Follow-up

Date	Skin Test (Same Batch of Old Tuberculin was used throughout except on 2/9/57 and 1/9/58)	Chest X-Ray Examination
1/12/50	Negative 1/100	Normal
7/12/50	B.C.G. vaccination	
26/2/51	Positive 1/3,000 (10 mm.)	"
8/7/52	—	"
18/11/53	Positive 1/3,000 (7 mm.)	"
16/2/55	Negative 1/3,000	"
15/5/56	Positive 1/100 (12 mm.)	"
23/7/57	Positive 1/3,000 (14 mm.)	"
	Negative 1/3,000	Reticular mottling right upper and mid-zones and left upper zone
2/9/57	Negative 1/1,000	Reticular mottling of whole of right lung and left upper and mid-zones
12/9/57	" 1/100	
1/9/58	Heaf test positive	
		Normal

In July, 1957, her Mantoux reaction (1/3,000) had become negative, and reticular mottling had appeared in the right upper and mid-zones and left upper zone of her chest x-ray picture. In September she was admitted to Plaistow Hospital Chest Unit under the care of one of us (P. E.).

The patient was asymptomatic. On examination small cervical and scalene glands were palpable and the liver edge was felt one fingerbreadth below the right costal margin. The spleen was just felt on inspiration. The results of further investigations were: haemoglobin, 12.6 g./100 ml.

(87%); W.B.C., from 5,000 to 9,000/c.mm., with a normal differential; E.S.R., 21 mm./hour (Westergren); total serum protein, 6.9 g./100 ml. (albumin 4.4 g., globulin 2.5 g.); serum protein electrophoretic pattern normal; serum calcium, 5 mEq/l.; inorganic phosphorus, 3.2 mEq/l.; serum bilirubin, 0.4 mg./100 ml.; alkaline phosphatase, 3.3 units; thymol turbidity, 4.5 units; zinc sulphate, 5.5 units; laryngeal swabs and gastric and tracheal lavages negative for acid-fast bacilli; x-ray film of the hands normal. The chest x-ray film showed reticular mottling involving the whole of the right lung field and the left upper and mid-zones (Fig. 1). A liver biopsy showed florid changes compatible

FIG. 1.—X-ray film showing reticular mottling involving almost all of right lung field and left upper and mid-zones.

FIG. 2.—Twelve months later. X-ray film showing complete resolution of lung reticulation.

with sarcoidosis; there were multiple granulomata exhibiting giant-cell formation, but no caseation was seen nor were acid-fast bacilli demonstrated. The Kveim test was negative, using two different potent antigens.

The patient was given no active treatment, and after her discharge from hospital in October, 1957, she returned to work. In December she continued to be asymptomatic, no abnormalities could be elicited clinically, and the chest x-ray film showed obvious evidence of resolution. Radiological improvement continued, and by September, 1958, the chest x-ray film was within normal limits. At that time the W.B.C. was 4,500/c.mm., with a normal differential; E.S.R., 6 mm./hour; total serum protein, 7.9 g./100 ml. (albumin 4.7 g., globulin 3.2 g.); serum protein electrophoretic pattern normal; serum calcium, 5 mEq/l.; and the Heaf tuberculin test was positive. Her chest x-ray picture (Fig. 2) has continued to be normal.

Discussion

Berg (1940) first described a regional lymphadenitis after B.C.G. vaccination, presenting the histological picture of sarcoidosis. Since then, as already mentioned, several references to sarcoidosis occurring in B.C.G.-vaccinated subjects have appeared in the world literature. A proportion of these cases fulfilled the histological criteria; in others the diagnosis was made on radiological and/or clinical grounds. Ganguin (1956) reported two vaccinated siblings exposed to tuberculosis, one of whom developed sarcoidosis, the other's illness being interpreted as atypical tuberculosis. The negative Kveim test in our case is still in keeping with the diagnosis. In a series of 110 Kveim tests the investigation has proved positive in 75% of patients with sarcoidosis (James, 1956). In a world survey of 560 patients with biopsy-confirmed sarcoidosis Siltzbach (1957) found that the positive Kveim reaction ranged from 65 to 92%.

The onset in most cases was within three years of inoculation. Fried and Genz (1958) have added a further 12 cases (six verified histologically), the ages of the patients at the time of vaccination being between 6 and 10 and at the onset of sarcoidosis between 12 and 21. The time interval after inoculation varied from 18 to 90 months, the average being five years. In our case the interval was seven years.

In the cases of Löfgren and Lundbäck (1952) spontaneous remissions often occurred, but in those of Fried and Genz (1958) only one is recorded. Five of the last-named authors' cases responded partially to antibiotic and hormonal therapy, but some of these might have improved had treatment not been given. In our case the recovery was spontaneous.

Patients with sarcoidosis are unable to develop and maintain skin sensitivity to tuberculin after B.C.G. vaccination (Israel *et al.*, 1950; Forgacs *et al.*, 1957). Many patients with sarcoidosis are known to have been tuberculin-positive prior to the onset of sarcoidosis. The sensitivity wanes and disappears when clinical symptoms of sarcoidosis appear, and when sarcoidosis subsides the sensitivity to tuberculin may return (Nitter, 1953). In some cases the previous history may reveal failure to convert to positive after vaccination (James, 1958). In the cases of Fried and Genz (1958) and in 17 of the 25 cases referred to by Löfgren and Lundbäck (1952) the tuberculin allergy became negative or distinctly weaker. Our patient followed the usual course; she developed positive skin sensitivity after vaccination, which was not maintained during the sarcoidosis, but reverted to positive when she recovered.

Mather *et al.* (1955) and Scadding (1956a) believe there is often a direct aetiological relationship between tuberculosis and sarcoidosis. Those who support the tuberculous aetiology would regard sarcoidosis as an example of tuberculous infection in which there is a high resistance and immunity, but in which immunological response has been associated with marked granulomatous formation (Robb-Smith, 1956). None the less, this combination in the same patient is not often found (Cowdell, 1954). In sarcoidosis the patient appears to have an enhanced capacity to destroy tubercle bacilli in the tissues, which accounts for the inability to identify them in sarcoid follicles. This cannot apply to acid-fast bacilli as a whole, as B.C.G. organisms remain alive at vaccination sites for as long a period in patients with sarcoidosis as in normal susceptible controls (Rostenberg *et al.*, 1953; Forgacs *et al.*, 1957).

It is possible that sarcoidosis and B.C.G. vaccination are coincidental (Törnell, 1954; James, 1958). The incidence of sarcoidosis as compared with tuberculosis is greater in vaccinated than in unvaccinated groups of patients, and the discrepancy is too great to be accidental (Fried and Genz, 1958). The incidence of post-B.C.G. sarcoidosis is higher than the normal morbidity rate for sarcoidosis in Switzerland (Birkhäuser, 1957). It is unlikely that the B.C.G. vaccination precipitates sarcoidosis in view of the long time interval between vaccination and the onset of sarcoidosis in our own and other writers' cases, although in leprosy there can be an interval of up to 20 years between contact and the development of granulomatous formation (Muir, 1948). Löfgren (1953) thinks vaccination may provoke sarcoidosis in patients when the disease is latent. It may well be that without B.C.G. vaccination these cases would normally have developed tuberculosis and that the vaccination protects and modifies the case to one of sarcoidosis. In this country more cases may appear as more school-leavers are vaccinated and followed up for several years. In children under 15 years of age only 113 cases of sarcoidosis had been recorded in a world survey of the literature (McGovern and Merritt, 1956).

Summary

A histologically proved case of sarcoidosis occurred in a girl given B.C.G. vaccination seven years previously. She recovered spontaneously, and the tuberculin sensitivity, which became negative during her illness, reverted to positive. The literature on the occurrence of sarcoidosis in B.C.G. vaccinated cases is reviewed. It is suggested that had she not been vaccinated she might have developed tuberculosis instead of sarcoidosis. The time interval was too long for vaccination to be regarded as the cause of the sarcoidosis.

We thank Dr. T. M. Pollock for kindly referring this case to us, and Dr. Sheila Sherlock and Dr. Geraint James for their help with the liver biopsy and Kveim tests respectively.

REFERENCES

- Berg, S. (1940). *Acta med. scand.*, **103**, 8. Quoted by E. Sommer, *Schweiz. Z. Tuberk.*, **12**, 163.
- Besnier, E. (1889). *Ann. Derm. Syph. (Paris)*, **10**, 333.
- Birkhäuser, H. (1957). *Schweiz. med. Wschr.*, **87**, 1434.
- Boeck, C. (1899). *J. cutan. Dis.*, **17**, 543.
- (1905). *Arch. Derm. Syph. (Wien)*, **73**, 71.
- Cowdell, R. H. (1954). *Quart. J. Med.*, **23**, 29.
- Forgacs, P., McDonald, C. K., and Skelton, M. O. (1957). *Lancet*, **1**, 188.
- Fried, K. H., and Genz, H. (1958). *Tuberk-Arzt.*, **12**, 558.
- Ganguin, H. G. (1956). *Allergie u. Asthma*, **2**, 88.
- Hutchinson, J. (1875-8). *Illustrations of Clinical Surgery*, **1**, 39. London.
- (1898). *Arch. Surg. (Lond.)*, **9**, 307.
- Israel, H. L., Sones, M., Stein, S. C., and Aronson, J. D. (1950). *Amer. Rev. Tuberc.*, **62**, 408.
- James, D. G. (1956). *Brit. med. J.*, **2**, 900.
- (1958). *Postgrad. med. J.*, **34**, 240.
- Jüngling, O. (1919). *Fortschr. Röntgenstr.*, **27**, 375.
- Larsen, A. K. (1950). *Nord. Med.*, **43**, 170.
- Löfgren, S. (1953). Quoted by Fried and Genz (1958).
- and Lundbäck, H. (1952). *Acta med. scand.*, **142**, 265.
- McGovern, J. P., and Merritt, D. H. (1956). *Advanc. Pediat.*, **8**, 97.
- Mather, G., Dawson, J., and Hoyle, C. (1955). *Quart. J. Med.*, **24**, 331.
- Muir, E. (1948). *Manual of Leprosy*, p. 10. Livingstone, Edinburgh.
- Nitter, L. (1953). *Acta radiol. (Stockh.)*, Suppl. 105.
- Pfisterer, R., Wespi, H., and Herzog, H. (1954). *Helv. med. Acta*, **21**, 439.
- Robb-Smith, A. H. T. (1956). *Proc. roy. Soc. Med.*, **49**, 1041.
- Rostenberg, A., Szymanski, F. J., Brebis, G. J., Haeberlin, J. B., and Seneear, F. E. (1953). *A.M.A. Arch. Derm. Syph.*, **67**, 306.
- Scadding, J. G. (1950). *Brit. med. J.*, **1**, 745.
- (1956a). *Proc. roy. Soc. Med.*, **49**, 799.
- (1956b). *Tubercle (Lond.)*, **37**, 371.
- Schaumann, J. (1914). *Sur le lupus pernio*. Paris.
- Siltzbach, L. E. (1957). *Amer. J. Med.*, **22**, 841.
- Törnell, E. (1954). *Nord. Med.*, **52**, 996.

CONTROLLED HYPOTHERMIA IN INFANTS AND CHILDREN

BY

DOUGLAS COHEN, M.S., F.R.A.C.S.

*Institute of Child Health and Congenital Heart Disease Unit,
Royal Alexandra Hospital for Children, Sydney*

AND

VICTOR HERCUS,* M.B., M.R.C.P.Ed., D.A.

*Unit of Clinical Investigation, Royal North Shore Hospital,
Sydney, and Congenital Heart Disease Unit,
Royal Alexandra Hospital for Children, Sydney*

The use of hypothermia now occupies a definite though restricted place in medicine. When we know more concerning oxygen transfer to the body cells, and its utilization by them during hypothermia, many of the present restrictions governing the use of lowered body temperatures may be removed, and the indications for its use may be extended considerably.

In cardiac surgery in children, when hypothermia is utilized, various technical requirements are desirable, the essential elements of which are:

1. A central body temperature which can be maintained accurately at any desired level, usually 30° C. This is estimated by the temperature recorded in the lower third of the oesophagus. Cooper and Kenyon have shown this to be a fairly accurate assessment of the central blood stream temperature during surface cooling, and we have also found that this temperature corresponds very closely to the temperature of the cerebral cortex.
2. A reasonable rate of fall in body temperature. This should be rapid enough to avoid the necessity for any extension of the normal operating-time.
3. Means whereby the body temperature may conveniently be raised or lowered while surgery is in progress.
4. Restoration of the body temperature to normal, as soon as possible after the completion of the operation, with minimal handling of the patient.
5. Accurate control of the rate of change of body temperature and facilities to stabilize it rapidly at any desired level, thus preventing any "overswing."

*Aided by a grant from the Royal Australasian College of Physicians.