participants were approved by the ethical committee of each participating centre. The patients with treatment failure received triple therapy at any time during the study if they asked for it, or at the end of the study if they did not wish to be enrolled in the follow-up study. Those who were enrolled in the second study received triple therapy at the end of follow-up or if the ulcer recurred. Furthermore, during the trial, we used a dual therapy with amoxicillin and not with clarithromycin so that in the case of treatment failure, clarithromycin-based triple therapy could be subsequently used without the risk of acquired resistance to clarithromycin and consequent reduced efficacy.¹

Dr Graham and Dr Yamaoka also suggest that rather than focusing on the similar efficacy of 1 and 2 weeks of triple therapy, we should have highlighted that the efficacy of this therapy is "unacceptably low", regardless of its duration. As we discuss in our paper, the eradication rates obtained in our study (1 week of triple therapy intent to treat 79.7%, per protocol 83.6%; 2 weeks of triple therapy intent to treat 81.7%, per protocol 84.9%) add to the body of evidence showing that the efficacy of triple therapy has undoubtedly decreased in recent years. However, these are not substantially below the threshold of 80% defined by the Maastricht III Consensus Report as clinically unacceptable for the treatment of *H. pylori* infection.² Moreover, triple therapy, referred to by Dr Graham and Dr Yamaoka as a "legacy" treatment, is the treatment recommended by the Maastricht III Consensus Report. Dr Graham is himself one of the authors of this report. We must agree with this recent recommendation, at least until the higher efficacy demonstrated for new treatments³ has been confirmed by other investigators and in large studies in different regions worldwide.

We argue that the importance of our study lies rather in its large sample size and demonstration that extending the duration of triple therapy from 1 to 2 weeks does not significantly increase the rate of *H. pylori* eradication (the difference in eradication rate between 1 and 2 weeks of triple therapy was 2%; 95% CI -4% to 8%). Our trial involved a large number of centres (n = 81) in the south, centre and north of Italy, and so patients were recruited from a broad population, providing a good basis for the generalisation of results. Before our study, two meta-analyses reported that compared with 1 week of treatment, 2 weeks of triple therapy increases the eradication rate by 9–12%.^{4,5} However, although the results of meta-analyses are relevant, they cannot substitute for large clinical trials. Indeed, in their meta-analysis, Calvet *et al.*⁴ specifically mentioned that their results suggest that larger studies comparing 1 and 2 weeks of therapy are needed. We think that our study has addressed this need.

Rocco Maurizio Zagari, Franco Bazzoli

Department of Internal Medicine and Gastroenterology,
University of Bologna, Bologna, Italy

Correspondence to: Professor F Bazzoli, Dipartimento di Medicina Interna e Gastroenterologia, Università di Bologna, Policlinico S Orsola, Via Massarenti 9, 40138 Bologna, Italy; franco.bazzoli@unibo.it

Competing interests: RMZ has received reimbursement from Janssen-Cilag and Abbot for attending symposia, and fees for speaking from AstraZeneca, Takeda and Abbott. FB has received fees for speaking from AstraZeneca and Takeda, reimbursement for organising education from Altana and has carried out consultancy work for AstraZeneca.

References

- 1 Bochenek WJ, Peters S, Fraga PD, *et al.* Eradication of *Helicobacter pylori* by 7-day triple therapy regimens combining pantoprazole with clarithromycin, metronidazole, or amoxicillin in patients with peptic ulcer disease: results of two double-blind, randomized studies. *Helicobacter* 2003;**8**:626–42.
- 2 Malfertheiner P, Megraud F, O'Morain C, *et al.* Current concepts in the management of *Helicobacter pylori* infection – the Maastricht III Consensus Report. *Gut* 2007;**56**:772–81.
- 3 De Francesco V, Zullo A, Margiotta M, *et al.* Sequential treatment for *Helicobacter pylori* does not share the risk factors of triple therapy failure. *Aliment Pharmacol Ther* 2004;**19**:407–14.
- 4 Calvet X, Garcia N, Lopez T, *et al.* A meta-analysis of short versus long therapy with a proton pump inhibitor, clarithromycin and either metronidazole or amoxicillin for treating *Helicobacter pylori* infection. *Aliment Pharmacol Ther* 2000;**14**:603–9.
- 5 Ford A, Moayyedi P. How can the current strategies for *Helicobacter pylori* eradication therapy be improved? *Can J Gastroenterol* 2003;**17**(Suppl B):36B–40B.

Risk of cancer and acid suppressant treatment

Garcia Rodriguez *et al.* (*Gut* 2006;**55**:1538–44) use data from the General Practitioner Research Database (GPRD) to report that the odds ratios (ORs) for incidences of non-cardia gastric cancer were increased fourfold in those who received acid suppressant treatment for ulcer indications, compared with those who did not, and fivefold for oesophageal adenocarcinoma in those with oesophageal indications. They consider that the increased risks reflect the indications for treatment and not drug toxicity. Although their conclusions may be right, their data raise some important concerns.

The GPRD has particular strengths in recording prescriptions, but case details can be less complete, which may explain why 482 of the oesophageal cancer cases were excluded because of unspecified histology, leaving 287 with adenocarcinoma for comparison with their controls without cancer. The assumption that these 287 were representative of other unidentified adenocarcinoma cases may or may not be sound.

Secondly, their paper suggests that gastro-oesophageal reflux or hiatal hernia is associated with this increased risk of oesophageal cancer, a conclusion that, if accepted, would result in radical reappraisal of current attitudes limiting cancer surveillance to patients with Barrett's oesophagus. We previously published data, 4 years ago in *Gut*¹ based on a surveillance of 18 000 patients who were prescribed omeprazole, and compared death rates from the main individual causes with those found nationally. Their dataset and ours were obtained by different methods, but they are, or should be, complementary, giving information on the risk of developing cancer and mortality ascribed to cancer (ours). We found 27 deaths from oesophageal cancer in 4 years of follow-up in 5517 patients with Barrett's disease, oesophageal stricture, ulcer or inflammation (OR 3.30 (95% CI 2.17 to 4.80)). By contrast, they found 16 cases among long-term users of acid suppressants, with all oesophageal indications (OR 5.42 (95% CI 3.13 to 9.29)). This OR is just compatible with ours of 3.30 in a restricted subset with severe oesophageal disease, but must be markedly higher if risk is, in fact, concentrated in a particular

subset with severe disease, such as that chosen by us. Table 5 in the paper indicates that only one of their oesophageal adenocarcinoma cases received long-term acid suppression for reflux as the sole indication, making it difficult to conclude that reflux alone is a risk factor, a finding consonant with ours, where 6 oesophageal cancer deaths were found in 6984 patients with hiatal hernia or reflux only OR 1.02 (95% CI 0.37 to 2.22).

Finally, they conclude that patients with ulcer indications treated by acid suppression have raised risks of gastric cardia and, particularly, non-cardia cancers. These findings are based on data from 522 gastric cancer cases, with 501 others excluded from consideration because gastric subsites were unclear. Table 7 shows that the raised ORs are derived from 3 cardia cases with ulcer indications, and 11 with non-cardia cancer, the OR for the cardia cancer cases being compatible with unity (OR 2.18, 95% CI 0.67 to 7.03). Retrospective consideration is always beguiling, particularly when derived from subset studies, and before accepting these, would it not have been sensible to compute risks for all gastric cancers, irrespective of subsite? We found that the chances of death due to gastric cancer fell to the population expectation after 2 years of follow-up in our total series of 51 cases, suggesting that any excess of earlier deaths in those who take drugs was likely to reflect acid suppressants prescribed for patients who already had gastric cancer, a conclusion consonant with results obtained in similar studies of cimetidine and published over 20 years ago.²

Garcia Rodriguez *et al.* agree with our earlier conclusion that confounding by treatment indication explains apparent increased risks of upper digestive cancer in those who receive acid suppressant treatment. However, their conclusions from data exploring oesophageal cancer risks associated with reflux and hiatal hernia, and gastric subsite risks with ulcer indications, seem insecure.

Michael Langman

Division of Medical Sciences, Medical School, University of Birmingham, Queen Elizabeth Hospital, Birmingham, UK

Richard Logan

Division of Public Health and Epidemiology, University of Nottingham Medical School, Queens Medical Centre, Nottingham, UK

Correspondence to: Professor M Langman, Division of Medical Sciences, Medical School, University of Birmingham, Queen Elizabeth Hospital, Birmingham B15 2TH, UK; m.j.s.langman@bham.ac.uk

Competing interests: None.

References

- 1 Bateman DN, Colin Jones D, Hartz S, *et al.* Mortality study of 18000 patients treated with omeprazole. *Gut* 2003;**52**:942–6.
- 2 Colin Jones DG, Langman MJ, Lawson DH, *et al.* Postmarketing surveillance of the safety of cimetidine: 12 month mortality report. *BMJ* 1983;**286**:1713–16.

Non-invasive measurement of small-bowel motility by MRI after abdominal surgery

Abdominal surgery under general anaesthesia causes postoperative ileus¹ with an extended recovery time.² Prolonged ileus creates major healthcare problems costing an estimated US\$

Figure 1 Repetitive acquisition of coronal two-dimensional balanced turbo field echo datasets of the same slice with 0.25 s acquisition time. Measurements of cross-sectional diameters of bowel. (A–C) Magnified image of a preterminal ileal segment showing (A) normal width, (B) contraction and (C) subsequent relaxation.

750 million per annum (£382 million, €563 million) in the US.³ Small-bowel motility is currently assessed by manometry, small-bowel transit time of radio-opaque markers and scintigraphy.⁴ All these methods are invasive (tube, radio-opaque markers or radio-labelled meal), unpopular with patients and restricted in their application to a few academic centres. Ultrafast magnetic resonance imaging (MRI) allows dynamic analysis of moving structures under physiological conditions, as reported for other organs.^{5,6}

In this prospective clinical study, 30 patients were examined 3–5 days after colorectal surgery (sigmoidectomy ($n=20$), diverticulitis or carcinoma; rectal resection ($n=8$),

carcinoma). Surgeries were performed under general anaesthesia combined with peridural block analgesia. For MRI (1.5 T), no oral preparation was administered to the patients. A coronal three-dimensional sequence covering the abdomen was acquired (turbo field echo (TFE), TR4.0, TE1.5, 80 slices, slice thickness = 1.5 mm). Small-bowel segments easily visible on the above sequence were subjected to a dynamic breath-hold coronal two-dimensional sequence (balanced TFE, TR2.7, TE1.3, matrix = 192×512 , slice-thickness = 10 mm) consisting of a single 10 mm thick slice repeated every 0.25 s for 17.5 s (fig 1).

To assess the motility, cross-luminal diameters of selected small-bowel segments were

measured on each of the dynamic images. The diameters were plotted over time and contraction-frequencies were assessed. The motility patterns were visually evaluated based on diameter, amplitude and frequency characteristics.

All 30 patients underwent MRI 4 (SD 1) days after surgery and dynamic sequences were successfully acquired in each. The mean diameter of the measured small-bowel segments was 15.9 (SD 2.5) mm (range 1–38.3 mm) and the mean amplitude was 10 (SD 4.7) mm (range 2–22.7 mm). Four clearly distinguishable motility patterns could be discerned (fig 2):

- (1) paralysis: dilated lumen, absent amplitudes, low contraction frequencies;
- (2) shivering: low amplitudes, high contraction frequencies;
- (3) uncoordinated: varying diameters, amplitudes and frequencies;
- (4) sinusoidal: regular amplitudes, constant contraction frequencies.

A statistically significant increase in uncoordinated and sinusoidal motility patterns was encountered during convalescence.

Assessment of small-bowel motility is still a largely unexplored field.⁷ MRI allows the static display of morphology and also the dynamic analysis of bowel motion using ultrafast scanning techniques.⁸ Our study demonstrates that MRI of small-bowel motility shortly after abdominal surgery is both feasible and highly

Figure 2 The four small-bowel motility patterns in patients examined 3–5 days after colorectal resection surgery. Measurements of cross-sectional diameters of the small-bowel loops plotted over time. (A) Paralysis: dilated lumen, absent amplitudes, low contraction frequencies; (B) shivering: low amplitudes, high contraction frequencies; (C) uncoordinated: varying diameters, amplitudes, frequencies; and (D) sinusoidal: regular amplitudes, constant contraction frequencies.

effective. To our knowledge, this study is the first to describe the non-invasive visualisation of small-bowel motility in a patient group.

The reduced distension did not impede the visualisation of the small-bowel wall as it was outlined by either air or gastrointestinal liquid. It had the added advantage of not affecting the bowel function by placing tubes or filling the bowel. Four distinct motility patterns could be distinguished in our patients: paralysis, shivering, uncoordinated motion and sinusoidal motion. A previous study on motility with healthy volunteers⁹ observed only the sinusoidal pattern, suggesting a clear difference in the motility patterns between postoperative patients and healthy volunteers. It may be hypothesised that motility recovers along a defined path during postoperative convalescence starting with slight shivering, changing into irregular movements and finally developing into full sinusoidal motion. Many factors may influence postoperative bowel motility, but the effect and relevance of each are complex and remain to some extent unknown.^{9,10}

The MRI technique described here is not limited to postoperative treatment, but can be applied to a wide range of motility diseases⁷ and common intestinal disorders with unknown pathomechanisms. Non-invasive MRI assessment of peristaltic motion allows rapid and reliable localisation and diagnosis of motility problems and thus could have a beneficial impact on treatment and recovery.

Michael A Patak

Institute of Diagnostic Radiology, Inselspital, University Hospital, Bern, Switzerland

Michael A Patak, Johannes M Froehlich, Constantin von Weymarn

MR-Research Group, Institute of Radiology, Cantonal Hospital, Winterthur, Switzerland

Stefan Breitenstein

Department of Surgery, University Hospital Zurich, Switzerland

Christoph L Zollkofer, Klaus-Ulrich Wentz

MR-Research Group, Institute of Radiology, Cantonal Hospital, Winterthur, Switzerland

Correspondence to: Dr M A Patak, Institute of Diagnostic Radiology, Inselspital, University Hospital, 3010 Bern, Switzerland; michael.patak@insel.ch

doi: 10.1136/gut.2007.120816

Competing interests: None.

References

- Behm B, Stollman N. Postoperative ileus: etiologies and interventions. *Clin Gastroenterol Hepatol* 2003;1:71–80.
- Waldhausen JH, Shaffrey ME, Skenderis BS II, et al. Gastrointestinal myoelectric and clinical patterns of recovery after laparotomy. *Ann Surg* 1990;211:777–84; discussion 785.
- Moss G, Regal ME, Lichtig L. Reducing postoperative pain, narcotics, and length of hospitalization. *Surgery* 1986;99:206–10.
- Camilleri M, Hasler WL, Parkman HP, et al. Measurement of gastrointestinal motility in the GI laboratory. *Gastroenterology* 1998;115:747–62.
- Schwizer W, Fox M, Steingotter A. Non-invasive investigation of gastrointestinal functions with magnetic resonance imaging: towards an "ideal" investigation of gastrointestinal function. *Gut* 2003;52(4 Suppl):iv34–9.
- Paetsch I, Jahnke C, Wahl A, et al. Comparison of dobutamine stress magnetic resonance, adenosine stress magnetic resonance, and adenosine stress

magnetic resonance perfusion. *Circulation* 2004;110:835–42.

- Jones MP, Wessinger S. Small intestinal motility. *Curr Opin Gastroenterol* 2006;22:111–16.
- Froehlich JM, Patak MA, von Weymarn C, et al. Small bowel motility assessment with magnetic resonance imaging. *J Magn Reson Imaging* 2005;21:370–5.
- Schnoor J, Unger JK, Kuepper T, et al. Effects of propofol and fentanyl on duodenal motility activity in pigs. *Can Vet J* 2005;46:995–1001.
- Husebye E. The patterns of small bowel motility: physiology and implications in organic disease and functional disorders. *Neurogastroenterol Motil* 1999;11:141–61.

Double resistance to imatinib and AMG 706 caused by multiple acquired *KIT* exon 17 mutations in a gastrointestinal stromal tumour

Gastrointestinal stromal tumours (GISTs) are rare mesenchymal neoplasms that arise in the wall of the gastrointestinal tract and originate from the interstitial cells of Cajal.¹ The tumours are characterised by the expression of the receptor tyrosine kinase *KIT* (CD117, c-Kit). Activating mutations of the *KIT* proto-oncogene are known to be associated with the tumorigenesis of most GIST.² The standard treatment for primary GIST is surgical resection, as GISTs are resistant to conventional

chemotherapy. The multitargeted tyrosine-kinase inhibitor (MKI) imatinib mesylate (Gleevec, Novartis, Basel, Switzerland) revolutionised the treatment for advanced disease. Although most patients benefit from imatinib, many subsequently develop acquired resistance.³ In patients developing resistance to imatinib, other MKIs provide clinical benefit. Sunitinib malate (Sutent, Pfizer, New York, USA) has demonstrated activity in patients with imatinib-resistant GIST,⁴ and other MKIs are currently in development, including AMG 706, an inhibitor of *KIT*, all vascular endothelial growth factor receptors and platelet-derived growth factor receptor. Preliminary data from recent clinical trials with AMG 706 showed an encouraging clinical benefit rate in patients with advanced high-dose imatinib-resistant GIST.⁵

We report on a 64-year-old woman with a 20 mm GIST of the small bowel (fig 1A, table 1), which was removed in September 2002.

In October 2003, peritoneal and liver metastases were recognised and treatment with imatinib (400 mg/day) was started, resulting in a complete metabolic response (on [¹⁸F] fluorodeoxyglucose positron emission tomography) 8 weeks later. Owing to the presence of a subsequent progressive mesenteric mass, the imatinib dose was increased (800 mg/day) in July 2004, leading to growth arrest. In April

Figure 1 (A) Histopathology of the high mitotic primary spindle cell type gastrointestinal stromal tumour (arrows: mitoses, H&E $\times 400$). (B) *KIT* exon 17 point mutation c.2466T→A (G) found in three specimens of the imatinib/AMG 706-resistant progressive tumour leading to Asn822Lys. (C) *KIT* exon 17 point mutations c.2453A→G, c.2467T→G, and c.2433T→C (silent) found in additionally investigated four specimens of the tumour relapse leading to Tyr823Asp and Lys818Arg.