ination showed, dense eosinophilic infiltration (>20 cells/mm³). She had never experienced dysphagia, vomiting or refusal of food. There was no allergy to inhaled or alimentary allergens. Serum IgE level was normal (54 IU/l) and skin tests for dust mite, graminaceae and common alimentary allergens (lactoproteins, ovalbumin, grain, rice and soja) were negative. A gluten-free diet was prescribed, but the girl admitted to not have followed it properly. After 1 year, EMAs were still weakly positive with persistent mucosal atrophy in the duodenum. At macroscopical and histological examination, EE remained almost unvaried, but the girl still had no EE-suggestive symptoms.

Case 3

A 19-year-old asymptomatic sister of a girl with CD, with positive DQ2 HLA and positive EMAs and tTG antibodies, underwent endoscopy to confirm CD. Endoscopy did not show macroscopic duodenal inflammation (but duodenal biopsy showed lymphocytic infiltration and initial mucosal atrophy), whereas on distal oesophageal mucosa white small plaques were reported, with bioptical examination revealing lymphocytic and granulocytic infiltration in the distal oesophagus, with eosinophilic count varying from 50 to 100/HPF. CD and EE were therefore diagnosed. Skin tests for common inhaled and alimentary allergens were negative, whereas only the patch test for nickel sulphate was positive. She was then prescribed a gluten-free diet and omeprazole for 3 months. A further clinical, endoscopic, bioptical and laboratory assessment was performed 7 months later. The girl was still asymptomatic, and she admitted to properly following the diet (except for possible external contaminations). Nonetheless, EMA and tTG were still weakly positive. Endoscopic examination showed a granular aspect of the distal oesophageal mucosa but no other macroscopic alterations. Biopsy specimens revealed lymphocytic epithelial infiltration with slightly dismorphic villi of the duodenal mucosa and an important eosinophilic count (90/HPF) in the distal oesophageal mucosa. Skin tests were repeated and tested negative, as were the radioallergosorbent tests for specific allergens. The histopathological diagnosis was of CD in infiltrative stadium and active EE. As the girl was asymptomatic, no specific EE treatment was started.

Conclusions and comments

The pathogenetic mechanisms that lead to EE are far from being understood. However, among paediatric patients, a large proportion of cases seem to be associated with food allergies and to respond to a specific elimination diet.⁴ High levels of circulating eosinophilic cells and IgE are usually found in patients affected by CD, thus supporting the hypothesis of an atopic allergic reaction as a possible pathophysiological trigger. Moreover, EE is often associated with conditions such as allergic rhinitis, asthma, atopic dermatitis and familiar history of atopy.^{1,3} Unlike most of better-known atopic syndromes, it is usually very difficult to identify the cause leading to EE (as non-food items are often involved); therefore immune modulating therapy (topical steroids, leucotrienes) is generally preferred to avoidance strategies (elementary or elimination diets), at least in the absence of indicative prick or radioallergosorbent test.

This is the first report of the association between EE and CD. Regardless of the intrinsic limitations of any isolated reports, our three cases suggest that EE may be a gluten-associated manifestation in subjects with CD. Between 2000 and 2006, 7440 endoscopies were performed at the Institute of Child Health, IRCCS Burlo Garofolo, Trieste, Italy, and at the Surgical and Endoscopic Digestive Unit, Ospedale Paediatrics, Bambino Gesù, Rome, Italy, leading to the diagnosis of 1919 cases of CD and 33 cases of EE. The prevalence of CD among patients with EE was 3 of 33 (9%), nine times higher than that expected in the general population (1:100).⁵ By contrast, the prevalence of EE among patients with CD was 3 of 1919 (1:640, 0.0015%), a 3.5 times higher prevalence than that calculated in a recent American study (1:2380, 0.00042%).³ In only one of the three cases that we have described, EE disappeared after the gluten-free diet, suggesting that EE may be a gluten-dependent—even if rare—manifestation of CD. In the two other cases, the gluten-free diet did not have any effect on the eosinophilic infiltrate but both girls were not compliant to the gluten-free diet. The association between EE and CD may be plausible, thus suggesting that one performs a duodenal biopsy in all patients with EE in association with anti-tTG and EMAs dosage.

These three reports further underline how EE may be completely asymptomatic even when endoscopic and histological diagnostic features are evident. In our clinical experience, some patients treated with either topical steroids and/or elimination diets became quickly asymptomatic while their endoscopic and histological alterations persisted (data not published). The possibility that EE may be completely asymptomatic raises the still open questions on its natural history, the goals and the best duration of treatment of this emerging clinical condition.

F Verzeegnassi, J Bua

Department of Paediatrics, Institute of Child Health IRCCS Burlo Garofolo, University of Trieste, Trieste, Italy

P De Angelis, L Dall'Oglio

Surgical and Endoscopic Digestive Unit, Ospedale Pediatrico Bambino Gesù, Rome, Italy

G Di Leo, A Ventura

Department of Paediatrics, Institute of Child Health IRCCS Burlo Garofolo, University of Trieste, Trieste, Italy

Correspondence to: Dr J Bua, Department of Paediatrics, Institute of Child Health, IRCCS Burlo Garofolo, Via dell'Istria 65/1, 34100 Trieste, Italy; jennybua@gmail.com

doi: 10.1136/gut.2006.117986

Competing interests: None.

Informed consent has been obtained for publication of the patients' details in this letter.

References

- Arora AS, Yamazaki K. Eosinophilic esophagitis: asthma of the esophagus? *Clin Gastroenterol Hepatol* 2004;**2**:523–30.
- Penttiuk SP, Miller CK, Kaul A. Eosinophilic esophagitis in infants and toddlers. *Dysphagia* 2007;**22**:44–8.
- Noel RJ, Putman PE, Rothenberg ME. Eosinophilic esophagitis. *N Engl J Med* 2004;**351**:940–1.
- Spergel JM, Beausoleil JL, Mascarenhas M, et al. The use of skin prick tests and patch tests to identify causative foods in eosinophilic esophagitis. *J Allergy Clin Immunol* 2002;**141**:576–81.

- Tommasini A, Not T, Kiren V, et al. Mass screening for coeliac disease using antihuman transglutaminase antibody assay. *Arch Dis Child* 2004;**89**:512–15.

Disease progression and liver cancer in the ferroportin disease

Ferroportin-associated iron overload (also known as the ferroportin disease) was clinically recognised in 1999,¹ and linked in 2001 to the A77D mutation of ferroportin (FPN).² Owing to the mild clinical expressivity reported in the literature, doubts have been raised on the penetrance of the genetic defect and the rationale for iron-removal therapy.³ However, so far, no prospective data have been collected pertaining to this disorder.

We have had the opportunity to study six members of a pedigree carrying the A77D mutation of FPN for 11–24 years in whom the disease was first described.^{1,2}

The proband (subject VI₃₃, table 1) was diagnosed at the age 59 years. After being on weekly phlebotomy for 2 years, he was on a maintenance programme for 18 years (one phlebotomy every 3 months). At the age of 79 years, he refused to continue on phlebotomy. In 2006, at the age of 83 years, hepatic ultrasound (US) examination of the proband identified two hyperechoic nodules in the liver; contrast-enhanced MRI scan confirmed the US examination findings and a US-guided liver biopsy documented a well-differentiated hepatocellular carcinoma (fig 1A), as shown by antihepatocyte antibody stain (not shown here), in the absence of cirrhosis. Strong p53-positive stain was also detected in the tumour specimen by using a specific antibody (fig 1B). The level of serum α fetoprotein was 5.0 ng/ml (normal value <10 ng/ml). The patient reported no alcohol intake for the past two decades. No antibodies against hepatitis C virus and hepatitis B virus (HBV) were found. However, HBV sequences corresponding to all the four different viral genomic regions tested were detected by nested PCR in the tumour biopsy specimen. In addition, HBV DNA was also found in the liver specimens collected at the age 59 and 79 years. The patient's liver nodules were treated by chemoembolisation.

All the proband's relatives, underwent regular phlebotomy, except for two who showed low adherence and, after reaching iron depletion, underwent phlebotomy occasionally, usually once a year (subject VI₃₃ and VI₃₆; table 1). Both patients presented signs of the progression of fibrosis as compared with the first assessment reported in 1999¹ (fig 1C,D), with appreciable increase in portal blood flow in subject VI₃₃ at colour Doppler US examination (velocity 17 cm/s), and liver iron content more than twofold the normal value.

In conclusion, the ferroportin disease may progress if not properly treated, and complicate with liver cancer even without cirrhosis, in the presence of cocarcinogenic factors, such as occult HBV infection. The hallmark of the classic FPN disease is marked iron accumulation in Kupffer cells.⁴ Kupffer cells are vital in producing fibrogenic mediators,⁵ in immunological tumour surveillance and disposal of transformed hepatocytes.^{6,7} Selective and massive iron overload might impair these activities and favour fibrogenesis and carcinogenesis. Macrophage iron overload is very resistant to iron withdrawal in this disorder, even in patients who are apparently well treated (fig 1E,F). As for the development of liver

Figure 1 Histological and imaging data from affected family members. (A,B) Specimens from the proband's tumour. (A) A well-differentiated hepatocellular carcinoma, intratumour fibrous tissue entrapping iron-laden non-parenchymal cells and necrosis areas. (B) In the nodule, most cells stain positive for p53 antibody (brown nuclei). Sirius red stain shows various degrees of fibrosis in (C) subject V₁₃₆ and (D) subject V₁₃₃. (E,F) MRI scans. T2-weighted gradient-echo sequences were used to detect iron accumulation. Owing to residual iron accumulation, significant decrease in MRI signal is still appreciable in the spleen and the spine of a patient who showed low compliance to therapy (E; subject V₃₃; and also, although to a lesser extent, in a well-treated subject with normal liver iron content (F; subject V₃₀).

cancer in the proband, we speculate that adequate phlebotomy "controlled" the oncogenic potential of HBV (which was present in the patient's liver since his first hospital admission at the age of 59 years). However, once the 20-year-long treatment was discontinued, the underlying hepatitis B infection could express its cofactorial oncogenic activity and contribute, with the rising iron burden, to

the development of neoplastic nodules. In this vein, iron-dependent loss in p53 growth-suppressor activity, also implicated in liver carcinogenesis in classic haemochromatosis,⁸ may have had a role and facilitated the HBV oncogenic potential.⁹ Our results prompt for a thorough lifelong maintenance therapy and regular screening for liver cancer in the patients with ferroportin disease.

Elena Corradini, Francesca Ferrara

Department of Internal Medicine, Center for Hemochromatosis, University Hospital of Modena, Modena, Italy

Teresa Pollicino

Department of Internal Medicine, Unit of Clinical and Molecular Hepatology, University Hospital of Messina, Messina, Italy

Alberto Vegetti, Gian Luca Abbati

Department of Internal Medicine, Center for Hemochromatosis, University Hospital of Modena, Modena, Italy

Luisa Losi

Department of Anatomy and Pathology, University Hospital of Modena, Modena, Italy

Giovanni Raimondo

Department of Internal Medicine, Unit of Clinical and Molecular Hepatology, University Hospital of Messina, Messina, Italy

Antonello Pietrangelo

Department of Internal Medicine, Center for Hemochromatosis, University Hospital of Modena, Modena, Italy

Correspondence to: Professor A Pietrangelo, Department of Internal Medicine, Center for Hemochromatosis, Policlinico, Via del Pozzo 71 41100 Modena, Italy; pietrangelo.antonello@unimo.it

doi: 10.1136/gut.2007.122549

Competing interests: None.

References

- 1 Pietrangelo A, Montosi G, Totaro A, *et al*. Hereditary hemochromatosis in adults without pathogenic mutations in the hemochromatosis gene [see comments]. *N Engl J Med* 1999;**341**:725–32.
- 2 Montosi G, Donovan A, Totaro A, *et al*. Autosomal-dominant hemochromatosis is associated with a mutation in the ferroportin (SLC11A3) gene. *J Clin Invest* 2001;**108**:619–23.
- 3 Devalia V, Carter K, Walker AP, *et al*. Autosomal dominant reticuloendothelial iron overload associated with a 3-base pair deletion in the ferroportin 1 gene (SLC11A3). *Blood* 2002;**100**:695–7.
- 4 Pietrangelo A. The ferroportin disease. *Blood Cells Mol Dis* 2004;**32**:131–8.
- 5 Friedman SL. Mechanisms of disease: mechanisms of hepatic fibrosis and therapeutic implications. *Nat Clin Pract Gastroenterol Hepatol* 2004;**1**:98–105.
- 6 Manifold IH, Triger DR, Underwood JC. Kupffer-cell depletion in chronic liver disease: implications for hepatic carcinogenesis. *Lancet* 1983;**2**:431–3.
- 7 Zhang W, Arai S, Sasaaki T, *et al*. The role of Kupffer cells in the surveillance of tumor growth in the liver. *J Surg Res* 1993;**55**:140–6.
- 8 Hussain SP, Raja K, Amstad PA, *et al*. Increased p53 mutation load in nontumorous human liver of Wilson disease and hemochromatosis: oxyradical overload diseases. *Proc Natl Acad Sci USA* 2000;**97**:12770–5.

Table 1 Biochemical and clinical data of the subjects with ferroportin-related iron overload

No	Subjects	Age at diagnosis (years)*	Years on phlebotomy	Serum ferritin level at present, (µg/l)	Liver iron concentration at present, (µmol/g)	Liver fibrosis at diagnosis	Liver fibrosis at present
1	V ₁₃	59	20	690	55	F1	F2
2	V ₁₂₅	34	15	650	35	F0	F0
3	V ₁₂₆	20	11	670	35	F0	F0
4	V ₁₃₀	37	12	410	28	F0	F0
5	V ₁₃₃	39	11	827	75	F0	F3
6	V ₁₃₆	34	15	750	65	F0	F2

Patient numbering is the same as that used in the original publication¹.

*All patients were male.

9 **Puisieux A**, Ponchel F, Ozturk M. p53 as a growth suppressor gene in HBV-related hepatocellular carcinoma cells. *Oncogene* 1993;8:487–90.

Which imaging modalities should be used for biliary strictures of unknown aetiology?

Why is it so difficult to determine the nature of isolated biliary strictures? In patients with no history of biliary surgery, most bile duct strictures have to be considered malignant until proven otherwise. This usually leads to extensive usage of diagnostic imaging—in many cases even to explorative laparotomy. Furthermore, we usually have a “luxury problem”: doctors have to choose the appropriate imaging technique out of the wide range of endoscopic and radiological modalities. For this purpose, doctors should have information about each technique. In a multimodal approach, we prospectively analysed sensitivity and specificity for differentiation between malignant and benign strictures, penetration into surrounding tissues, vascular invasiveness and staging of lymph node.

We included patients with jaundice of unknown aetiology and suspected biliary stricture. According to the study protocol, patients underwent diagnostic laparotomy; alternatively, a long follow-up of more than 12 months was accepted. Patients were not included in case of metastases. We examined patients applying endoscopic retrograde cholangiopancreatography (ERCP), intraductal ultrasonography (IDUS), endoscopic ultrasound (EUS), multidetector CT (MDCT) and MRI.

In our study, a total of 30 patients were enrolled. According to the histopathological results following laparotomy, we found 21 (70%) patients with malignancies. For analysis of penetration into the surrounding tissue, vascular infiltration and lymph node invasiveness, we included only patients who underwent surgery as gold standard ($n=28$). ERCP and IDUS, EUS and MDCT proved to be significantly superior in differentiating benign from malignant strictures as compared with MRI (table 1). Penetration into the surrounding tissue was best shown by ERCP and IDUS, with a sensitivity of 88% (95% CI 64% to 99%), which was significantly better than MRI (24%, 95% CI 7% to 50%). With respect to vascular infiltration and lymph node invasiveness, we found only a slight difference in favour of MDCT compared with the other techniques.

In 2002, Rösch *et al.*¹ performed the first study with a multimodal approach. Since imaging modalities have clearly improved, there is a strong need for further prospective studies assessing different imaging methods adapted to the present state-of-the-art. According to our data, differentiation between malignant and benign strictures is most suitable by ERCP and IDUS. Compared with MRI, all three imaging techniques proved to be significantly superior in their ability to define potential malignancy of biliary strictures. Furthermore, ERCP supplemented by IDUS showed a high sensitivity rate—with a significant difference compared with MRI—in the assessment of tumour penetration into the surrounding tissue. With respect to vascular infiltration and lymph node invasiveness, no statistically significant difference was documented between the applied methods.

The development and improvement in MRI technology led to various assessments by

Table 1 Ability of diagnosing malignancy in biliary strictures of unknown aetiology

	Sensitivity (%)	p<0.05	Specificity (%)	p<0.05
ERCP and IDUS	95	0.02	89	NS
EUS	81	0.039	89	NS
MDCT	76	0.041	78	NS
MRI	48		78	NS

ERCP, endoscopic retrograde cholangiopancreatography; EUS, endoscopic ultrasound; IDUS, intraductal ultrasonography; MDCT, multidetector CT.

several study groups comparing MRCP with ERCP or with EUS.^{1–5} The study group of Rösch stated that after an imaging technique has newly been introduced, scientific evaluations more typically achieve excellent results.¹ With wider use of the method, however, the initial enthusiasm decreases and less impressive accuracy rates are published.^{1–8} This phenomenon could also be substantiated by the data of our study.

We are aware that the analysis performed in this study was not entirely fair since three cross-sectional techniques and one ductal imaging technique supplemented with a limited cross-sectional modality were compared. Furthermore, according to the study protocol, there has been a bias in our study: since some of the patients were referred to our hospital because of acute cholangitis, plastic prosthesis had to be placed immediately during ERCP, which made it difficult for the further applied diagnostic imaging modalities. Under ideal conditions, the diagnostic results might probably have been better for these techniques. However, our results may mirror adequately the clinical reality in most of the patients with biliary strictures of unknown aetiology.

According to our results, the combination of ERCP and IDUS seems to be the most precise imaging modality in diagnosing biliary strictures. For a more complete preoperative evaluation (eg, lymph node invasion), a further imaging modality such as EUS⁹ or MDCT should be added.

Acknowledgements

DD thanks Professor Gary L Powell for helpful and valuable suggestions regarding the manuscript and true friendship for over 20 years.

Dirk Domagk*

Department of Medicine B, University of Muenster, Muenster, Germany

Johannes Wessling*

Department of Clinical Radiology, University of Muenster, Muenster, Germany

Beate Conrad

Department of Medicine B, University of Muenster, Muenster, Germany

Roman Fischbach

Department of Clinical Radiology, University of Muenster, Muenster, Germany

Christina Schleicher

Department of General Surgery, University of Muenster, Muenster, Germany

Werner Böcker

Gerhard-Domagk-Institute of Pathology, University of Muenster, Muenster, Germany

Norbert Senninger

Department of General Surgery, University of Muenster, Muenster, Germany

Achim Heinecke

Department of Medical Statistics, University of Muenster, Muenster, Germany

Walter Heindel

Department of Clinical Radiology, University of Muenster, Muenster, Germany

Wolfram Domschke, Torsten Kucharzik

Department of Medicine B, University of Muenster, Muenster, Germany

Correspondence to: Dr D Domagk, Department of Medicine B, University of Muenster, Albert-Schweitzer-Street 33, D-48129 Muenster, Germany; domagkd@uni-muenster.de

doi: 10.1136/gut.2007.123315

*These authors equally contributed to the study.

Competing interests: None.

References

- 1 Rösch T, Meining A, Frühmorgen S, *et al.* A prospective comparison of the diagnostic accuracy of ERCP, MRCP, CT, and EUS in biliary strictures. *Gastrointest Endosc* 2002;55:870–6.
- 2 Domagk D, Wessling J, Reimer P, *et al.* Endoscopic retrograde cholangiopancreatography, intraductal ultrasonography and magnetic resonance cholangiopancreatography in bile duct strictures: a prospective comparison of imaging diagnostics with histopathological correlation. *Am J Gastroenterol* 2004;99:1684–9.
- 3 Adamek HE, Albert J, Breer H, *et al.* Pancreatic cancer detection with magnetic resonance cholangiopancreatography and endoscopic retrograde cholangiopancreatography: a prospective controlled study. *Lancet* 2000;356:190–3.
- 4 Adamek HE, Albert J, Weitz M, *et al.* A prospective evaluation of magnetic resonance cholangiopancreatography in patients with suspected bile duct obstruction. *Gut* 1998;43:680–3.
- 5 Scheiman JM, Carlos RC, Barnett JL, *et al.* Can endoscopic ultrasound or magnetic resonance cholangiopancreatography replace ERCP in patients with suspected biliary disease? A prospective trial and cost analysis. *Am J Gastroenterol* 2001;96:2900–4.
- 6 Sahai AV, Devonshire D, Yeoh KG, *et al.* The decision-making value of magnetic resonance cholangiopancreatography in patients seen in a referral center for suspected biliary and pancreatic disease. *Am J Gastroenterol* 2001;96:2074–80.
- 7 Varghese JC, Farrell MA, Courtney G, *et al.* A prospective comparison of magnetic resonance cholangiopancreatography with endoscopic retrograde cholangiopancreatography in the evaluation of patients with suspected biliary tract diseases. *Clin Radiol* 1999;54:513–20.
- 8 Bipat S, Phoa SS, van Delden OM, *et al.* Ultrasonography, computed tomography and magnetic resonance imaging for diagnosis and determining resectability of pancreatic adenocarcinoma: a meta-analysis. *J Comput Assist Tomogr* 2005;29:438–45.
- 9 Rösch T, Hofrichter K, Frimberger E, *et al.* ERCP or EUS for tissue diagnosis of biliary strictures? A prospective comparative study. *Gastrointest Endosc* 2004;60:390–6.