Published in final edited form as:

Neurocomputing. 2007 June ; 70(10-12): 2050–2054.

Neuromodulation of short-term synaptic dynamics examined in a mechanistic model based on kinetics of calcium currents

Lian Zhou^a, Shunbing Zhao^a, and Farzan Nadim^{a,b,*}

^a Department of Biological Sciences, Rutgers University, 101 Warren St., Newark, NJ 07102, USA

^b Department of Mathematical Sciences, NJIT, 323 MLK Blvd., Newark, NJ 07102, USA

Abstract

Network plasticity arises in large part due to the effects of exogenous neuromodulators. We investigate the neuromodulatory effects on short-term synaptic dynamics. The synapse from the lateral pyloric (LP) to the pyloric dilator (PD) neuron in the pyloric network of the crab *C. borealis* has both spike-mediated and non-spike-mediated (graded) components. Previous studies have shown that the graded component of this synapse exhibits short-term depression. Recent results from our lab indicate that in the presence of neuromodulatory peptide proctolin, low-amplitude presynaptic stimuli switch the short-term dynamics of this graded component from depression to facilitation. In this study, we show that this facilitation is correlated with the activation of a presynaptic inward current that is blocked by Mn^{2+} suggesting that it is a slowly-accumulating Ca^{2+} current. We modify a mechanistic model of synaptic release by assuming that the low-voltage-activating Ca^{2+} current in our system is composed of two currents with fast (I_{CaF}) and slow (I_{CaS}) kinetics. We show that if proctolin adjusts the activation rate of I_{CaS} , this leads to accumulation of local intracellular Ca^{2+} in response to multiple presynaptic voltage stimuli which, in turn, results in synaptic facilitation. Additionally, we assume that proctolin increases the maximal conductances of Ca^{2+} currents in the model, consistent with the increased synaptic release found in the experiments. We find that these two presynaptic actions of proctolin in the model are sufficient to describe its actions on the short-term dynamics of the LP to PD synapse.

Keywords

proctolin; graded release; short-term plasticity; central pattern generator; stomatogastric

Introduction

Neuromodulation can reconfigure a network to produce a multitude of outputs [1]. The crustacean stomatogastric nervous system (STNS) is one of the most extensively researched neural systems in studying the effects of neuromodulation. There are several reports of the actions of neuromodulators on intrinsic neuronal properties and synaptic strength in the STNS (see [2] for a review) yet there are few reports of neuromodulatory effects on the short-term dynamics of synaptic transmission. A recent report from our laboratory showed that neuromodulation can alter short-term synaptic dynamics to the extent that a depressing synapse can become facilitating [3]. Such a drastic shift in dynamics can have significant consequences

*Corresponding Author: Farzan Nadim, farzan@njit.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

for the role of that synapse in network activity. In this study, we use a combination of experiments and modeling to explore the synaptic mechanisms underlying the neuromodulator effects on the short-term dynamics of pyloric synapses.

We investigate the effects of the neuropeptide proctolin on the dynamics of the inhibitory synapse from the lateral pyloric (LP) to the pyloric dilator (PD) neuron in the rhythmic pyloric network of the crab. This synapse has both spike-mediated and non-spike-mediated (graded) components and is the only chemical feedback to the pyloric pacemaker neurons. The graded component of this synapse demonstrates short-term depression in control saline, yet, in the presence of proctolin, low-amplitude (<30 mV) presynaptic stimulation causes the graded component of the LP to PD synapse to facilitate [3]. In contrast, with high-amplitude (>30 mV) stimuli the synapse remains depressing in the presence of proctolin, albeit enhanced in strength. We show that the switch to facilitation is correlated with the activation of a slowly activating presynaptic inward current. This inward current is blocked by Mn^{2+} , suggesting that it is a slowly-accumulating Ca^{2+} current activated by proctolin.

We propose that the switch from depression to facilitation in the LP to PD synapse is due to modifications in presynaptic Ca^{2+} current kinetics. Ca^{2+} channels can be divided into low-voltage-activated (LVA) and high-voltage-activated (HVA) channels. LVA channels such as T-type Ca^{2+} currents begin to activate ~ -60 mV, peak at ~ -30 mV and their amplitude decreases with further depolarization [4]. Since LVA channels have a low activation threshold, they support action potential generation, regulate subthreshold oscillatory activities, produce graded transmission and modulate spike-mediated transmission [5]. HVA Ca^{2+} currents activate ~ -40 to -10 mV, peak at ~ 0 mV [6], and are believed to underlie spike-mediated transmitter release [6].

LVA Ca^{2+} currents are usually inactivating. Many LVA Ca^{2+} current consists of two kinetically distinct components; one that activates/inactivates rapidly (I_{CaF}), another that activates/inactivates slowly (I_{CaS}) [5,7]. We propose that the LP neuron also has two LVA Ca^{2+} currents. Additionally, we predict that proctolin not only adjusts the amplitude of these two currents but also causes I_{CaS} to activate even more slowly, which leads to accumulation of total Ca^{2+} current and facilitation of synaptic release from LP. Using a model of synaptic release, we demonstrate that these proposed effects of proctolin explain the switch in the LP to PD synapse from depression to facilitation, in response to low-amplitude presynaptic stimuli.

Model

The synaptic release model is modified from our previous model which simulates the effect of neurotransmitter release around the readily releasable pool (RRP) [8]. Our previous model involved only a single non-inactivating LVA Ca^{2+} current. In the current model, we include two LVA Ca^{2+} currents with realistic kinetics. Specifically, the LVA Ca^{2+} currents comprise a fast (I_{CaF}) and a slow (I_{CaS}) activating/inactivating component which are kinetically distinct. Additionally, in the current model, the HVA Ca^{2+} current (I_{CaH}) also plays a role in the graded release, especially with high-amplitude presynaptic voltage pulses. Briefly, the model is described as follows:

a) The Ca^{2+} currents: The total Ca^{2+} current underlying graded release is $I_{CaG} = I_{CaF} + I_{CaS} + I_{CaH}$, where I_{CaF} and I_{CaS} are the fast and slow LVA calcium current respectively, I_{CaH} is the HVA calcium current. The I_{CaF} and I_{CaS} are modeled as $I_{Ca} = G_{max} m h (V_{pre} - E_{Ca})$ and I_{CaH} is modeled as $I_{Ca} = G_{max} m (V_{pre} - E_{Ca})$, where G_{max} is the maximum conductance (G_{max} in μS : **Control**: I_{CaS} 0.002, I_{CaF} 0.01, I_{CaH} 0.014; **Proctolin**: I_{CaS} 0.008, I_{CaF} 0.0175, I_{CaH} 0.018), V_{pre} is the presynaptic membrane potential and E_{Ca} is the reversal potential of Ca^{2+} (fixed at 100 mV), m is the activation gate and h is the inactivation gate, the calculation of m and h is described below.

A set of differential equations are used to describe activation and inactivation gate of the above calcium currents: $dx/dt = (x_{\infty}(V_{pre}) - x)/\tau(V_{pre})$, $x = m, h$, where $x_{\infty}(V) = 1/(1 + \exp((V - V_x)/k))$ denotes the steady-state activation/inactivation process with parameter values V_x (in mV) I_{CaS} : $m -35, h -27$; I_{CaF} : $m -30, h -45$; I_{CaH} : $m -22.5$ and k (in mV) I_{CaS} : $m -2.0, h 10.0$; I_{CaF} : $m -3.0, h 0.2$; I_{CaH} : $m -6.0$. The time constants are modeled as $\tau(V) = \tau_l + (\tau_h - \tau_l)/(1 + \exp(-(V + 35)/10))$ with parameter values **Control**: τ_l (in msec) I_{CaS} : $m 50, h 200$; I_{CaF} : $m 1, h 200$; I_{CaH} : $m 1$ and τ_h (in msec) I_{CaS} : $m 50, h 5$; I_{CaF} : $m 100, h 5$; I_{CaH} : $m 1$. **Proctolin**: τ_l (in msec) I_{CaS} : $m 1000, h 5000$ and τ_h (in msec) I_{CaS} : $m 1000, h 5$; others unchanged.

b) The local intracellular calcium $[Ca^{2+}]_i$: Given the total Ca^{2+} current I_{Cag} , $[Ca^{2+}]_i$ is described by $d[Ca^{2+}]_i/dt = (-\lambda I_{Cag} - [Ca^{2+}]_i)/\tau$, where $\lambda = 11.0 \mu M/nA$ and $\tau = 1.0$ msec.

c) Synaptic release: The rate of change of the number of vesicles (N) in the RRP depends on the interaction of a supply rate of new vesicles $P(N)$ and a release rate $R(N)$: $dN/dt = P(N) - R(N)$. The supply rate is dependent on internal concentration of Ca^{2+} :

$$P(N) = \alpha \frac{[Ca^{2+}]_i + a_1}{[Ca^{2+}]_i + a_2} (N_{max} - N)$$

where $\alpha = 0.05 \text{ msec}^{-1}$, $a_1 = 2 \mu M$, $a_2 = 100 \mu M$ and N_{max} is the maximum number of vesicles in the RRP (80). $R(N)$ is given by $R(N) = \gamma N [Ca^{2+}]_i^4$ where $\gamma = 5 \times 10^{-7} \text{ msec}^{-1} \mu M^{-4}$.

d) The postsynaptic response: The postsynaptic potential is given by

$C \frac{dV}{dt} = -G_{syn}(t)(V - V_{syn}) - g_m(V - V_{rest})$ where C is the membrane capacitance (1nF), V_{syn} is the synaptic reversal potential (set at -80 mV), g_m is the intrinsic membrane conductance (set at $0.0258 \mu S$), and V_{rest} is the membrane resting potential (set at -55 mV). The postsynaptic conductance change is calculated as $G_{syn}(t) = \int_0^t mG_{syn}(t')R(N)dt'$ where mG_{syn} is the miniature synaptic conductance (due to a single vesicle) which is obtained from fits to data [8].

Biological Methods

Experiments were carried on adult male crabs (*C. borealis*) purchased from local distributors (Newark, NJ). Details of experimental measurements are identical to those described in [9]. Spontaneous activity in the pyloric network was blocked by superfusion with saline containing $0.1 \mu M$ TTX (Biotium, Hayward, CA). (TTX does not block graded synaptic release in this system.) For measurements of synaptic resonance the presynaptic LP neuron was voltage clamped with two electrodes (TEVC) at a holding potential of -60 mV and multiple low- (20 mV) or high-amplitude (60 mV) voltage pulses were applied. The postsynaptic response was measured in the PD neuron in current clamp mode, held at a resting potential of ~ -60 mV. All data was recorded after 15 minutes of drug application or 45 minutes of wash. Data were acquired at 5 kHz and analyzed using the PClamp 9.2 software (Molecular Devices, Union City, CA).

Results

To characterize short-term plasticity of the LP to PD synapse, we injected a series of multiple low-amplitude (20 mV, V_{LP} in Fig. 1) and high-amplitude (not shown) voltage pulses into the LP neuron and recorded the postsynaptic potential in the PD neuron. The experiment was performed with two different sets of treatment. First, recordings were done in both control saline (no modulators) and in the presence of 10^{-6} M proctolin (Fig. 1 A, B). This proctolin concentration was used because it has been shown to best replicate the effects of proctolin release from endogenous modulatory neurons on the pyloric network [10]. Second, the

experiments were performed in normal Ca^{2+} or after blocking all Ca^{2+} currents (and therefore synaptic transmission), in both control and proctolin, by substituting the Ca^{2+} (13.0 mM Ca^{2+} , the normal concentration in crab saline) with Mn^{2+} (12.9 mM Mn^{2+} and 0.1 mM Ca^{2+}). In all conditions we made simultaneous measurements of the presynaptic current (Fig. 1, I_{LP}) and the postsynaptic potential (Fig. 1, V_{PD}). The difference between the presynaptic currents measured in Ca^{2+} saline and in Mn^{2+} saline is a putative calcium current (Fig. 1, $I_{LP}(\text{Ca}^{2+}-\text{Mn}^{2+})$). In the control saline (with normal Ca^{2+}), either the synaptic response showed depression (not shown, but see [3]) or no apparent synaptic plasticity was observed (Fig. 1A, V_{PD}). In contrast, the synaptic response showed facilitation in proctolin saline (with normal Ca^{2+}) (Fig. 1B, V_{PD}). V_{PD} is not shown in Mn^{2+} saline because there was no synaptic transmission in Mn^{2+} saline either in control or in proctolin. The putative calcium current was small in control conditions and its amplitude showed no obvious variation among the different voltage pulses (Fig. 1A, $I_{LP}(\text{Ca}^{2+}-\text{Mn}^{2+})$). However, in the presence of proctolin, this current increased in amplitude with each subsequent pulse, indicating accumulation of Ca^{2+} currents (Fig. 1B, arrows in $I_{LP}(\text{Ca}^{2+}-\text{Mn}^{2+})$ trace). This accumulation of inward currents was correlated with the facilitation of the IPSPs in the postsynaptic PD neuron (Fig. 1B, V_{PD} trace).

In contrast with the responses measured using low-amplitude (20 mV) pulses, with high-amplitude (60 mV) pulses there was little or no accumulation of the putative Ca^{2+} current with multiple pulses in either control or proctolin. Additionally, in both conditions the IPSPs were always depressing (not shown, but see [3]).

The main goal of the model is to explain how proctolin causes the short-term synaptic dynamics of the LP to PD IPSPs to switch from depression (or no change) to facilitation in response to low-amplitude presynaptic stimuli, whereas, in response to high-amplitude stimuli, the synapse remains depressing both in control and in proctolin. Additionally, the model demonstrates the enhancement of the IPSP amplitudes in proctolin, in response to any stimulus.

We incorporate the effect of proctolin in the model by modifying the activation and inactivation of the slow LVA Ca^{2+} current as well as the maximum conductances of all three Ca^{2+} currents. We propose that proctolin increases the maximum conductance of the HVA Ca^{2+} current because our experimental measurements have shown that proctolin also enhances spike-mediated release [8] which depends on this current. To simulate realistic Ca^{2+} channel kinetics, the LVA currents were set to peak at -30 mV while HVA current peaked around 0 mV. There was a decrease in both LVA Ca^{2+} current amplitudes with high-amplitude voltage pulses, which resulted from fast inactivation at high presynaptic voltages (Fig. 2A, B, control LVA and HVA).

To explain the effect of proctolin on short-term synaptic dynamics we assumed that proctolin modified the kinetics of the model I_{CaS} by increasing the time constant of I_{CaS} activation from 50 to 1000 ms and the inactivation time constant (at low voltages) from $\tau_1=200$ to $\tau_1=5000$ ms in the presence of proctolin. Consequently, in the presence of proctolin, I_{CaS} accumulated with multiple low-amplitude presynaptic voltage pulses, as observed in the biological recordings (Fig. 1B). Furthermore, the maximum conductances of all Ca^{2+} currents were increased with proctolin compared to control (Fig. 2A, B: proctolin LVA and HVA). Due to the new kinetics of I_{CaS} , the total Ca^{2+} current I_{CaT} also accumulated with multiple low-amplitude presynaptic voltage pulses in the presence of proctolin and its amplitude was increased (Fig. 2, I_{CaT}).

In our model of synaptic release, the total effect of the short term synaptic plasticity is determined by the interaction between the intracellular presynaptic Ca^{2+} concentration $[\text{Ca}^{2+}]_i$ and the number of vesicles N in the RRP. In control conditions, N is largely depleted with multiple high-amplitude voltage pulses, which results in the depression shown in the

postsynaptic IPSPs in the PD neuron (Fig 2, control V_{PD}). There is also depression in control conditions with low-amplitude voltage pulses but to a much smaller extent. Proctolin leads to the accumulation of $[Ca^{2+}]_i$ with multiple low-amplitude voltage pulses, which in turn gives rise to the facilitation of the IPSPs in the PD neuron (Fig 2, proctolin V_{PD}). In contrast, with multiple high-amplitude voltage pulses, there is synaptic depression which is again attributed to the depletion of the number of vesicles N in the RRP and the decrease of LVA Ca^{2+} current amplitudes.

Discussion

Short-term synaptic dynamics such as depression and facilitation play an important role in network operation, yet their contribution to network activity is unclear. In a previous study, we produced a model of synaptic release for the LP to PD synapse in the rhythmically active crab pyloric network based on changes in intrinsic low- (LVA) and high-voltage activated (HVA) Ca^{2+} currents [8]. The current study modifies this model by incorporating realistic Ca^{2+} currents into the model to examine the role of extrinsic neuromodulation on short-term synaptic plasticity. We demonstrate that the short-term dynamics of the LP to PD synapse change from depression or stationary dynamics to facilitation in the presence of the neuropeptide proctolin. Additionally, we show that the appearance of synaptic facilitation is correlated with a presynaptic inward current that accumulates with multiple voltage pulses and is blocked if bath Ca^{2+} is substituted with Mn^{2+} . This current appears with voltage pulses from -60 to -40 mV, suggesting that it is a LVA Ca^{2+} current. Additionally, proctolin enhances synaptic release in response to presynaptic stimuli of any amplitude [3].

We incorporate one HVA and two LVA (I_{CaF} and I_{CaS}) currents in our model to properly model spike-mediated (high-threshold) and graded (low-threshold) synaptic release. We then demonstrate that the actions of proctolin on the synapse can be explained by two simple mechanisms. First, we assume that proctolin enhances the maximum conductances of all presynaptic calcium currents, thus increasing release probability. Second, we demonstrate that the appearance of synaptic facilitation in the presence of proctolin can be explained if the kinetics of I_{CaS} is slowed down by proctolin. Although it is possible that the kinetics of the other Ca^{2+} currents are also changed by proctolin, the change in synaptic dynamics as well as the cumulative inward current (Fig. 1B) were sufficiently explained by the change in the I_{CaS} kinetics. With high-amplitude presynaptic stimuli, the synapse shows depression in both control saline and in the presence of proctolin [3]. Short-term synaptic depression in response to high-amplitude presynaptic stimuli in the model is due to depletion of vesicles, both in control and in proctolin, as described previously [8]. We would like to point out that there are other potential mechanisms that may explain how proctolin enhances synaptic release. The model presented here is not meant to rule out other mechanisms but to provide a predictive hypothesis for the synaptic response in more realistic conditions where both spike-mediated and graded release are present and modulated by proctolin. This predictive model would provide grounds for further experimental exploration.

The importance of short-term synaptic dynamics in network activity has been demonstrated in computational and experimental studies [11,12]. In addition, many studies show that synaptic dynamics are also subject to neuromodulation [13]. In the current study we used a mechanistic model of synaptic release to explore the underlying mechanisms of the effects of neuromodulation on short-term synaptic dynamics. To understand how neuromodulators reshape network output, however, it is necessary to explore how actions of the neuromodulator on synaptic and cellular components are integrated at the network level. The pyloric network could prove to be an ideal model for such a study because the actions of the neuropeptide proctolin on pyloric neurons have been incorporated into realistic biophysical models [14] and the current study provides the first detailed model of proctolin actions on pyloric synapses.

Acknowledgements

Supported by NIH grant MH-60605 (FN).

References

1. Marder E, Thirumalai V. Cellular, synaptic and network effects of neuromodulation. *Neural Netw* 2002;15:479–93. [PubMed: 12371506]
2. Nusbaum MP, Beenhakker MP. A small-systems approach to motor pattern generation. *Nature* 2002;417:343–50. [PubMed: 12015615]
3. Atamturktur S, Manor Y, Nadim F. Proctolin enables a shift in synaptic dynamics from depression to facilitation due to the presynaptic waveform amplitude in a rhythmic network. *Soc Neurosci Abst* 2004;30
4. Perez-Reyes E. Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol Rev* 2003;83:117–61. [PubMed: 12506128]
5. Ivanov AI, Calabrese RL. Modulation of spike-mediated synaptic transmission by presynaptic background Ca^{2+} in leech heart interneurons. *J Neurosci* 2003;23:1206–18. [PubMed: 12598609]
6. Lacinova L. Voltage-dependent calcium channels. *Gen Physiol Biophys* 2005;24(Suppl 1):1–78. [PubMed: 16096350]
7. Joksovic PM, Bayliss DA, Todorovic SM. Different kinetic properties of two T-type Ca^{2+} currents of rat reticular thalamic neurones and their modulation by enflurane. *J Physiol* 2005;566:125–42. [PubMed: 15845580]
8. Zhou L, LoMauro R, Nadim F. The interaction between facilitation and depression of two release mechanisms in a single synapse. *Neurocomputing* 2006;69:1001–5.
9. Manor Y, Nadim F. Synaptic depression mediates bistability in neuronal networks with recurrent inhibitory connectivity. *J Neurosci* 2001;21:9460–70. [PubMed: 11717380]
10. Blitz DM, Christie AE, Coleman MJ, Norris BJ, Marder E, Nusbaum MP. Different proctolin neurons elicit distinct motor patterns from a multifunctional neuronal network. *J Neurosci* 1999;19:5449–63. [PubMed: 10377354]
11. Cook DL, Schwindt PC, Grande LA, Spain WJ. Synaptic depression in the localization of sound. *Nature* 2003;421:66–70. [PubMed: 12511955]
12. Tsodyks M, Uziel A, Markram H. Synchrony generation in recurrent networks with frequency-dependent synapses. *J Neurosci* 2000;20:1–5. [PubMed: 10627575]
13. Kreitzer AC, Regehr WG. Modulation of transmission during trains at a cerebellar synapse. *J Neurosci* 2000;20:1348–57. [PubMed: 10662825]
14. Soto-Treviño C, Rabbah P, Marder E, Nadim F. Computational model of electrically coupled, intrinsically distinct pacemaker neurons. *J Neurophysiol* 2005;94:590–604. [PubMed: 15728775]

Biographies

Lian Zhou is currently a graduate student in the Department of Biological Sciences at Rutgers University-Newark.

Shunbing Zhao is currently a graduate student in the

Department of
Biological
Sciences at
Rutgers
University-
Newark.

**Farzan
Nadim** is
currently a
professor in
the
Department of
Mathematical
Sciences at
New Jersey
Institute of
Technology
and the
Department of
Biological
Sciences at
Rutgers
University-
Newark. He
received his
Ph.D. in
Mathematics
from Boston
University and
was a post-
doctoral fellow
in
neurobiology
at Emory
University and
Brandeis
University.

Fig 1.

Facilitation of the LP to PD inhibitory postsynaptic potentials (V_{PD} trace) in proctolin is associated with the activation of a slow Ca^{2+} -like inward current. The presynaptic currents in the LP neuron are measured in Ca^{2+} saline (black) and in Mn^{2+} saline (gray) with a train of low-amplitude presynaptic voltage pulses. The difference between the two measured currents (bottom trace) shows a new inward current activated in proctolin. The amplitude of this current increases with each pulse (arrows).

Fig. 2.

Model of Ca^{2+} currents underlying graded release. **(A)** Low-amplitude voltage pulses in the presynaptic LP neuron activate both I_{CaS} and I_{CaF} LVA Ca^{2+} currents. This leads to an increase in intracellular Ca^{2+} levels ($[\text{Ca}^{2+}]_i$), which causes the synaptic release ($R(N)$) and variations in the number of available vesicles N in the RRP. The postsynaptic response (V_{PD}) is obtained based on $R(N)$. Due to fast inactivation, I_{CaS} and I_{CaF} become smaller with high-amplitude voltage pulses. In contrast, the HVA Ca^{2+} current I_{CaH} becomes larger with high-amplitude voltage pulses. In response to both low- and high-amplitude presynaptic pulses, there is depletion of the available vesicles N in the RRP. Consequently, the IPSPs in the PD neuron show short-term depression (V_{PD}). The extent of depression is larger in response to high-amplitude stimuli. $I_{\text{CaG}} = I_{\text{CaS}} + I_{\text{CaF}} + I_{\text{CaH}}$. **(B)** Proctolin changes the activation time constant of I_{CaS} , which leads to the accumulation of I_{CaS} under low voltage pulses. Proctolin also increases the maximum conductance of I_{CaS} , I_{CaF} and I_{CaH} . These two changes result in the accumulation of $[\text{Ca}^{2+}]_i$ with low-amplitude presynaptic voltage pulses, which gives rise to the facilitation of the IPSPs in the PD neuron (V_{PD}). In contrast, with high-amplitude voltage pulses, there is extensive depletion of N in the RRP and a decrease of LVA Ca^{2+} current. These result in depression of the IPSPs (V_{PD}). Note, however, that in both cases the synaptic response in proctolin is larger than control conditions.