We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer. Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.

FAMILIAL HYPERCHOLESTEROLAEMIA

Cascade testing is tried and tested and cost effective

Wald et al suggest that screening children for high cholesterol at the time of immunisation, and then testing the parents of affected children, is an effective screening method for familial hypercholesterolaemia.¹

Earlier data from our group showed that universal screening was much less cost effective than cascade testing (family tracing from affected patients). A recent pilot study of cascade testing at five NHS trusts, funded by the Department of Health (www.fhcascade.org.uk), has confirmed cascading to be feasible, cost effective, and acceptable to both patients and clinicians.² Our recommendations to the department include that, because of the degree of overlap in cholesterol concentrations in affected and unaffected individuals, DNA testing is required to underpin a diagnosis of familial hypercholesterolaemia. The use of cholesterol testing as proposed by Wald et al will lead either to many false positive diagnoses in children, causing unnecessary anxiety, testing, and costs, or to true positives being missed because the cut-off point is set too high.

Around 15 000 patients with familial hypercholesterolaemia (of the estimated 100 000 cases) attend lipid clinics in the United Kingdom and cascading from these known cases should be the first approach as half of their first degree relatives will be affected.³ This has been evaluated and should take priority for resources.⁴

However, finding more new patients will become important after the cascade work, which is estimated to find half of UK cases. The proposal to screen infants has clinical merit but would need to be piloted carefully. Approaches that circumvent using children

as the primary contact warrant further investigation, for example, using data from general practice records, which recent data suggest to be feasible,⁵ or testing those who have had a heart attack at an early age.

S Gaye Hadfield project coordinator, FH cascade testing audit, London IDEAS Genetics Knowledge Park

Steve E Humphries BHF professor of cardiovascular genetics, Department of Cardiovascular Genetics, British Heart Foundation Laboratories, Royal Free and University College Medical School, London WC1E 6JJ rmhaseh@ucl.ac.uk

Competing interests: None declared.

- 1 Wald DS, Bestwick JP, Wald NJ. Child-parent screening for familial hypercholesterolaemia: screening strategy based on a meta-analysis. *BMJ* 2007;335:599-603. (22 September.)
- 2 Marks D, Wonderling D, Thorogood M, Lambert H, Humphries SE, Neil AW. Cost-effectiveness analysis of different approaches of screening for familial hypercholesterolaemia. *BMJ* 2002;324:1303-6.
- 3 Marks D, Thorogood M, Farrer M, Humphries SE. Census of clinics providing specialist lipid services in the United Kingdom. *J Public Health Med* 2004;26:353-4.
- 4 Bhatnagar D, Morgan J, Siddiq S, Mackness MI, Miller JP, Durrington PN. Outcome of case finding among relatives of patients with known heterozygous familial hypercholesterolaemia. *BMJ* 2000;321:1497-500.
- 5 Gray J, Jaiyeola A, Whiting M, Model M, Wierzbicki A. Identifying patients with familial hypercholesterolaemia in primary care: an informatics-based approach in one primary care centre. *Heart* 2007 Jun 17; Epub ahead of print

Child-parent screening may have adverse psychological effects

Wald et al do not mention the potential adverse psychological impact of child-parent screening for familial hypercholesterolaemia.¹

No matter how well these issues are communicated, this test will exacerbate the cholesterol neurosis that much of the population already seems to have. General practitioners in particular are aware that a screening programme which sounds fine in theory can translate into anxiety in patients. Throw children into the mix and the anxieties will be even higher.

Telling parents that their child has high cholesterol will worry them, which makes putting things into perspective difficult. It risks casting a shadow over childhood because parents view their progeny as threatened or fragile. This will be compounded by the fact that no immediate therapeutic action can be taken. Parents will feel that their child is vulnerable and that nothing can be done in the immediate term.

LETTERS

At best this will make them uncomfortable, at worst it could blight their parenthood.

Wald et al suggest delaying treatment until adulthood. At which point—after a childhood of being less “well” than their peers and overprotected by their parents—they may or may not decide to be treated and may or may not be happy that this test was imposed on them. Perhaps their thwarted efforts to obtain life insurance will help them make up their minds.

Keith A Hopcroft general practitioner
Laindon Health Centre, Basildon, Essex SS15 5TR
keithhopcroft@supanet.com

Competing interests: None declared.

- 1 Wald DS, Bestwick JP, Wald NJ. Child-parent screening for familial hypercholesterolaemia: screening strategy based on a meta-analysis. *BMJ* 2007;335:599-603. (22 September.)

See bmj.com for authors' replies

DIRECT TO CONSUMER ADVERTISING

The world of the market place

Magrini and Font discuss direct to consumer advertising,¹ but drug promotion at an earlier stage may have already influenced the views of the public and patients.

Current European legislation on drug promotion makes it difficult to define what promotion by a pharmaceutical company before market authorisation entails. Many drug companies employ marketing companies to promote new drugs. The recent premature promotion of Herceptin for early breast cancer is viewed as exemplary by the industry and to be copied.

Releasing early results from high profile trials at major cancer conferences can gain worldwide publicity. When supported by website press releases to inform the stock market, one sided information is freely available in the market place.

The small patient group “Women Fighting for Herceptin” was supported by a leading marketing company.² The ensuing national media campaign pushed politicians into supporting the unlicensed use of Herceptin and hence undermined the UK Medicines Act and the role of the National Institute for Health and Clinical Excellence (NICE).³ Many healthcare commissioners caved in under the subsequent pressure and agreed early funding.

The drive for pharmaceutical company profits is changing the tactics of information use. This is more likely to undermine public trust in the industry than help patients.

Jonathan V Howell consultant in public health
West Midlands Specialised Commissioning Team, Edwin House, Burton on Trent DE14 2WF
jonathan.howell@lycos.com

Competing interests: JWH advised healthcare commissioners about the funding of unlicensed Herceptin. His complaint about the promotion of Herceptin was upheld.

- 1 Magrini N, Font M. Direct to consumer advertising of drugs in Europe. *BMJ* 2007;335:526. (15 September.)
- 2 Boseley S. The selling of a wonder drug. *Guardian* 2006;29 March. www.guardian.co.uk/science/2006/mar/29/medicineandhealth.health
- 3 Department of Health. Speech by Rt Hon Patricia Hewitt MP, Secretary of State for Health, 25 October 2005: Breast cancer awareness. Especially paragraphs 18 and 21. www.dh.gov.uk/en/News/Speeches/Speecheslist/DH_4121929

MEDICAL MIGRATION

Full house

Winyard argues that the BMA is being “ambiguous” in criticising both the medical training application service (MTAS) fiasco and the changes in immigration policy.¹ He implies that it is incongruent to do so, as stopping immigration is necessary for the system to run adequately. Although this is somewhat true, the BMA is not lobbying for the rights of future immigrants, but for the rights of current immigrants who have been unfairly forced into exile by the new immigration rules. The BMA deserves nothing but praise for this.

He also mentions that the United Kingdom currently has “the worst of all worlds,” as it invests in expanding medical schools but still recruits people from overseas, only to fail to give them all a job down the line. But this is actually quite clever. In the end, what you get is a surplus of intelligent, well intentioned, overqualified people with no transferable skills, thousands of pounds in debt, desperately looking for a job in the one company that has the monopoly for employing their services. It's the best of all worlds, and any company in the private sector would kill for this.

The way forward is clear. The house is full, so please shut all the doors. All of them. Not only permit-free training, but highly skilled migrant programmes, professional and linguistic assessment board tests, and all entry to UK medical schools by overseas students aspiring to stay in the UK. The fact that all these structures still offer false hope (and charge heavily for the privilege) is nothing but cruel. But what about those already inside? Would it be too much to let them stay? Even if it means a little more

competition? After all, they did help to build the house.

Jorge Zimbron academic foundation year 2
Institute of Psychiatry at the Maudsley, King's College, London SE5 8AF jorgezimbron@doctors.net.uk
Competing interests: JZ is an overseas UK graduate unlikely to be able to apply for specialist training next year.

- 1 Winyard G. Medical immigration: the elephant in the room. *BMJ* 2007;335:593-5. (22 September.)

The real elephant in the room

Winyard has missed the issue of non-European Union UK medical graduates.¹ These students make up about 15% of medical undergraduates, and UK medical schools depend on their financial backing to survive. Each student would have forked out at least £80 000 (€114 000; \$162 000) by year 5, excluding the cost of living. When these students first chose to study in Britain, the rules would have allowed them fair access to postgraduate training.

Since April 2006, with the abolition of permit-free training, these students are left in limbo, with a strong feeling of breach of moral contract. I fail to understand why these non-EU students are included in the figures used by the Department of Health and the Foreign Office when accounting for UK graduates but are judged as “international medical graduates” when applying for jobs.

Also it costs the taxpayer about £250 000 to train each one.

Mohammad Farhad Peerali fourth year medical student
University of Sheffield, Sheffield S10 2TN
mda04mfp@shef.ac.uk

Competing interests: MFP is an international medical student in a British medical school.

- 1 Winyard G. Medical immigration: the elephant in the room. *BMJ* 2007;335:593-5. (22 September.)

“BARE BELOW THE ELBOWS”

A cheap soundbite

We were astounded to read that the Department of Health plans to ban doctors from wearing white coats because of the risk of cross infection.^{1 2} The package describes a “bare below the elbows” dress code (short sleeves, no wrist watch, no jewellery, no ties when carrying out clinical activity).

What is the evidence for this? It is impossible to say because this policy is based on two as yet unpublished reviews. Surely, they should have been made available first for discussion. We doubt that evidence is convincing as the working group's first conclusion is that there is no conclusive evidence that work clothes

are a significant risk in terms of spreading infection, and the policy document ends with, “it seems unlikely that uniforms are a significant source of cross-infection.”

The document considers it good practice for medical staff to wear short sleeved garments and not white coats because “cuffs become heavily contaminated and are more likely to come into contact with patients.” Presumably this conclusion is based on bacteria such as *Staphylococcus aureus* being found on a quarter of white coats.³ But these coats were worn for up to eight weeks and not changed daily. Why? Because hospitals no longer provide clean white coats.

But maybe we should not be astounded. Dirty hospitals are expensive to sort out, whereas “bare below the elbows” is a cheap soundbite that saves money. But the policy ignores the advantages of white coats and name badges—ease of identification and cleaning, protection, and usefulness for carrying things. Most patients in the UK prefer doctors to wear white coats,⁴ and doctors wear white coats in most countries.⁵

White coats should not be banned. Yes, use plastic aprons when examining patients, but insist that you are provided with a clean white coat every day.

Competing interests: None declared.

Adam Magos, consultant gynaecologist
a.magos@medsch.ucl.ac.uk

Allan Maclean professor of obstetrics and gynaecology

Daryll Baker consultant vascular surgeon

Nicholas Goddard consultant orthopaedic surgeon

Olagunju Ogunbiyi consultant colorectal surgeon

Royal Free Hospital, London NW3 2QG

- 1 White coats to go in cleanliness bid. *BMJ* 2007;335:582. (22 September.)
- 2 Department of Health. *An evidence base for developing local policy*. London: DOH, 2007.
- 3 Wong D, Nue K, Hollis P. Microbial flora on doctors' white coats. *BMJ* 1991;303:1602-4.
- 4 Tiwari A, Abesinghe N, Hall A, Perera P, Ackroyd JS. Should doctors wear white coats? The patient's perspective. *J Eval Clin Pract* 2001;7:343-5.
- 5 Major K, Hayase Y, Balderrama D, Lefor AT. Attitudes regarding surgeons' attire. *Am J Surg* 2005;190:103-9.

‘It’s soap on a rope’