Published in final edited form as:

Cell Calcium. 2007 ; 42(4-5): 409–417.

Neuronal calcium channels: Splicing for optimal performance

Annette C Gray¹, Jesica Raingo, and Diane Lipscombe²

Department of Neuroscience, Brown University, Providence, RI 02912

Abstract

Calcium ion channels coordinate an astounding number of cellular functions. Surprisingly, only 10 $\text{Ca}_v\alpha_1$ subunit genes encode the structural cores of all voltage-gated calcium channels. What mechanisms exist to modify the structure of calcium channels and optimize their coupling to the rich spectrum of cellular functions? Growing evidence points to the contribution of post-translational alternative processing of calcium channel RNA as the main mechanism for expanding the functional potential of this important gene family. Alternative splicing of RNA is essential during neuronal development where fine adjustments in protein signaling promote and inhibit cell-cell interactions and underlie axonal guidance. However, attributing a specific functional role to an individual splice isoform or splice site has been difficult. In this regard, studies of ion channels are advantageous because their function can be monitored with precision, allowing even subtle changes in channel activity to be detected. Such studies are especially insightful when coupled with information about isoform expression patterns and cellular localization.

In this paper we focus on two sites of alternative splicing in the N-type calcium channel $\text{Ca}_v2.2$ gene. We first describe cassette exon 18a that encodes a 21 amino acid segment in the II-III intracellular loop region of $\text{Ca}_v2.2$. Here we show that e18a is up-regulated in the nervous system during development. We discuss these new data in light of our previous reports showing that e18a protects the N-type channel from cumulative inactivation. Second, we discuss our published data on exons e37a and e37b, which encode 32 amino acids in the intracellular C-terminus of $\text{Ca}_v2.2$. These exons are expressed in a mutually exclusive manner. Exon e37a-containing $\text{Ca}_v2.2$ mRNAs and their resultant channels express at higher density in dorsal root ganglia and, as we showed recently, e37a increases N-type channel sensitivity to G protein-mediated inhibition, as compared to generic e37b-containing N-type channels.

Alternative splicing in voltage-gated calcium channels

Human, rat, and mouse genomes contain ten genes that encode $\text{Ca}_v\alpha_1$ subunits. The nervous system expresses nine of these genes. $\text{Ca}_v\alpha_1$ genes are large, spanning 100 - 800 kb of human genome sequence and containing upwards of 50 exons [1-4]. The large size of these genes provides ample opportunity for alternative splicing. Based on already known splice sites, the theoretical number of splice isoforms possible from a single $\text{Ca}_v\alpha_1$ subunit exceeds 1,000 [3, 4]. Other genes, far smaller than $\text{Ca}_v\alpha_1$ genes, have the potential to generate a similar or greater number of splice isoforms [5], and in some genes the number of possible isoforms is in the tens of thousands [6].

²Correspondence: Diane Lipscombe, Department of Neuroscience, Sidney E Frank Hall for Life Sciences, 185 Meeting Street, Providence RI 02912, USA, Diane_Lipscombe@brown.edu, Tel: 401 863 1092, Fax: 401 863 1074

¹Current address: Department of Biology, Brandeis University, Waltham, MA 02454

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Alternative splicing occurs at sites important for controlling calcium channel activity, creating an array of functionally distinct calcium channels. Thus each neuron, in theory, can optimize calcium channel activity for the specific task at hand by orchestrating appropriate exon selection. Placing a specific channel isoform within a specific cell type can in turn provide information about the requirements of particular calcium signaling cascades. Furthermore, studies of alternative splicing can and have revealed the location of key domains on calcium channels that regulate their activity.

Investigators have shown that the structural changes generated by alternative splicing can affect channel biophysics, density, targeting, post-translational modification, and coupling to downstream signaling pathways. Functional sites of alternative splicing involve regulation that depends on tissue type, cell type, stage of development, and neuronal activity [7-22]. Importantly, the two sites of alternative splicing we discuss here, in the II-III intracellular loop region (e18a) and in the C-terminus (e37a and e37b), are conserved across a number of mammalian $\text{Ca}_v2.2$ genes including human, rat, and mouse (see Fig.1 for e18a sequence comparisons and see reference [7] for e37a and e37b comparisons). We reason that such sites of alternative splicing, that are under cellular control and that are evolutionarily conserved, must be located in key domains essential for regulating N-type channel function and likely evolved to contribute functional advantage to the organisms in which they occur [23,24].

The $\text{Ca}_v2.2$ N-type calcium channel

Our interest in the N-type calcium channel stems from its established role in the control of transmitter release from many different types of neurons [25,26]. The N-type calcium channel is also implicated in synaptogenesis and regulation of gene expression [27,28]. N-type calcium channels differ functionally across cell types and sub-cellular compartments within a given cell, suggestive of molecular and structural heterogeneity. Differences in N-type channel inactivation, single-channel conductance, gating, sensitivity to G-protein-mediated modulation, and ability to couple to transmitter release have been documented [29-39]. Several factors support functional diversity among voltage-gated calcium channels, including association with different auxiliary subunits, post-translational modifications, and association with target proteins and signaling molecules [40-47]. These processes are also controlled by alternative splicing [2,3,47-50].

Cassette exons in the II-III loops of Ca_v2 channels

The intracellular loops connecting homologous domains II and III ($\text{L}_{\text{II-III}}$) of pore-forming $\text{Ca}_v\alpha_1$ subunits are of special interest. This region supports key cellular functions and links calcium channels to downstream effector proteins. In skeletal muscle, the $\text{L}_{\text{II-III}}$ of $\text{Ca}_v1.1$ binds directly to ryanodine receptors on the sarcoplasmic reticulum, an interaction necessary and sufficient for excitation-contraction coupling [51-53]. In the case of Ca_v2 voltage-gated calcium channels, II-III loops connect channels to sites of synaptic vesicle fusion via binding to SNARE (soluble N-ethyl-maleimide-sensitive fusion (NSF) attachment protein receptor) proteins [54], critical for coupling N-type and P/Q-type calcium channels to transmitter release and for targeting channels to presynaptic nerve terminals [48,54-61]. The $\text{L}_{\text{II-III}}$ region of $\text{Ca}_v2.2$ channels also regulates cumulative inactivation of channels in response to stimulus trains [54,62,63].

Isoforms of $\text{Ca}_v2.1$, $\text{Ca}_v2.2$, and $\text{Ca}_v2.3$ exist that differ within the $\text{L}_{\text{II-III}}$ [16,64] and differ in their abilities to bind presynaptic SNARE proteins [48]. Our group has focused on cassette exons, e18a, located between exons 18 and 19 in $\text{L}_{\text{II-III}}$ of mammalian $\text{Ca}_v2.2$ and $\text{Ca}_v2.3$ genes that, as described below, are evolutionary conserved (Fig. 1). We showed previously that e18a in $\text{Ca}_v2.2$ protects the channel from inactivation. Specifically, e18a-containing $\text{Ca}_v2.2$ channels are less sensitive to closed-state inactivation, and therefore calcium influx during action

potential trains is expected to be sustained relative to that through Ca_v2.2 channels lacking e18a [62].

Human Ca_v2 genomic sequences

Our interest in e18a was motivated from our analyses of various Ca_v2 gene sequences. Mammalian Ca_v2.2, and Ca_v2.3, but not Ca_v2.1, genomic sequences contain a putative cassette exon between constitutive exons, e18 and e19. In rat, e18a resides within a large 6125 bp intron between e18 and e19 of the Ca_v2.2 gene [15]. We found a nearly identical cassette exon (61 of 63 bp), flanked by consensus gt-ag splice junctions, within the 9775 bp intron between e18 and e19 of the human Ca_v2.2 gene (see Fig. 1) [3]. We also identified the location of a 57 nucleotide exon within a 6221 bp intron of the human Ca_v2.3 gene consistent with previous studies of Ca_v2.3 mRNAs [65,66]. Despite similarity in size and position, exons 18a of Ca_v2.2 and Ca_v2.3 contain distinct nucleotide sequences with only 25% homology and encode dissimilar amino acid sequences (Fig. 1). A similar analysis of the Ca_v2.1 human gene revealed a much smaller intron (~ 1 kb) between e18 and e19, and no evidence for a putative cassette exon.

We also examined the genomic sequences of Ca_v2.2 and Ca_v2.3 for *Pan troglodytes*, *Bos taurus*, *Mus musculus*, *Canis familiaris*, and *Gallus gallus*. All these species contain the alternatively expressed e18a of Ca_v2.2 and Ca_v2.3 (Fig. 1B) and exon sequences were highly conserved. By our analysis intracellular regions of the channels, including the C-terminus and loops connecting homologous domains, were $71.8 \pm 3.0\%$ similar among species. In comparison, Ca_v2.2 e18a nucleotide sequences shared $96.2 \pm 0.5\%$ similarity among the seven species investigated here, significantly greater than the average sequence conservation within constitutively expressed exons that encode intracellular domains of the channel ($p < 0.0001$).

Expression of Ca_v2.2 and Ca_v2.3 alternatively spliced exons in the human nervous system

We next used RT-PCR to establish that e18a of Ca_v2.2 and Ca_v2.3 are expressed in human neuronal tissue, and we also confirmed the lack of alternative splicing in the same region of Ca_v2.1 (Fig. 2A). We found Ca_v2.2 e18a expression in human whole brain and spinal cord (Fig. 2A), contrary to reports that this exon is not expressed in human cells [12]. Notably, the proportion of Ca_v2.2 e18a-containing mRNAs was higher in adult spinal cord as compared to whole brain. Furthermore, Ca_v2.2 e18a was not detected in fetal whole brain. Thus, Ca_v2.2 e18a expression in the human is both regionally and developmentally regulated. Ca_v2.3 e18a is also alternatively spliced in human tissues (Fig. 2A) [66]. Interestingly, as Ca_v2.2 e18a increased during development, there appeared to be a concomitant decrease in Ca_v2.3 e18a expression. Inclusion of e18a in Ca_v2.2 results in N-type calcium channels that are less sensitive to closed-state inactivation [62]. In contrast, Ca_v2.3 channels that contain e18a exhibit increased sensitivity to calcium-dependent current enhancement [67,68] and stimulation by phorbol esters [69].

Developmental regulation of Ca_v2.2 e18a

We next examined the developmental expression of Ca_v2.2 exon 18a in the rat superior cervical ganglia, a tissue that expresses relatively high levels of Ca_v2.2 and that contains a relatively homogenous class of neurons (Fig. 2B) [15]. We quantified e18a splice isoforms by ribonuclease protection from P1, P5, P18, and adult superior cervical ganglia. We used an antisense probe that was complementary to constitutively expressed exons 15-18 and the alternatively spliced e18a. Ca_v2.2 mRNAs that contain e18a fully protected the probe (upper band, Fig. 2B), while Ca_v2.2 mRNAs that lack the alternatively spliced exon yielded a partially protected probe (lower band, Fig. 2B). In the adult ganglia, e18a was present in $67 \pm 4\%$ of all Ca_v2.2 mRNAs (Fig. 2B). By comparison, the relative abundance of e18a was $54 \pm 5\%$ at P18, $25 \pm 4\%$ at P5, and $16 \pm 4\%$ at P1 ($n = 3$ separate experiments for all ages). The adult expression

level of e18a is significantly different than that in P1 and P5 SCG ($P < 0.0003$), and the proportion of e18a mRNA transcripts increased >4-fold between P1 and adulthood. The overall expression level of $\text{Ca}_v2.2$ mRNA decreased between P1 and P18, in comparison to GAPDH expression levels (Fig. 2B, bottom gels). This is consistent with an increase in the number of non-neuronal cells in the maturing SCG.

During the first three weeks of postnatal life, while e18a expression continuously increased in our assay (Fig. 2), SCG neurons undergo a variety of developmental changes. By birth, the majority of SCG neurons have differentiated after migrating into the ganglia, and they begin to form synaptic connections with target structures soon after they arrive. The vast majority of these connections have formed by P21 [70]. Rat SCG neurons rely exclusively on the $\text{Ca}_v2.2$ N-type calcium channel to control transmitter release [71]. Our data, therefore, indicate that $\text{Ca}_v2.2$ N-type channels that contain the 21 amino acids encoded by e18a likely control transmitter release from mature SCG terminals and are specifically upregulated during the period of exuberant ganglion cell synaptogenesis for this purpose.

We note that alternatively spliced exons e18a within $\text{Ca}_v2.2$ and $\text{Ca}_v2.3$ show a high level of amino acid sequence conservation across mammalian and avian species (Fig. 1) when compared to constitutively expressed exons in this region of the genes. Others have also reported an exon homologous to mammalian e18a in the pufferfish $\text{Ca}_v2.2$ ortholog [72]. These data signify that additional evolutionary pressure exists to preserve e18a sequences and suggest an important functional role for $\text{Ca}_v2.2$ and $\text{Ca}_v2.3$ isoforms that contain the additional 21 and 19 amino acids encoded by their respective exons. Interestingly, our data point to the possibility that a common mechanism regulates expression of e18a exons in $\text{Ca}_v2.2$ and $\text{Ca}_v2.3$ based on their approximate reciprocal expression patterns. Specifically, we find that e18a containing $\text{Ca}_v2.2$ mRNAs are more abundant in the mature brain (Fig. 1), peripheral ganglia [15], and spinal cord (Fig. 1), whereas, $\text{Ca}_v2.3$ mRNAs containing e18a are more abundant in fetal brain (Fig. 1). Further, $\text{Ca}_v2.3$ e18a mRNAs are virtually absent from the peripheral nervous system (unpublished observations, ACG and DL). One tantalizing possibility is that e18a controls targeting of $\text{Ca}_v2.2$ and $\text{Ca}_v2.3$ channels to sites of transmitter release given that this exon modifies amino acid composition in the II-III linker region. Although e18a is absent from $\text{Ca}_v2.1$, others have reported distinct $\text{L}_{\text{II-III}}$ isoforms of this channel that might arise from alternative splicing of a different exon. These isoforms of $\text{Ca}_v2.1$ differ in their interactions with SNARE proteins and sub-cellular expression patterns [48,64]. Regulated expression of $\text{L}_{\text{II-III}}$ isoforms could represent a mechanism to limit overall Ca_v2 channel density at the synapse during maturation.

Mutually exclusive exons e37a and e37b in the C-terminus of $\text{Ca}_v2.2$

The C-terminus of Ca_v2

The C-termini of Ca_v2 genes coordinate various aspects of calcium channel function, including inactivation, modulation by G-proteins, modulation by calmodulin, and protein-protein interactions that regulate activity and/or target the channel to specific cellular compartments [9,37,47,49,73-76]. Alternative splicing in exon 46 of $\text{Ca}_v2.2$ affects the ability of G-proteins to modulate the N-type calcium channel [49] and alters sub-cellular targeting [47,73]. Mutually exclusive e37a and e37b are found in $\text{Ca}_v2.1$, $\text{Ca}_v2.2$, and $\text{Ca}_v2.3$ genes (Figure 3A). The amino acid compositions of these exons differ among the Ca_v2 genes, and the expression pattern of $\text{Ca}_v2.2$ e37a and e37b [7] is distinct from the reported expression pattern of the equivalent exons in $\text{Ca}_v2.1$ [9], suggesting that they serve different functions. Notably, we have shown that $\text{Ca}_v2.2\text{e}[37a]$ is preferentially expressed in capsaicin-responsive neurons of the dorsal root ganglia, implying a role in nociception [7].

Exon 37a is expressed preferentially in dorsal root ganglia

Ca_v2.2 channels localize to synapses of nociceptive neurons in the dorsal horn laminae I/II [77] and dominate in the control of transmitter release from these presynaptic terminals [78-80]. Mice lacking Ca_v2.2 have impaired nociception [81-84], and N-type channel blockers administered intrathecally, are powerful analgesics in a variety of animal pain models [85-89]. Some of the analgesic properties of morphine and other opiates are mediated via G-protein-dependent inhibition of N-type channels in the dorsal horn. A sub-set of nociceptive neurons preferentially express Ca_v2.2e[37a] mRNAs [7,90]. This novel Ca_v2.2 splice isoform supports larger N-type currents in nociceptive neurons, as compared to the more prevalent splice isoforms of Ca_v2.2 that contain the alternative e37b [7,91]. Larger currents associated with e37a-containing channels impact the amount of calcium entering cells in response to a variety of action potential waveforms [91]. More recently, we showed that e37a increases the sensitivity of the N-type channel to the voltage-independent form of G-protein modulation [92]. Collectively, our findings provide potentially important insights into the cellular mechanisms that control N-type channel activity in nociception and indicate that a sub-set of nociceptive neurons express a form of the N-type channel that is highly sensitive to inhibition by neurotransmitters and hormones.

G-protein-dependent modulation of N-type channel gating

Having shown preferential expression of e37a in nociceptors of dorsal root ganglia, we considered what features of N-type channels in these cells are especially prominent. Numerous G-protein-coupled neurotransmitter receptors suppress transmitter release from many neurons by inhibiting N-type calcium channels [30,44,45,93]. G-protein-dependent inhibition can be voltage-dependent and mediated by direct interaction of G-protein $\beta\gamma$ subunit dimers with Ca_v2.2 subunits [43,44,94]. This inhibition is membrane delimited, slows activation of the N-type current, and can be relieved by a strong depolarizing pre-pulse[44]. Voltage-dependent inhibition is most effective against calcium entry triggered by single action potentials. It is relieved by a train of action potentials [94-96]. Another form of G-protein-dependent inhibition of the N-type current is voltage-independent and this is an especially prominent feature of N-type channels in nociceptors[97]. This form of inhibition is not relieved by strong depolarizing pre-pulses and may involve kinase-dependent phosphorylation by protein kinase C [98] and tyrosine kinases [99]. The contributions of voltage-dependent and voltage-independent inhibition vary with transmitter receptor, cell type, and level of syntaxin 1A [43,44,100-103]. However, the molecular mechanisms that set the level and type of transmitter-mediated inhibition of the N-type channel are unknown. Our recent findings show that the degree and type of G protein-mediated inhibition of the N-type channel expressed in nociceptors is regulated by the alternative splice site, e37a and e37b, in the C-terminus of the Ca_v2.2 subunit.

Alternative splicing controls G protein–dependent inhibition of N-type calcium channels in nociceptors

We were motivated to test the role of e37a in G protein-mediated inhibition of the N-type channel because we had shown enrichment of e37a in nociceptors, where N-type channels are highly sensitive to inhibition by G protein-coupled receptors. We found that inclusion of e37a in Ca_v2.2 created an inhibitory pathway that was independent of voltage and that substantially increased the sensitivity of Ca_v2.2 channels to modulation by opiates and GABA [92]. Voltage-dependent inhibition was present in e37a and e37b-containing N-type channels and was unaffected by alternative splicing at this site in the C-terminus. We also showed that both forms of inhibition, voltage-dependent and voltage-independent, utilize PTX-sensitive G_i and G_o class G proteins, but the two pathways diverge downstream of the trimeric G protein. As described by many investigators, voltage-dependent inhibition requires the G $\beta\gamma$ dimer binding to the Ca_v2.2 channel and is independent of kinase activation [44,45] (Fig. 3B). By contrast,

voltage-independent inhibition proceeds independent of G $\beta\gamma$ and requires pp60c-src tyrosine kinase activation [92,99] (Fig. 3B). Interestingly, there is a tyrosine residue encoded in e37a and absent in e37b, that is essential for supporting voltage-independent inhibition of Ca $_v$ 2.2e [37a] channels (Fig. 3A). GABA, opiates and probably other transmitters and drugs can utilize this pathway to inhibit Ca $_v$ 2.2 channels by a mechanism that is independent of stimulus intensity [95,96,104] (Fig. 3B).

Our studies of alternative splicing illustrate how investigations of evolutionarily conserved, natural variants of Ca $_v$ 2.2 can uncover critical domains in N-type channels that regulate their activity. The e18a splice site illustrates that alternative splicing of certain exons in Ca $_v$ 2.2 is under developmental control and possibly coordinated with splicing of the homologous site in Ca $_v$ 2.3. Collectively, our studies suggest that N-type channels become less sensitive to inactivation following trains of stimuli during development. Alternative splicing of e37a on the other hand is under cell-specific control. Knowledge that e37a-containing Ca $_v$ 2.2 channels are preferentially expressed in nociceptors provided the functionally relevant context to understand the significance of our findings.

We still know very little of the factors that regulate alternative splicing of voltage-gated calcium channels. Of the proteins that have been identified that regulate neuron-specific alternative splicing, only the Nova2 RNA binding protein has been linked to Ca $_v$ 2.2 [105]. As factors that regulate expression of alternatively spliced exons in Ca $_v$ 2.2 channels are identified, we should begin to gain a better understanding of the mechanisms that exist to regulate calcium channel function and to control synaptic efficacy.

Acknowledgements

This work was supported by Howard Hughes Medical Institute Predoctoral Fellowship (A.C.G.) and National Institutes of Health Grants: NS29967 and NS055251 (D.L.).

Citations

1. Soldatov NM. Genomic structure of human L-type Ca $_v$ 2+ channel. *Genomics* 1994;22:77–87. [PubMed: 7959794]
2. Lipscombe D, Pan JQ, Gray AC. Functional diversity in neuronal voltage-gated calcium channels by alternative splicing of Ca(v)alpha1. *Mol Neurobiol* 2002;26:21–44. [PubMed: 12392054]
3. Lipscombe D, Castiglioni AJ. Alternative splicing in voltage gated calcium channels. In: McDonough, SI., editor. *Calcium Channel Pharmacology*. Kluwer Academic/Plenum; New York: 2004. p. 369–409.
4. Emerick MC, Stein R, Kunze R, McNulty MM, Regan MR, Hanck DA, Agnew WS. Profiling the array of Ca(v)3.1 variants from the human T-type calcium channel gene CACNA1G: alternative structures, developmental expression, and biophysical variations. *Proteins* 2006;64:320–42. [PubMed: 16671074]
5. Missler M, Sudhof TC. Neurexins: three genes and 1001 products. *Trends Genet* 1998;14:20–6. [PubMed: 9448462]
6. Schmucker D, Clemens JC, Shu H, Worby CA, Xiao J, Muda M, Dixon JE, Zipursky SL. Drosophila Dscam is an axon guidance receptor exhibiting extraordinary molecular diversity. *Cell* 2000;101:671–84. [PubMed: 10892653]
7. Bell TJ, Thaler C, Castiglioni AJ, Helton TD, Lipscombe D. Cell-specific alternative splicing increases calcium channel current density in the pain pathway. *Neuron* 2004;41:127–38. [PubMed: 14715140]
8. Angelotti T, Hofmann F. Tissue-specific expression of splice variants of the mouse voltage-gated calcium channel alpha2/delta subunit. *FEBS Lett* 1996;397:331–7. [PubMed: 8955374]
9. Bourinet E, Soong TW, Sutton K, Slaymaker S, Mathews E, Monteil A, Zamponi GW, Nargeot J, Snutch TP. Splicing of alpha 1A subunit gene generates phenotypic variants of P- and Q-type calcium channels. *Nat Neurosci* 1999;2:407–15. [PubMed: 10321243]

10. Diebold RJ, Koch WJ, Ellinor PT, Wang JJ, Muthuchamy M, Wieczorek DF, Schwartz A. Mutually exclusive exon splicing of the cardiac calcium channel alpha 1 subunit gene generates developmentally regulated isoforms in the rat heart. *Proc Natl Acad Sci U S A* 1992;89:1497–501. [PubMed: 1311102]
11. Gidh-Jain M, Huang B, Jain P, Battula V, el-Sherif N. Reemergence of the fetal pattern of L-type calcium channel gene expression in non infarcted myocardium during left ventricular remodeling. *Biochem Biophys Res Commun* 1995;216:892–7. [PubMed: 7488209]
12. Kaneko S, Cooper CB, Nishioka N, Yamasaki H, Suzuki A, Jarvis SE, Akaike A, Satoh M, Zamponi GW. Identification and characterization of novel human Ca(v)2.2 (alpha 1B) calcium channel variants lacking the synaptic protein interaction site. *J Neurosci* 2002;22:82–92. [PubMed: 11756491]
13. Lin Z, Haus S, Edgerton J, Lipscombe D. Identification of functionally distinct isoforms of the N-type Ca²⁺ channel in rat sympathetic ganglia and brain. *Neuron* 1997;18:153–66. [PubMed: 9010213]
14. Lin Z, Lin Y, Schorge S, Pan JQ, Beierlein M, Lipscombe D. Alternative splicing of a short cassette exon in alpha1B generates functionally distinct N-type calcium channels in central and peripheral neurons. *J Neurosci* 1999;19:5322–31. [PubMed: 10377343]
15. Pan JQ, Lipscombe D. Alternative splicing in the cytoplasmic II-III loop of the N-type Ca channel alpha 1B subunit: functional differences are beta subunit-specific. *J Neurosci* 2000;20:4769–75. [PubMed: 10864934]
16. Soong TW, DeMaria CD, Alvania RS, Zweifel LS, Liang MC, Mittman S, Agnew WS, Yue DT. Systematic identification of splice variants in human P/Q-type channel alpha1(2.1) subunits: implications for current density and Ca²⁺-dependent inactivation. *J Neurosci* 2002;22:10142–52. [PubMed: 12451115]
17. Vignes S, Chabret C, Valentin S, Valmier J. Rat embryonic hippocampal neurons express a new class A calcium channel variant. *Neurosci Lett* 1998;258:37–40. [PubMed: 9876046]
18. Vignes S, Gastaldi M, Chabret C, Massacrier A, Cau P, Valmier J. Regulation of calcium channel alpha(1A) subunit splice variant mRNAs in kainate-induced temporal lobe epilepsy. *Neurobiol Dis* 1999;6:288–301. [PubMed: 10448056]
19. Welling A, Ludwig A, Zimmer S, Klugbauer N, Flockerzi V, Hofmann F. Alternatively spliced IS6 segments of the alpha 1C gene determine the tissue-specific dihydropyridine sensitivity of cardiac and vascular smooth muscle L-type Ca²⁺ channels. *Circ Res* 1997;81:526–32. [PubMed: 9314833]
20. Takimoto K, Li D, Hershman KM, Li P, Jackson EK, Levitan ES. Decreased expression of Kv4.2 and novel Kv4.3 K⁺ channel subunit mRNAs in ventricles of renovascular hypertensive rats. *Circ Res* 1997;81:533–9. [PubMed: 9314834]
21. Vendel AC, Terry MD, Striegel AR, Iverson NM, Leuranguer V, Rithner CD, Lyons BA, Pickard GE, Tobet SA, Horne WA. Alternative splicing of the voltage-gated Ca²⁺ channel beta4 subunit creates a uniquely folded N-terminal protein binding domain with cell-specific expression in the cerebellar cortex. *J Neurosci* 2006;26:2635–44. [PubMed: 16525042]
22. Yang Y, Chen X, Margulies K, Jeevanandam V, Pollack P, Bailey BA, Houser SR. L-type Ca²⁺ channel alpha 1c subunit isoform switching in failing human ventricular myocardium. *J Mol Cell Cardiol* 2000;32:973–84. [PubMed: 10888251]
23. Lareau LF, Green RE, Bhatnagar RS, Brenner SE. The evolving roles of alternative splicing. *Curr Opin Struct Biol* 2004;14:273–82. [PubMed: 15193306]
24. Lipscombe D. Neuronal proteins custom designed by alternative splicing. *Curr Opin Neurobiol* 2005;15:358–63. [PubMed: 15961039]
25. Reid CA, Bekkers JM, Clements JD. Presynaptic Ca²⁺ channels: a functional patchwork. *Trends Neurosci* 2003;26:683–7. [PubMed: 14624853]
26. Dunlap K, Luebke JI, Turner TJ. Exocytotic Ca²⁺ channels in mammalian central neurons. *Trends Neurosci* 1995;18:89–98. [PubMed: 7537420]
27. Brosenitsch TA, Katz DM. Physiological patterns of electrical stimulation can induce neuronal gene expression by activating N-type calcium channels. *J Neurosci* 2001;21:2571–9. [PubMed: 11306610]

28. Jones OT, Bernstein GM, Jones EJ, Jugloff DG, Law M, Wong W, Mills LR. N-Type calcium channels in the developing rat hippocampus: subunit, complex, and regional expression. *J Neurosci* 1997;17:6152–64. [PubMed: 9236227]
29. Plummer MR, Logothetis DE, Hess P. Elementary properties and pharmacological sensitivities of calcium channels in mammalian peripheral neurons. *Neuron* 1989;2:1453–63. [PubMed: 2560643]
30. Lipscombe D, Kongsamut S, Tsien RW. Alpha-adrenergic inhibition of sympathetic neurotransmitter release mediated by modulation of N-type calcium-channel gating. *Nature* 1989;340:639–42. [PubMed: 2570354]
31. Delcour AH, Lipscombe D, Tsien RW. Multiple modes of N-type calcium channel activity distinguished by differences in gating kinetics. *J Neurosci* 1993;13:181–94. [PubMed: 8380849]
32. Rittenhouse AR, Hess P. Microscopic heterogeneity in unitary N-type calcium currents in rat sympathetic neurons. *J Physiol* 1994;474:87–99. [PubMed: 8014899]
33. Jones SW, Marks TN. Calcium currents in bullfrog sympathetic neurons. I. Activation kinetics and pharmacology. *J Gen Physiol* 1989;94:151–67. [PubMed: 2478659]
34. Plummer MR, Hess P. Reversible uncoupling of inactivation in N-type calcium channels. *Nature* 1991;351:657–9. [PubMed: 1646965]
35. Cox DH, Dunlap K. Inactivation of N-type calcium current in chick sensory neurons: calcium and voltage dependence. *J Gen Physiol* 1994;104:311–36. [PubMed: 7807051]
36. Delmas P, Abogadie FC, Buckley NJ, Brown DA. Calcium channel gating and modulation by transmitters depend on cellular compartmentalization. *Nat Neurosci* 2000;3:670–8. [PubMed: 10862699]
37. Beedle AM, McRory JE, Poirot O, Doering CJ, Altier C, Barrere C, Hamid J, Nargeot J, Bourinet E, Zamponi GW. Agonist-independent modulation of N-type calcium channels by ORL1 receptors. *Nat Neurosci* 2004;7:118–25. [PubMed: 14730309]
38. Poncer JC, McKinney RA, Gahwiler BH, Thompson SM. Either N- or P-type calcium channels mediate GABA release at distinct hippocampal inhibitory synapses. *Neuron* 1997;18:463–72. [PubMed: 9115739]
39. Poncer JC, McKinney RA, Gahwiler BH, Thompson SM. Differential control of GABA release at synapses from distinct interneurons in rat hippocampus. *J Physiol* 2000;528(Pt 1):123–30. [PubMed: 11018111]
40. Birnbaumer L, Qin N, Olcese R, Tareilus E, Platano D, Costantin J, Stefani E. Structures and functions of calcium channel beta subunits. *J Bioenerg Biomembr* 1998;30:357–75. [PubMed: 9758332]
41. Klugbauer N, Marais E, Hofmann F. Calcium channel alpha2delta subunits: differential expression, function, and drug binding. *J Bioenerg Biomembr* 2003;35:639–47. [PubMed: 15000524]
42. Walker D, De Waard M. Subunit interaction sites in voltage-dependent Ca²⁺ channels: role in channel function. *Trends Neurosci* 1998;21:148–54. [PubMed: 9554724]
43. Strock J, Diverse-Pierluissi MA. Ca²⁺ channels as integrators of G protein-mediated signaling in neurons. *Mol Pharmacol* 2004;66:1071–6. [PubMed: 15269290]
44. Ikeda SR, Dunlap K. Voltage-dependent modulation of N-type calcium channels: role of G protein subunits. *Adv Second Messenger Phosphoprotein Res* 1999;33:131–51. [PubMed: 10218117]
45. Hille B, Beech DJ, Bernheim L, Mathie A, Shapiro MS, Wollmuth LP. Multiple G-protein-coupled pathways inhibit N-type Ca channels of neurons. *Life Sci* 1995;56:989–92. [PubMed: 10188803]
46. Striessnig J. Targeting voltage-gated Ca²⁺ channels. *Lancet* 2001;357:1294. [PubMed: 11421217]
47. Maximov A, Bezprozvanny I. Synaptic targeting of N-type calcium channels in hippocampal neurons. *J Neurosci* 2002;22:6939–52. [PubMed: 12177192]
48. Rettig J, Sheng ZH, Kim DK, Hodson CD, Snutch TP, Catterall WA. Isoform-specific interaction of the alpha1A subunits of brain Ca²⁺ channels with the presynaptic proteins syntaxin and SNAP-25. *Proc Natl Acad Sci U S A* 1996;93:7363–8. [PubMed: 8692999]
49. Simen AA, Lee CC, Simen BB, Bindokas VP, Miller RJ. The C terminus of the Ca channel alpha1B subunit mediates selective inhibition by G-protein-coupled receptors. *J Neurosci* 2001;21:7587–97. [PubMed: 11567049]

50. Sakurai T, Hell JW, Woppmann A, Miljanich GP, Catterall WA. Immunochemical identification and differential phosphorylation of alternatively spliced forms of the alpha 1A subunit of brain calcium channels. *J Biol Chem* 1995;270:21234–42. [PubMed: 7673157]
51. Rios E, Brum G. Involvement of dihydropyridine receptors in excitation-contraction coupling in skeletal muscle. *Nature* 1987;325:717–20. [PubMed: 2434854]
52. Mori Y, Niidome T, Fujita Y, Mynlieff M, Dirksen RT, Beam KG, Iwabe N, Miyata T, Furutama D, Furuichi T, et al. Molecular diversity of voltage-dependent calcium channel. *Ann N Y Acad Sci* 1993;707:87–108. [PubMed: 9137545]
53. Flucher BE, Franzini-Armstrong C. Formation of junctions involved in excitation-contraction coupling in skeletal and cardiac muscle. *Proc Natl Acad Sci U S A* 1996;93:8101–6. [PubMed: 8755610]
54. Catterall WA. Interactions of presynaptic Ca²⁺ channels and snare proteins in neurotransmitter release. *Ann N Y Acad Sci* 1999;868:144–59. [PubMed: 10414292]
55. Leveque C, el Far O, Martin-Moutot N, Sato K, Kato R, Takahashi M, Seagar MJ. Purification of the N-type calcium channel associated with syntaxin and synaptotagmin. A complex implicated in synaptic vesicle exocytosis. *J Biol Chem* 1994;269:6306–12. [PubMed: 8119979]
56. Sheng ZH, Rettig J, Takahashi M, Catterall WA. Identification of a syntaxin-binding site on N-type calcium channels. *Neuron* 1994;13:1303–13. [PubMed: 7993624]
57. Tobi D, Wiser O, Trus M, Atlas D. N-type voltage-sensitive calcium channel interacts with syntaxin, synaptotagmin and SNAP-25 in a multiprotein complex. *Receptors Channels* 1998;6:89–98. [PubMed: 9932286]
58. Mochida S, Westenbroek RE, Yokoyama CT, Zhong H, Myers SJ, Scheuer T, Itoh K, Catterall WA. Requirement for the synaptic protein interaction site for reconstitution of synaptic transmission by P/Q-type calcium channels. *Proc Natl Acad Sci U S A* 2003;100:2819–24. [PubMed: 12601156]
59. Mochida S, Sheng ZH, Baker C, Kobayashi H, Catterall WA. Inhibition of neurotransmission by peptides containing the synaptic protein interaction site of N-type Ca²⁺ channels. *Neuron* 1996;17:781–8. [PubMed: 8893034]
60. Rettig J, Heinemann C, Ashery U, Sheng ZH, Yokoyama CT, Catterall WA, Neher E. Alteration of Ca²⁺ dependence of neurotransmitter release by disruption of Ca²⁺ channel/syntaxin interaction. *J Neurosci* 1997;17:6647–56. [PubMed: 9254677]
61. Harkins AB, Cahill AL, Powers JF, Tischler AS, Fox AP. Deletion of the synaptic protein interaction site of the N-type (Cav2.2) calcium channel inhibits secretion in mouse pheochromocytoma cells. *Proc Natl Acad Sci U S A* 2004;101:15219–24. [PubMed: 15471993]
62. Thaler C, Gray AC, Lipscombe D. Cumulative inactivation of N-type Ca_v2.2 calcium channels modified by alternative splicing. *Proc Natl Acad Sci U S A* 2004;101:5675–9. [PubMed: 15060274]
63. Degtiar VE, Scheller RH, Tsien RW. Syntaxin modulation of slow inactivation of N-type calcium channels. *J Neurosci* 2000;20:4355–67. [PubMed: 10844004]
64. Sakurai T, Westenbroek RE, Rettig J, Hell J, Catterall WA. Biochemical properties and subcellular distribution of the BI and rB isoforms of alpha 1A subunits of brain calcium channels. *J Cell Biol* 1996;134:511–28. [PubMed: 8707834]
65. Pereverzev A, Klockner U, Henry M, Grabsch H, Vajna R, Olyschlager S, Viatchenko-Karpinski S, Schroder R, Hescheler J, Schneider T. Structural diversity of the voltage-dependent Ca²⁺ channel alpha1E-subunit. *Eur J Neurosci* 1998;10:916–25. [PubMed: 9753159]
66. Williams ME, Marubio LM, Deal CR, Hans M, Brust PF, Philipson LH, Miller RJ, Johnson EC, Harpold MM, Ellis SB. Structure and functional characterization of neuronal alpha 1E calcium channel subtypes. *J Biol Chem* 1994;269:22347–57. [PubMed: 8071363]
67. Pereverzev A, Leroy J, Krieger A, Malecot CO, Hescheler J, Pfitzer G, Klockner U, Schneider T. Alternate splicing in the cytosolic II-III loop and the carboxy terminus of human E-type voltage-gated Ca(2+) channels: electrophysiological characterization of isoforms. *Mol Cell Neurosci* 2002;21:352–65. [PubMed: 12401453]
68. Leroy J, Pereverzev A, Vajna R, Qin N, Pfitzer G, Hescheler J, Malecot CO, Schneider T, Klockner U. Ca²⁺-sensitive regulation of E-type Ca²⁺ channel activity depends on an arginine-rich region in the cytosolic II-III loop. *Eur J Neurosci* 2003;18:841–55. [PubMed: 12925010]

69. Klockner U, Pereverzev A, Leroy J, Krieger A, Vajna R, Pfitzer G, Hescheler J, Malecot CO, Schneider T. The cytosolic II-III loop of Cav2.3 provides an essential determinant for the phorbol ester-mediated stimulation of E-type Ca²⁺ channel activity. *Eur J Neurosci* 2004;19:2659–68. [PubMed: 15147300]
70. Vidovic M, Hill CE, Hendry IA. Developmental time course of the sympathetic postganglionic innervation of the rat eye. *Brain Res* 1987;429:133–8. [PubMed: 3567657]
71. Hirning LD, Fox AP, McCleskey EW, Olivera BM, Thayer SA, Miller RJ, Tsien RW. Dominant role of N-type Ca²⁺ channels in evoked release of norepinephrine from sympathetic neurons. *Science* 1988;239:57–61. [PubMed: 2447647]
72. Minovitsky S, Gee SL, Schokrpur S, Dubchak I, Conboy JG. The splicing regulatory element, UGCAUG, is phylogenetically and spatially conserved in introns that flank tissue-specific alternative exons. *Nucleic Acids Res* 2005;33:714–24. [PubMed: 15691898]
73. Maximov A, Sudhof TC, Bezprozvanny I. Association of neuronal calcium channels with modular adaptor proteins. *J Biol Chem* 1999;274:24453–6. [PubMed: 10455105]
74. Hibino H, Pironkova R, Onwumere O, Vologodskaya M, Hudspeth AJ, Lesage F. RIM binding proteins (RBPs) couple Rab3-interacting molecules (RIMs) to voltage-gated Ca(2+) channels. *Neuron* 2002;34:411–23. [PubMed: 11988172]
75. Krovetz HS, Helton TD, Crews AL, Horne WA. C-Terminal alternative splicing changes the gating properties of a human spinal cord calcium channel alpha 1A subunit. *J Neurosci* 2000;20:7564–70. [PubMed: 11027215]
76. Liang H, DeMaria CD, Erickson MG, Mori MX, Alseikhan BA, Yue DT. Unified mechanisms of Ca²⁺ regulation across the Ca²⁺ channel family. *Neuron* 2003;39:951–60. [PubMed: 12971895]
77. Kerr LM, Filloux F, Olivera BM, Jackson H, Wamsley JK. Autoradiographic localization of calcium channels with [125I]omega-conotoxin in rat brain. *Eur J Pharmacol* 1988;146:181–3. [PubMed: 2450766]
78. Maggi CA, Tramontana M, Cecconi R, Santicoli P. Neurochemical evidence for the involvement of N-type calcium channels in transmitter secretion from peripheral endings of sensory nerves in guinea pigs. *Neurosci Lett* 1990;114:203–6. [PubMed: 1697665]
79. Holz, GG; Dunlap, K.; Kream, RM. Characterization of the electrically evoked release of substance P from dorsal root ganglion neurons: methods and dihydropyridine sensitivity. *J Neurosci* 1988;8:463–71. [PubMed: 2448433]
80. Heinke B, Balzer E, Sandkuhler J. Pre- and postsynaptic contributions of voltage-dependent Ca²⁺ channels to nociceptive transmission in rat spinal lamina I neurons. *Eur J Neurosci* 2004;19:103–11. [PubMed: 14750968]
81. Hatakeyama S, Wakamori M, Ino M, Miyamoto N, Takahashi E, Yoshinaga T, Sawada K, Imoto K, Tanaka I, Yoshizawa T, Nishizawa Y, Mori Y, Niidome T, Shoji S. Differential nociceptive responses in mice lacking the alpha(1B) subunit of N-type Ca(2+) channels. *Neuroreport* 2001;12:2423–7. [PubMed: 11496122]
82. Kim C, Jun K, Lee T, Kim SS, McEnery MW, Chin H, Kim HL, Park JM, Kim DK, Jung SJ, Kim J, Shin HS. Altered nociceptive response in mice deficient in the alpha(1B) subunit of the voltage-dependent calcium channel. *Mol Cell Neurosci* 2001;18:235–45. [PubMed: 11520183]
83. Saegusa H, Kurihara T, Zong S, Kazuno A, Matsuda Y, Nonaka T, Han W, Toriyama H, Tanabe T. Suppression of inflammatory and neuropathic pain symptoms in mice lacking the N-type Ca²⁺ channel. *Embo J* 2001;20:2349–56. [PubMed: 11350923]
84. Saegusa H, Matsuda Y, Tanabe T. Effects of ablation of N- and R-type Ca(2+) channels on pain transmission. *Neurosci Res* 2002;43:1–7. [PubMed: 12074836]
85. Bowersox SS, Gadbois T, Singh T, Pettus M, Wang YX, Luther RR. Selective N-type neuronal voltage-sensitive calcium channel blocker, SNX-111, produces spinal antinociception in rat models of acute, persistent and neuropathic pain. *J Pharmacol Exp Ther* 1996;279:1243–9. [PubMed: 8968347]
86. Bowersox SS, Luther R. Pharmacotherapeutic potential of omega-conotoxin MVIIA (SNX-111), an N-type neuronal calcium channel blocker found in the venom of *Conus magus*. *Toxicon* 1998;36:1651–8. [PubMed: 9792182]

87. Bowersox SS, Miljanich GP, Sugiura Y, Li C, Nadasdi L, Hoffman BB, Ramachandran J, Ko CP. Differential blockade of voltage-sensitive calcium channels at the mouse neuromuscular junction by novel omega-conopeptides and omega-agatoxin-IVA. *J Pharmacol Exp Ther* 1995;273:248–56. [PubMed: 7714772]
88. Matthews EA, Dickenson AH. Effects of spinally delivered N- and P-type voltage-dependent calcium channel antagonists on dorsal horn neuronal responses in a rat model of neuropathy. *Pain* 2001;92:235–46. [PubMed: 11323145]
89. Vanegas H, Schaible H. Effects of antagonists to high-threshold calcium channels upon spinal mechanisms of pain, hyperalgesia and allodynia. *Pain* 2000;85:9–18. [PubMed: 10692598]
90. Zamponi GW, McCleskey EW. Splicing it up. A variant of the N-type calcium channel specific for pain. *Neuron* 2004;41:3–4. [PubMed: 14715128]
91. Castiglioni AJ, Raingo J, Lipscombe D. Alternative splicing in the C-terminus of Ca_v2.2 controls expression and gating of N-type calcium channels. *J Physiol* 2006;576:119–34. [PubMed: 16857708]
92. Raingo J, Castiglioni AJ, Lipscombe D. Alternative splicing controls G protein-dependent inhibition of N-type calcium channels in nociceptors. *Nature Neuroscience*. 2007In press
93. Dunlap K, Fischbach GD. Neurotransmitters decrease the calcium component of sensory neurone action potentials. *Nature* 1978;276:837–9. [PubMed: 31570]
94. Elmslie KS. Neurotransmitter modulation of neuronal calcium channels. *J Bioenerg Biomembr* 2003;35:477–89. [PubMed: 15000517]
95. Park D, Dunlap K. Dynamic regulation of calcium influx by G-proteins, action potential waveform, and neuronal firing frequency. *J Neurosci* 1998;18:6757–66. [PubMed: 9712647]
96. Brody DL, Patil PG, Mülle JG, Snutch TP, Yue DT. Bursts of action potential waveforms relieve G-protein inhibition of recombinant P/Q-type Ca²⁺ channels in HEK 293 cells. *J Physiol* 1997;499(Pt 3):637–44. [PubMed: 9130160]
97. Diverse-Pierluissi M, Dunlap K. Distinct, convergent second messenger pathways modulate neuronal calcium currents. *Neuron* 1993;10:753–60. [PubMed: 8097399]
98. Rane SG, Dunlap K. Kinase C activator 1,2-oleoylacetyl glycerol attenuates voltage-dependent calcium current in sensory neurons. *Proc Natl Acad Sci U S A* 1986;83:184–8. [PubMed: 2417236]
99. Diverse-Pierluissi M, Remmers AE, Neubig RR, Dunlap K. Novel form of crosstalk between G protein and tyrosine kinase pathways. *Proc Natl Acad Sci U S A* 1997;94:5417–21. [PubMed: 9144252]
100. Jarvis SE, Magga JM, Beedle AM, Braun JE, Zamponi GW. G protein modulation of N-type calcium channels is facilitated by physical interactions between syntaxin 1A and Gbetagamma. *J Biol Chem* 2000;275:6388–94. [PubMed: 10692440]
101. Jarvis SE, Zamponi GW. Distinct molecular determinants govern syntaxin 1A-mediated inactivation and G-protein inhibition of N-type calcium channels. *J Neurosci* 2001;21:2939–48. [PubMed: 11312277]
102. Lu Q, AtKisson MS, Jarvis SE, Feng ZP, Zamponi GW, Dunlap K. Syntaxin 1A supports voltage-dependent inhibition of alpha1B Ca²⁺ channels by Gbetagamma in chick sensory neurons. *J Neurosci* 2001;21:2949–57. [PubMed: 11312278]
103. Stanley EF, Mirotznik RR. Cleavage of syntaxin prevents G-protein regulation of presynaptic calcium channels. *Nature* 1997;385:340–3. [PubMed: 9002518]
104. Brody DL, Yue DT. Relief of G-protein inhibition of calcium channels and short-term synaptic facilitation in cultured hippocampal neurons. *J Neurosci* 2000;20:889–98. [PubMed: 10648693]
105. Ule J, Jensen KB, Ruggiu M, Mele A, Ule A, Darnell RB. CLIP identifies Nova-regulated RNA networks in the brain. *Science* 2003;302:1212–5. [PubMed: 14615540]

Figure 1.

A, Alternatively spliced cassette exons, e18a, are found within human $Ca_v2.2$ and $Ca_v2.3$ genes. Partial gene structures, spanning e17 through e20, of human $Ca_v2.1$, $Ca_v2.2$, and $Ca_v2.3$ are illustrated. Exons are illustrated as gray (constitutive exons) or blue (alternatively spliced exons, e18a of $Ca_v2.2$ and $Ca_v2.3$) boxes, and introns are depicted as lines. Alternate 5' donor ($Ca_v2.1$ e17) and 3' acceptor ($Ca_v2.2$ e19) sites are indicated by superscript 'a'. We performed BLAST searches of the publicly available human genome sequence with published mRNA sequences for $Ca_v2.1$ (Accession #X99897), $Ca_v2.2$ (#M94172, #AF222338), and $Ca_v2.3$ (#L27745). Intron and exon sizes are drawn to scale, as indicated. **B**, Partial alignments of the deduced amino acid sequence of $Ca_v2.2$ (top) and $Ca_v2.3$ (bottom), including the alternatively spliced exon, e18a (blue and white), show limited variability across species. Abutting sequence of e18 and e19 (gray and black) are also included. $Ca_v2.2$ gene sequences were identified from the following depositions: *P. troglodytes*, Accession #NW_111847; *B. Taurus*, #NW_001024540; *C. familiaris*, #NW_876333; *R. rattus*, #NW_047651; *M. musculus*, #NT_039206; *G. gallus*, #NC_006104. $Ca_v2.3$ genes were found in the following sequence depositions: *P. troglodytes*, Accession #NW_101951; *B. Taurus*, #NW_972257; *C. familiaris*, #NW_876323; *R. rattus*, #NW_047398; *M. musculus*, # NT_039184; *G. gallus*, #NC_006095.

Figure 2.

A, Regulated alternative splicing of Ca_v2.2 and Ca_v2.3 e18a occurs in the human central nervous system. Using primers that flank e18a, we performed RT-PCR for Ca_v2.1, Ca_v2.2, and Ca_v2.3 with RNA preparations from human fetal and adult brain as well as human adult spinal cord. RT-PCR of Ca_v2.1 (*top*) in this region results in one product of the appropriate size (455 bp). Ca_v2.2 primers (*middle*) produced a single product from fetal brain, compared to two products that differ by 63 bps from adult brain and spinal cord (+e18a, 351 bp; Δe18a, 288 bp). Two products (+e18a, 508 bp; Δe18a, 451 bp) result from the Ca_v2.3 PCR (*bottom*) for all three RNA sources. **B**, Ca_v2.2 e18a expression increases during the first three postnatal weeks in rat SCG. We examined the developmental expression of e18a by RNase Protection Assay with the RPA III Kit (Ambion) and biotinylated antisense probes. *Top*, histogram indicates the percent of e18a-containing mRNA (upper band) as compared to the total signal from protected fragments for P1, P5, P18, and adult SCG RNA. Compared to P1 and P5 SCG, P18 and Adult SCG contain significantly greater proportions of e18a-containing Ca_v2.2 mRNA ($p < 0.002$). *Bottom*, representative gels showing typical results for three probes: *top* - a probe designed to examine alternative splicing of e18a (240 min exposure), *middle* - pan- Ca_v2.2 probe (180 min exposure), *bottom* - GAPDH probe (10 min exposure). In the upper gel, the fully protected product contains e18a (381 bp; gray and blue bar), and the partially protected product (lower band) lacks e18a (318 bp; gray bar). GAPDH and pan- Ca_v2.2 probes were hybridized to SCG RNA in the same RPA reaction. P18 and adult Ca_v2.2 expression levels normalized to GAPDH are lower than those at P1 (Ca_v2.2: GAPDH ratios: P1 = 0.71 ± 0.06 , P5 = 0.61 ± 0.10 , P18 = 0.37 ± 0.09 , adult = 0.30 ± 0.07 , mean $\pm$ SE; $p < 0.05$).

Figure 3.

A, Amino acid sequence alignments of exons e37a and e37b of human Ca_v2.1, Ca_v2.2, and Ca_v2.3 genes. Accession numbers were: AC011446 (Ca_v2.1), AY211499 (Ca_v2.2e[37a]), AF055477 (Ca_v2.2e[37b]), and AF239258S3 (Ca_v2.3). Red font labels amino acids that differ between e37a and e37b. A tyrosine kinase consensus site is indicated by asterisk at position 1747 in Ca_v2.2e[37a]. Tyrosine 1747 is essential for G-protein mediated voltage-independent inhibition of N-type Ca currents. **B**, Schematic showing the two inhibitory pathways that couple to N-type channels in nociceptors. The voltage-dependent inhibitory pathway couples to both Ca_v2.2e[37a] and Ca_v2.2e[37b] channels. Voltage-dependent inhibition utilizes pertussis toxin-sensitive G proteins and requires Gβγ dimers. Voltage-independent inhibition only couples to Ca_v2.2e[37a] channels. This pathway uses pertussis toxin-sensitive G proteins, is independent of Gβγ dimers, depends on *src* pp60 tyrosine kinase, and requires a tyrosine at position 1747 in Ca_v2.2.