LETTER TO THE EDITOR

On the age-dependent association between cancer of the breast and endometrium

Sir – In their recent paper on their follow-up (cohort) study of 60,065 newly diagnosed breast cancer patients, Adami *et al.* (1987) reported that 260 endometrial cancers had been diagnosed after the breast cancer diagnosis compared to an expected number, based on national rates, of 151.1 (relative risk = $260/151.1 = 1.72$). This relative risk for the subsequent incidence of endometrial cancer in breast cancer patients increased from 0.95 in patients diagnosed with breast cancer before age 40 to 2.47 in patients diagnosed at 80 years of age or older. The authors go on from this to state that 'there is no obvious common risk factor related to reproductive life which could give a reasonable explanation for [this observed] age-dependent association between cancer of the breast and cancer of the endometrium', and imply that this age dependence is evidence against the 'traditional... paradigm' of seeking the causes of (and link between) these two cancers in terms of 'hormonal expressions of characteristics of reproductive life or... dietary habits'. Their results are, however, in good agreement with what we would predict from our current understanding of the (hormonal) epidemiology of these two cancers.

There are a number of well-established risk factors for endometrial cancer. Low parity, late menopause and oestrogen replacement therapy (ERT) all increase the risk of endometrial cancer, while use of combination-type oral contraceptives (COCs) decreases the risk of the disease [see Kelsey *et al.* (1982) and Henderson *et al.* (1982) for references]. Obesity is a major risk factor both in pre-menopausal and post-menopausal women [see La Vecchia *et al.* (1982) and Henderson *et al.* (1983) for references]. These risk factors are all explicable in terms of the 'unopposed oestrogen hypothesis' for endometrial cancer (Siiteri, 1978; Niskier *et al.*, 1980; Henderson *et al.*, 1982).

Late age at first birth, late age at menopause and use of ERT all increase the risk of breast cancer (Thomas, 1984; Pike & Ross, 1984). We noted above that two of these three breast cancer risk factors, *viz.* late menopause and ERT, are also risk factors for endometrial cancer; late first birth is too, since late first birth is correlated with low parity. COC use does not, however, protect against breast cancer, and may in some circumstances increase the risk of breast cancer (McPherson & Drife, 1986). The sharp distinction between the protective effect of COC use against endometrial cancer and the lack of any such effect on breast cancer risk clearly shows that the endocrinology of endometrial cancer and breast cancer are not the same. Most important for an understanding of the results of Adami *et al.* (1987) is the finding that obesity is only associated with an increased risk of breast cancer in older post-menopausal women (Kelsey *et al.*, 1981; Lubin *et al.*, 1985). In pre-menopausal women obesity has been found, in fact, to be associated with a decreased risk of breast cancer (Willett *et al.*, 1985).

In the pre-menopausal period obesity acts contrariwise as a risk factor in endometrial cancer and breast cancer. Pre-menopausal breast cancer patients will tend to be of low parity but non-obese, so that their risk of endometrial cancer will probably be little different from expected as their low parity will slightly increase their risk while their non-obesity

will decrease it. Post-menopausal breast cancer patients will tend to be of low parity and of late menopause, and to be obese: their risk of endometrial cancer will be increased (relative to expected) on all counts. The association of breast cancer with obesity only becomes clear five to ten years after menopause (Kelsey *et al.*, 1981; Lubin *et al.*, 1985) when the protective effect evident in the pre-menopausal period has had time to be reversed, so that the greatly increased risk of endometrial cancer associated with obesity will become evident only in older post-menopausal breast cancer patients. This is precisely what Adami *et al.* (1987) observed.

It may be argued that the above reasoning will not hold in Sweden because Adami *et al.* (1977) did not find any relationship between weight and breast cancer in post-menopausal Swedish women. We have no explanation for their negative results, nor for the fact that Adami *et al.* (1978) also reported finding no significant relationship between any reproductive factor and risk of breast cancer in their study population. However, since the study we are discussing here (Adami *et al.*, 1987) has no data on body weight, we feel that it is reasonable to explain the results in terms of our general understanding of breast cancer.

Yours etc.,

T. Key and M.C. Pike,
ICRF Cancer Epidemiology Unit,
Gibson Building,
Radcliffe Infirmary,
Oxford OX2 6HE, UK.

References

- ADAMI, H.-O., RIMSTEN, A., STENKVIST, B. & VEGELIUS, J. (1977). Influence of height, weight and obesity on risk of breast cancer in an unselected Swedish population. *Br. J. Cancer*, **36**, 787.
- ADAMI, H.-O., RIMSTEN, A., STENKVIST, B. & VEGELIUS, J. (1978). Reproductive history and risk of breast cancer. A case-control study in an unselected Swedish population. *Cancer*, **41**, 747.
- ADAMI, H.-O., KRUSEMO, U.B., BERGKVIST, L., PERSSON, I. & PETTERSSON, B. (1987). On the age-dependent association between cancer of the breast and of the endometrium. A nationwide cohort study. *Br. J. Cancer*, **55**, 77.
- HENDERSON, B.E., ROSS, R.K., PIKE, M.C. & CASAGRANDE, J.T. (1982). Endogenous hormones as a major factor in human cancer. *Cancer Res.*, **42**, 3232.
- HENDERSON, B.E., CASAGRANDE, J.T., PIKE, M.C., MACK, T., ROSARIO, I. & DUKE, A. (1983). The epidemiology of endometrial cancer in young women. *Br. J. Cancer*, **47**, 749.
- KELSEY, J.L., FISCHER, D.B., HOLFORD, T.R. & 4 others (1981). Exogenous estrogens and other factors in the epidemiology of breast cancer. *J. Natl Cancer Inst.*, **67**, 327.
- KELSEY, J.L., LI VOLSI, V.A., HOLFORD, T.R. & 5 others (1982). A case-control study of cancer of the endometrium. *Am. J. Epidemiol.*, **116**, 333.
- LA VECCHIA, C., FRANCESCHI, S., GALLUS, G. & 4 others (1982). Oestrogens and obesity as risk factors for endometrial cancer in Italy. *Int. J. Epidemiol.*, **11**, 120.
- LUBIN, F., RUDER, A.M., WAX, Y. & MODAN, B. (1985). Overweight and changes in weight throughout adult life in breast cancer etiology. *Am. J. Epidemiol.*, **122**, 579.
- MCPHERSON, K. & DRIFE, J.O. (1986). The pill and breast cancer: Why the uncertainty? *Br. Med. J.*, **2**, 709.
- NISKIER, J.A., HAMMOND, G.L., DAVIDSON, B.J. & 4 others (1980). Serum sex-hormone-binding globulin capacity and the percentage of free estradiol in postmenopausal women with and without endometrial carcinoma. *Am. J. Obstet. Gynecol.*, **138**, 637.

A letter in similar vein to that of Key and Pike has also been received from Dr F. de Waard, Dept. of Epidemiology, Rijksinstituut voor Volksgezondheid en Milieuhygiene, The Netherlands.

- PIKE, M.C. & ROSS, R.K. (1984). Breast cancer. *Br. Med. Bull.*, **40**, 351.
- SHITERI, P.K. (1978). Steroid hormones and endometrial cancer. *Cancer Res.*, **38**, 4360.
- THOMAS, D.B. (1984). Do hormones cause breast cancer? *Cancer*, **53**, 595.

- WILLETT, W.C., BROWNE, M.L., BAIN, C. & 6 others (1985). Relative weight and risk of breast cancer among premenopausal women. *Am. J. Epidemiol.*, **122**, 731.

Dr Adami & colleagues reply:

Sir, – We appreciate the response to our somewhat provocative challenge of the 'general understanding' of breast and endometrial cancer and have read the letter of Drs Key and Pike with great interest. We should like to offer two comments. The first is a more general one, which concerns different approaches to dealing with a large amount of complex and contradictory data. The second, more specific comment refers to what we actually know so far about the pattern of risk factors for breast cancer in the population studied.

Risk factors for cancer of the endometrium and of the breast have been studied extensively during the last 15 to 20 years. Numerous associations have been revealed and those which have emerged with a reasonable degree of consistency are often referred to as 'well established'. Proceeding from these associations, some of which are mentioned by Key and Pike, it has been an attractive goal to find unifying concepts which link, for example, hormonal derivatives of risk factors to the probability of development of the disease. Most hypotheses related to such attempts have the postulate in common that excess oestrogenic stimulation – for example due to high circulating levels or increased bioavailability of oestrogens or insufficient opposing effects of progesterone – is an important determinant for neoplastic transformation, particularly if occurring during periods of assumed increased susceptibility of the target organ.

We agree with Key and Pike that the findings concerning relations to cancer of the endometrium have been fairly consistent. However, as far as breast cancer is concerned, the pattern is frustratingly complex and equivocal. A scrutiny of available data has shown that the number of studies in which the association between breast cancer and the 'established risk factors' (e.g. early menarche, low parity, late first birth, late menopause, short duration of breast feeding) has been found to be weak or absent is greater than can be explained by methodological flaws and type II errors. The resulting uncertainty might be an important reason why the same factors are actually being studied over and over again. The lack of association between age at first birth and risk of breast cancer, which we found at the beginning of this decade (Adami *et al.*, 1980) has now been confirmed in Denmark (Ewertz, 1987), Norway (Kvåle *et al.*, 1987) and Sweden (Adami, unpublished).

It is an attractive challenge to attempt to reduce this confusing pattern to a single unifying hypothesis. The risk involved in this approach is that when the same risk factors

are selected and approximately the same hypothesis formulated a number of times, a conformistic situation is created in which the large inconsistencies in available evidence become neglected. This will hamper a critical assessment of current research strategies and delay the advancement of more fruitful hypotheses.

More specifically, the 'general understanding' to which Key and Pike refer, requires that obesity be a risk factor for postmenopausal breast cancer – and preferably that it protect against premenopausal occurrence in the Swedish population. We agree that our previous entirely negative finding in a relatively small case control study needs to be confirmed (Adami *et al.*, 1977). Confirmatory results to be published have emerged, however, from two other sets of data. Thus, a recent case-control study in women younger than 45 years (Meirik *et al.*, 1986) failed to reveal any association between obesity and breast cancer (unpublished). Likewise, weight was analysed in a nested case-control study within a cohort of women who had received climacteric oestrogen treatment (Persson *et al.*, 1983; Bergkvist, 1987). Using weight below 60 kg as a reference, the relative risk in women who weighed 90 kg or more was 0.55 (95% confidence interval 0.15–2.00). When weight was included as a continuous variable in a logistic regression model, the relative risk per kg increase was 0.99 (0.98–1.02) (Bergkvist, unpublished). These findings are not unique for the Swedish population. The literature offers in fact only highly equivocal support for the finding that is crucial for the claim by Key and Pike, namely that obesity entails an increased risk of developing breast cancer. And in the positive reports, the possible confounding effect of dietary patterns needs to be clarified.

From these considerations, we still believe that critical judgement of available evidence justifies the conclusion that the common aetiological mechanisms of breast and endometrial cancer are poorly understood within the general framework ('the traditional paradigm') of associations with obesity, the reproductive characteristics and exogenous hormones.

Yours etc'.

H.-O. Adami¹, L. Bergkvist¹, I. Persson² & B. Pettersson³
Departments of ¹Surgery; ²Gynecology and Obstetrics
and Oncology and ³Division of Gynecologic Oncology
University Hospital
Uppsala

References

- ADAMI, H.-O., RIMSTEN, Å., STENKVIST, B. & VEGELIUS, J. (1977). Influence of height, weight and obesity on risk of breast cancer in an unselected Swedish population. *Br. J. Cancer*, **36**, 787.
- ADAMI, H.-O., HANSEN, I., JUNG, B. & RIMSTEN, A.J. (1980). Age at first birth, parity and risk of breast cancer in a Swedish population. *Br. J. Cancer*, **42**, 651.
- BERGKVIST, L. (1987). *Menopausal Estrogens and Breast Cancer*. (Thesis). Uppsala University, Uppsala, ISBN 91-554-2053-2.
- EWERTZ, M. (1987). Effect of pregnancies on breast cancer risk in Denmark. *EORTC Breast Cancer Working Conference*, abstract B 1.6.
- KVÅLE, G. & HEUCH, J. (1987). A prospective study of reproductive factors and breast cancer. II. Age at first and last birth. *Am. J. Epidemiol.* (in press).
- MEIRIK, O., LUND, E., ADAMI, H.-O., BERGSTROM, R., CHRISTOFFERSEN, T. & BERGSJO, P. (1986). Oral contraceptive use and breast cancer in young women. *Lancet*, **i**, 650.
- PERSSON, I., ADAMI, H.-O., JOHANSSON, E.D.B., LINDBERG, B., MANELL, P. & WESTERHOLM, B. (1983). A cohort study of oestrogen treatment and the risk of endometrial cancer: Evaluation of a method and its applicability. *Eur. J. Clin. Pharmacol.*, **25**, 625.