- HADDOW, A., HARRIS, R. J. C., KON, G. A. R., AND ROE, E. M. F.—(1948) *Phil. Trans.*, A, **241**, 147.
- Idem*, SCOTT, C. M., AND SCOTT, J. D.—(1937) *Proc. Roy. Soc.*, B, **122**, 477.
- HANCE, R. T., AND MURPHY, J. B.—(1926) *J. exp. Med.*, **44**, 339.
- HOROWITZ, N. H., HOULAHAN, M. B., HUNGATE, M. G., AND WRIGHT, B.—(1946) *Science*, **104**, 233.
- KOLLER, P. C.—(1934) *Genetics*, **16**, 447.
- LACASSAGNE, A., BUU-HOI, AND CAGNIANT, P.—(1944) *C.R. Soc. Biol. Paris*, **138**, 16.
- MATHER, K., AND STONE, L. M. A.—(1933) *J. Genet.*, **28**, 1.
- MULLER, H. J.—(1928) *Proc. nat. Acad. Sci. Wash.*, **14**, 714.
- NETTLESHIP, A., AND HENSHAW, P. S.—(1943) *J. nat. Cancer Inst.*, **4**, 309.
- PATERSON, E., HADDOW, A., APTHOMAS, I., AND WATKINSON, J. M.—(1946) *Lancet*, i, 677.
- PROSSER, C. L., PAINTER, E. E., LISCO, H., BRUES, A. M., JACOBSON, L. O., AND SWIFT, M. N.—(1947) *Radiology*, **49**, 299.
- STRONG, L. C.—(1945) *Proc. nat. Acad. Sci. Wash.*, **31**, 290.
- TATUM, E. L.—(1947) *Ann. N.Y. Acad. Sci.*, **49**, 97.

CARCINOGENIC ACTIVITY OF LIPOID SUBSTANCES.

I. HIEGER.

From the Chester Beatty Research Institute, Royal Cancer Hospital, London, S.W. 3.

Received for publication January 18, 1949.

THE investigation of carcinogenic substances which have been obtained from human tissues (Kleinenberg, Neufach and Shabad, 1940) and from other biological sources has been the subject of earlier publications from this Institute (Hieger, 1940, 1941, 1946, 1947), reporting that :

1. A carcinogen is present in extracts from human subjects who have died of cancer *or* of other causes.
2. The carcinogen is found in the unsaponifiable part of a number of tissues, such as liver, lung, kidney and muscle.
3. It is found in the cholesterol-rich part of the unsaponifiable fraction, which contains c. 85 per cent of the digitonin precipitable sterol and is referred to below as "Fraction A" or "Stage A," or "— 10° fraction," since it crystallizes out at — 10° when the unsaponifiable fraction of the tissue (doubly saponified) is crystallized from methyl alcohol.
4. Commercial cholesterol containing 95–98 per cent of the pure sterol is carcinogenic, and its potency is of the same order as that of the active fractions from human tissues.

With very few exceptions the carcinogen in biological material is slow acting (Table V), by comparison with synthetic carcinogens. Hence the tests on animals are lengthy.

Since some of the experiments described here were in progress when the papers referred to above were published and others have not before been described in detail, a certain amount of repetition is unavoidable.

1. *Technique used in these experiments for testing for carcinogenic activity.*

Unless otherwise stated, the substance is injected subcutem. (15 per cent solution in lard) into mice of C57 strain or of mixed commercial stock. The injection of 0.1–0.4 ml. is repeated fortnightly until a reservoir remains under the skin; injections are suspended as long as a nodule can be detected by touch. All tumours referred to are spindle-cell sarcomas arising at the site of injection (Fig. 1–8). Precautions were taken to prevent contamination by carcinogens from outside (Hieger, 1946).

2. *Control experiment on reagents used in the saponification of tissue by potassium hydroxide and alcohol.*

Since the great majority of the carcinogenic fractions had undergone saponification at least once, control experiments on the process itself acting on non-carcinogenic tissue components were made. The saponification conditions were described in detail (Hieger, 1946); about 1500 g. tissue is refluxed on the water bath with 400 g. KOH and 500 c.c. distilled alcohol for 2–3 hours and then extracted several times with ether. Carcinogens might be formed during saponification from precursors in the tissues or from the reagents themselves.

In the earlier experiments the alcohol was industrial spirit (the manufacturers state that it consists of ethyl and methyl alcohols in the ratio of 19:1 with 5 per cent water), which in separate experiments was found to darken rapidly on refluxing with KOH owing to resinous condensation of aldehyde. Acetaldehyde was therefore refluxed with KOH and spirit under the same conditions as those of tissue saponification. The resinous product was extracted from the saponification mixture with ether and injected into mice in the usual way. In another experiment the resinous product before injection into mice was mixed with cholesterol, which was intended to act as a carrier, in order to imitate rather more closely the products and conditions after tissue saponification (Experiments 1 and 2). All the tests on resin gave negative results.

Although in the earlier experiments spirit was used for the saponification, a number of unsaponifiable fractions made from adrenals, cancer liver and non-cancer liver (see note to experiment 3a for a definition of these terms) were found to be non-tumour-producing, which suggests that carcinogens or precursors are

DESCRIPTION OF PLATES.

FIG. 1.—Sarcoma in a C57 ♂ mouse at site of injection of commercial cholesterol in lard (15 per cent). 11th month. Experiment 10d. This tumour is now in its 63rd generation. $\times 55$.

FIG. 2.—Same tumour as Fig. 1. $\times 290$.

FIG. 3.—Sarcoma produced by commercial cholesterol in a C57 ♂ mouse. 20th month. Experiment 10d. $\times 55$.

FIG. 4.—Same tumour as Fig. 3. $\times 290$.

FIG. 5.—Sarcoma produced by commercial cholesterol in a MRC ♂ mouse. 23rd month. Experiment 10d. $\times 55$.

FIG. 6.—Sarcoma produced by cod liver oil in a C57 ♀ mouse. 17th month. Experiment 10b. $\times 55$.

FIG. 7.—Sarcoma produced by a fraction from lung-kidney-muscle of non-cancerous human subjects. The fraction had been saponified 3 times, crystallized from MeOH 3 times and once from acetone. 16th month. Stock ♂ mouse. Experiment 3f. $\times 55$.

FIG. 8.—Sarcoma produced by lard in an old C57 ♀ mouse which was 16 months at the beginning of the experiment. 6th month. $\times 55$.

either absent in the reagents or are present only occasionally. If the carcinogen were derived from the spirit a constant carcinogenic factor would be present. An alternative explanation would be that, in the experiments just mentioned, the mice happened to be non-sensitive. The question of sensitivity will be discussed more fully later on.

3. *Fractionation of cholesterol rich (c. 85 per cent) carcinogenic fraction from human tissues.*

Since the carcinogen was concentrated in the c. 85 per cent cholesterol fraction of the unsaponifiable part of tissue, a number of sub-fractions were prepared and tested on mice, thus:

(a) Eighty-five per cent cholesterol fraction (fraction A) from pooled cancer livers was recrystallized from hot MeOH and cooled to room temperature instead of to -10° ; crystals and mother liquors were tested separately on 20 and 15 C57 mice respectively. In the 20th month a single tumour appeared in the "crystals" series (Table I).

Note.—In this paper "cancer liver" indicates the liver of a human subject who has died of cancer of *any* part of the body. "Non-cancer liver" indicates the liver of a human subject who has died of any cause *other* than cancer.

(b) Fraction A from mixed cancer livers, partly the same as those used for the preparation in (a), was crystallized from 80 per cent aqueous alcohol after adsorption from petrol ether on to a large column of Al_2O_3 eluted with petrol ether-benzene-MeOH (80-20-20). The 80 per cent aqueous alcohol crystals (glistening plates m.p. $130-135^{\circ}$) were tested on 10 C57 mice; the mother liquors sufficed for 5 mice. In the 19th month a sarcoma appeared in the crystal series.

(c) A more detailed fractionation of fraction A from mixed cancer livers was carried out as in Flow Sheet I.

The fractions A, B, C, D, E gave no tumours, although of 40 mice initially, 24 survived 21 months; for survival rate see Table I. The possible reasons for the absence of tumours are discussed below.

(d) A similar fractionation was made on another batch of cancer livers, and mice were injected with material from G, C, F, D, E and H. No tumours arose, although of 50 mice, 48 survived for 12 months, and 32 for 18 months. The test on F was repeated on a further 10 mice in which one tumour arose after 2 years.

(f) The -10° fraction from human non-cancer lung-kidney-muscle (fraction A) had given 3 tumours in 25 mice, and was considered a potent carcinogen-containing fraction. It had been in the cold room for over 3 years and was then fractionated according to a simplified scheme (Flow Sheet 2).

Of 52 mice at the beginning of the tests, 43 survived 12 months and 26, 18 months. Four sarcomas appeared, 3 in fraction G and 1 in C.

(g) The crude unsaponifiable fraction (M) of mixed livers which had been used in experiment (d) was fractionated and tested at J, K, L, G, A, C, F, M, I, using in all 110 mice, which lived well; 92 survived 12 months and 40, 18 months. Two tumours were produced, one by the fraction

FLOW SHEET I.

FLOW SHEET 2.

obtained at the same level as in (d) (i.e. mother liquors of acetone crystals, (F)) and one by the crystals from which these mother liquors had separated (G).

(h) *Fractionation of 85 per cent cholesterol fraction via acetylation (in collaboration with Dr. C. W. Shoppee).*—In two earlier papers (Hieger, 1940, 1946) it was reported that the unsaponifiable fraction of the livers of Bantu natives contained carcinogen. A large batch of the crude unsaponifiable matter was worked up to stage A; a part was kept for injection, and another part was acetylated and crystallized by careful seeding. The crystalline acetate was hydrolysed and the regenerated cholesterol tested. 20 mice were employed in each test. In the 12th month one sarcoma appeared in the A series, and in the 19th month one sarcoma developed in the cholesterol group.

(i) *Cholesterol-rich fraction from cream (cow's milk).*—Kennaway (unpublished communication) had found that the unsaponifiable part of cow's milk produced a sarcoma by injection into C_3H mice. A more concentrated source of the unsaponifiable fraction was obviously milk fat. 3 kg. cream was therefore saponified twice, extracted and crystallized from MeOH to give stage A fraction which was injected into 20 mice. Stage L fraction, i.e. mother liquors of A, were also injected into 10 mice. Two tumours appeared in the crystalline A fraction series in the 13th and 19th months; no tumours were found at the site of injection in the L series.

(j) *Bile and gall bladder as sources of carcinogen.*—According to one hypothesis of human cancer, carcinogens are formed as a result of faulty steroid metabolism, consequently bile, which is a rich source of steroids, might well contain the aberrant factor.

Forty-five gall bladders from cancerous subjects were used; the bile was not separated, and some of the bladders were almost empty. Bladders and bile were saponified and worked up as before and the fractions tested at C, D, E, F. The experimental mice lived well; after 29 months a mouse of the F series (Flow Sheet I) developed a sarcoma at the site of injection.

(k) Liver from a lymphosarcomatous human subject gave a highly carcinogenic unsaponifiable fraction in some of the earlier experiments (Hieger, 1940), and the tissues of humans with this type of cancer may possibly be rich sources of carcinogen. Three livers from lymphosarcoma cases were saponified, fractionated into stages A, J, M, L, and injected into 6, 5, 11 and 6 mice of crossed C3H $\times$ C57 strain, which survived well, but no tumours developed.

Experiment 4.

The 85 per cent cholesterol fraction (A) of cancer livers which had given 2 sarcomas in 10 stock mice was stored in the cold room for $2\frac{1}{4}$ years, then tested on 10 C57 mice. No tumours arose, although 3 mice survived 24 months.

Experiment 5.

Dr. Mead, of British Drug Houses Ltd., kindly sent us for testing 5 fractions consisting chiefly of cholesterol obtained from cattle liver in the preparation of vitamin A concentrates. 119 mice were used in all for these tests but they did not live well; 60 survived for 12 months and 18 for 18 months; no tumours resulted.

Experiment 6.—To test if commercial cholesterol could replace the “cholesterol-rich (c. 85 per cent) fraction.”

Mixed human livers, some cancer, some non-cancer, were saponified and worked up to give the -10° (i.e. 85 per cent cholesterol) fraction (A) and also the mother liquors of this fraction (L). Commercial cholesterol was then added to L to give the same concentration of sterol as constituted the unsaponifiable fraction before the separation of the crystalline part A. Three series of mice were set up:

- (1) injected with -10° fraction (A);
- (2) „ „ mother liquors of -10° , i.e. L;
- (3) „ „ L to which the appropriate amount of cholesterol had been added.

Of 32 mice originally 28 survived 12 months, and 14, 18 months; no tumours resulted.

7. Effect of solvent.

Sesame oil is commonly used as a solvent for the injection of compounds undergoing tests for carcinogenic activity. Gardner (unpublished communication) reported that he found a single sarcoma in control mice injected with this solvent. Steiner, Warren Stanger and Bolyard (1947) obtained one sarcoma in

145 mice injected with tricaprylin, and Harris (1941) found that wheat germ oil gave rise to sarcoma on injection into rats and mice.

Two series of mice were set up ; one was injected with sesame oil containing 10 per cent commercial cholesterol, and one with the oil alone as control. One sarcoma appeared in the first series in the 14th month in a C57 female, and one sarcoma of fibrous type in the second in the 22nd month in a stock female (Experiment 7, Table I).

When over 300 control tests had been carried out on lard alone without the production of a single tumour, a sarcoma arose at the site of injection in a C57 mouse first injected with lard when already 16 months old. The latent period was 6 months, the shortest observed here in all the tests on carcinogens of biological origin.

Old mice were used in this experiment to see whether the prolonged latent period for tissue carcinogens was equivalent to the interval required for some essential preliminary change in the tissue which prepares it for carcinogenesis, and whether the natural ageing of tissue could in some degree be a substitute for the latent interval.

Earlier experiments in this Institute had shown that sarcoma may occur in rats injected with lard alone when strict precautions were taken against contamination by hydrocarbons (Burrows, Hieger and Kennaway, 1932, 1936).

8. *Fatty extract of Walker rat tumour.*

Aptekman, King and Lewis (1943) induced sarcoma in rats by injection of the fatty fraction of Walker rat tumours. Their results were confirmed here by tests where the tumours were dried by storing in distilled spirit and extracting with benzene. The crude benzene extract was evaporated, and the residue without added solvent (lard) injected subcutaneously into 10 wistar rats. A parallel group of 10 rats were injected subcutaneously with lard alone. After two years one rat in each series developed a fibro-sarcoma, which grew with difficulty on transplantation. The tumour in the lard series was more malignant than the tumour in the Walker extract series. Both tumours arose in the subcutaneous tissue at the same level as the injection site, but on the opposite side of the body. This effect has been observed several times in experiments where the injected substance is very fluid, and is probably pressed round the subcutaneous spaces by the movements of the animal.

9. *Miscellaneous tissues.*

The ether extract of minced mammae from lactating mice (DBA of high mammary cancer strain) proved carcinogenic, for it gave 1 sarcoma in 10 mice (Hieger, 1941). An ether extract of the lactating mamma of the rabbit was evaporated and the oily material injected into 20 mice, but no tumours resulted. Two sarcomas were obtained by injection of an ether extract of human cancer liver (Hieger, 1946).

10. *Carcinogenic activity of commercial cholesterol.*

The carcinogenic activity of commercial cholesterol (about 95-98 per cent pure sterol) has been briefly described (Hieger, 1947). Tests of the commercial product dissolved in lard were begun when it was found that the fractions of

TABLE I.—*Sarcoma Production by Injection of Lipoid Substances.*

Experiment.	Substance.	Mouse strain.	Mice at start.	Survivors (months).							Total matings.
				12.	15.	18.	21.	24.	30.		
3a	Livers from cancerous subjects. Fraction A: crystallized } from MeOH } → Crystals → Mother liquors	C57	20 15	14 14	13 11	10 10	*9 8	5 0	1 0	1 0	
3b	Livers from cancerous subjects. Fraction A: passed through Al ₂ O ₃ column, } crystallized from 80% } alcohol } → Crystals → Mother liquors	C57	10 5	8 4	8 4	8 4	*6 4	5 4	0 0	1 0	
3c	Livers from cancerous subjects. Fraction A " B " C " D " I " (See Flow Sheet 1)	C57	9 11 10 5 5	8 10 10 5 4	6 10 10 5 4	5 10 10 4 3	2 10 5 4 3	1 9 0 4 0	0 0 0 0 ..	0 0 0 0 0	
3d	Livers from cancerous subjects. Fraction G " C " F " E " D " I " (See Flow Sheet)	C57	10 10 20 10 5 5	9 10 17 10 5 5	9 10 16 10 5 5	8 10 8 8 5 0	7 8 7* 5 0 0	1 6 3 4	0 0 0 0	0 0 1 0 0 0	
3f	Lung-kidney-muscle from non-cancerous subjects. Fraction A " G " D " C " F	Stock	10 10 10 12 10	9 9 9 8 8	4 8 7 7* 8	3 5 6 5 7	2 5 2 3 3	0 *0* 0 0 0		0 3 0 1 0	
3g	Livers from cancerous and non-cancerous subjects. Fraction M " A " K " J " G " L " D " F " C " (See Flow Sheet)	Stock	15 15 10 15 10 15 10 5 15	10 12 8 *8 12 12 10 4 14	9 9 7 5 7 10 9 3 10	4 5 3 5 5 5 2 5	1 0 0 1 0 2 0 2	1 *0 1	0 0	 1	
3h	Livers from Bantu natives. Fraction A acetylated, cholesterol acetate crystallized out and cholesterol regenerated.	C57	20	8*	5	2	2	..	..	1	
3i	Cream (cow's milk). Fraction A " L	C57	20 10	13* 9	11 9	11* 4	10 0	4 0	0 ..	2 0	

TABLE I—(cont.).

Experiment.	Substance.	Mouse strain.	Mice at start.	Survivors (months).							Total sarcomas.
				12.	15.	18.	21.	24.	30.		
3j	Bile and gall bladders.	Fraction L " K " C " F	C57 " " "	10 10 10 10	8 8 10 8	4 8 8 7	3 6 3 6	3 5 2 5	0 0 2 3	0 0 0 1	
3k	Livers from lymphosarcoma cases.	Fraction M " J " L " A	Hybrid " " "	6 6 6 11	5 5 3 9			4 3 2 7	3 2 1 2	0 0 0 0	
6	Mixed livers (cancer + non-cancer). Substitution of A (i.e. 86% cholesterol fraction) by commercial cholesterol (C.C.).	Fraction A " L L + C.C.	C57 " "	10 10 12	10 9 9	9 8 8	3 5 4	2 0 [*] 0	0 .. 0	0 0 0	
7	Commercial cholesterol (C.C.) + Sesame oil (S.O.).	C.C. S.O. 10% 90% 0 100%	C57 Stock C57 Stock	10 10 10 10	16 15 13 12	2 2 2 2	1 1 [*] — —	— — — —		1 1 1 1	
10a	Commercial cholesterol re-fluxed with KOH and (a) alcohol, or (b) spirit.	a b	C57 Stock C57 Stock	10 10 10 10	9 7 10 9	3 0 10 0	2 .. 10 7	1 1 2 2	0 .. 0 0	0 0 0 2†††	
10b	Commercial cholesterol (C.C.) + Cod liver oil (C.L.O.), or linseed oil (L.O.).	Final mixtures made up as 15% solutions in lard.									
		C.C. C.L.O. L.O.									
	I 1 g. 12 "drops"	0	C57	10	10	10	9	9	5*	3	
	II 1 g. 2 "	0	"	10	10	10	9	8	8	0†	
	III 1 g. 0 12 drops		"	10	10	10	10	8	4	0	
	IV 1 g. 0 2 "		"	10	10	9	7	6	3	0	
	V 0 1.0 c.o.	1.0 c.o.	"	10	9	4	2	1	..	0	
	VI 0.5 g. 1.0 "	1.0 "	"	10	10	10	8	6	6	0	
	VII 6 g. 0	0	C57 Stock	15 15	20 13	14 10*	14 10	.. 4		1 3 (Stock ♂)	
	VIII 6 g. 1.2 c.o.	0	C57 Stock	15 15	13 8	10* 8	10 7			3 (Stock ♂, ♂, ♂)	
	IX 1.2 g. 6 "	0	C57 Stock	15 15	8 12	8 12*	7 9			1††† 1† (C57 ♂)	
	X 0 6 "	0	C57 Stock	15 15	12 12	12* 9	9 ..			1† 3 (C57 ♀)	

TABLE I—(cont.).

Experiment.	Substance.	Mouse strain.	Mice at start.	Survivors (months).							Total sarcomas.		
				12.	15.	18.	21.	24.	30.				
10c	Commercial cholesterol crystallized from acetone	Least soluble fraction {	C57	10	13*	11	**11	..	..	..	} 3 (Stock ♂, ♀ and C57 ♀)		
			Stock	10	..	..	..	..	..	..			
		Mod. soluble	C57	11	15*	12	9	..	..	..	} (Stock ♂)		
			Stock	10	..	..	..	..	..	..		..	
	Mother liquors	C57	12	13	12	10	..	..	..	..	..		
		Stock	10	..	..	..	..	..	..	..	..		
10d	Commercial cholesterol§	C57	5	..	*4	1	0	..	..	..	..	1	
		"	5	..	0	..	..	..	..	..	..	0	
		"	6	..	3	3	3	3	3*	0	..	..	1
		"	6	..	4	4	0	..	..	..	..	..	0
		"	6	..	0	..	..	..	..	..	..	..	0
		"	6	..	0	..	..	..	..	..	..	..	0
		CBA	5	..	3	2	0	..	..	..	..	..	0
		"	5	..	5	2	1	0	..	..	..	..	0
		"	5	..	5*	4	4	1	0	..	..	..	1
		"	5	..	5	5	2*	1*	..	..	..	..	2
		"	5	..	5	2	1	0	..	..	..	..	0
		"	5	..	2	0	..	..	..	..	..	..	0
		"	6	..	3	3	3	0	..	..	..	..	0
		"	5	..	4	3	2	2*	0	..	..	..	1
		"	5	..	5	4	3	2*	2	0	..	..	1
		"	5	..	4	0	..	..	..	..	..	..	..
		"	5	..	4	4	..	..	..	..	..	..	..
		"	5	..	0	..	..	..	..	..	..	..	..
		"	5	..	3	0	..	..	..	..	..	..	..
		"	5	..	4	3	..	..	..	..	..	..	..
"	4	..	4	3*	2*	..	..	..	..	..	2		
"	"	5	..	4	3*	..	..	..	..	..	1		
"	"	5	..	4	3*	..	..	..	..	..	..		

* Each asterisk represents one sarcoma. The position of the asterisk on the time-grid indicates the latent period.

† Indicates histological changes at the site of injection suggestive of incipient sarcoma production. Thus in bottom line of Experiment 10a (10 Stock mice) sarcomas arose in 2 mice and suggestive changes were observed in 4 others.

§ For a more complete statement of the results with cholesterol see Table II.

tissues which were richest in cholesterol were also the most potent carcinogenically. Up to the present 25 sarcomas have been obtained in 436 mice by commercial cholesterol or some fraction of it, in solution in lard or in other solvents (Table I (10d), and Table II).

TABLE II.—*Sarcoma Production by Commercial Cholesterol.*

	Material *	Number of mice.		Sarcomas.
		Initial	12 months.	
1. Commercial cholesterol	.	213	77	11
2. " " in sesame oil	.	20	16	1
3. " " + cod liver oil	.	100	71	7
4. " " (after refluxing with KOH in spirit (or alcohol))	.	40	35	2
5. " " crystallized from acetone	.	63	41	4
Total	.	436	240	25

* Injections of 15 per cent solutions in lard except for No. 2.

10a. *Effect on cholesterol of refluxing with KOH and alcohol.*

The possibility that carcinogens are formed from cholesterol during saponification was tested as follows: 10 g. of commercial cholesterol (i.e. about the amount present in a human liver) was refluxed for 2½ hours with 400 g. KOH, 1 l. distilled alcohol or spirit and enough water to imitate the conditions in the saponification of a liver; the mixture was extracted with ether and injected into mice as before. A preliminary statement (Hieger, 1947) was made, but further series of mice have been used (10a in Table I).

Four groups of mice were set up:

- a. 10 C57 mice injected with cholesterol saponified in distilled spirit.
- b. 10 stock " " " " "
- c. 10 C57 " " " " alcohol.
- d. 10 stock " " " " "

In the 14th month in group *b* 2 mice developed sarcomas, and pre-neoplastic changes were seen at the site of injection in 4 other mice.

Thus in group *b*, 6 of 10 mice had either developed sarcoma or might have done so if they had lived longer. The two sarcomatous mice and two of the incipiently sarcomatous mice were from the same box of 5 mice.

One cannot say why only group *b* responded so readily, but (1) in a fair number of experiments the saponification of tissue has been done in spirit but the products have produced no tumours, showing that spirit is not essential for carcinogenesis by the unsaponifiable product; (2) the absorption spectra of the two final products (cholesterol saponified in spirit and in alcohol) are closely similar; and (3) while the substance was strongly carcinogenic in group *b* (stock mice), the same preparation was inactive in group *a* (C57). Consequently it must be supposed that the *b* group of mice were either—

(1) Peculiar genetically; (2) peculiar congenitally; (3) had become infected or (4) had become contaminated with carcinogen.

(4) is, on the whole, not very probable, because:

The 1946 paper reported that in tests on 18 subfractions of the unsaponifiable fraction of human tissue, partly from cancerous, partly from non-cancerous

subjects, of the total yield of 8 sarcomas, 7 were produced by fraction A made from 3 different sources and 1 from an M fraction (Flow Sheet 1). The concentration of 7/8ths of the sarcomas in one particular fraction strongly suggests that the fraction itself was responsible and not a chance contamination, for it would be highly improbable that contamination would occur only in the boxes, holding 5 mice each, containing those treated with the A fraction.

The data in Table III are the results when a fraction was divided into 3 or more sub-fractions. The concentration of tumour production in a few fractions, particularly F, G and A, supports the conclusion in the foregoing paragraph, namely, that contamination will not explain satisfactorily the incidence of sarcomas. Contamination only in those boxes where these particular fractions were used would be very unlikely, since each box contains 5 mice initially and the experiments on these fractions were started over a period of 4 years, during which time the boxes were changed many times (Table III).

TABLE III.—*Showing Trend of Concentration of Carcinogenic Factor During Fractionation.*

Fraction (see Flow Sheet 1).	Number of samples derived from different batches of tissue.	Number of mice at start.	Survivors at 12 months.	Number of sarcomas.		Number of samples which gave positive results.
					%.	
A .	11 .	175 .	110 .	10	6	5
M .	4 .	52 .	41 .	1	2	1
J .	4 .	40 .	25 .	0	0	0
L .	8 .	141 .	101 .	0	0	0
G .	3 .	30 .	26 .	4	13	2
C .	5 .	57 .	52 .	1	2	1
F .	4 .	45 .	37 .	3	7	3
I .	2 .	10 .	9 .	0	0	0
D .	3 .	25 .	24 .	0	0	0
E .	1 .	10 .	10 .	0	0	0
K .	2 .	20 .	18 .	0	0	0

10b. *Effect of addition of oils on carcinogenic potency of commercial cholesterol.*

The activity of commercial cholesterol suggested several possibilities; for example :

- (a) That cholesterol is indeed a feeble carcinogen.
- (b) That commercial cholesterol as well as the cholesterol-rich fraction obtained from a number of sources contains a small amount of a potent carcinogen.
- (c) That a combination of cholesterol, as carrier, and an active agent are essential for carcinogenesis.
- (d) That all the preparations of cholesterol used are contaminated with atmospheric carcinogen (i.e. benzpyrene from soot).
- (e) That carcinogens of biological origin such as lipids and particularly preparations rich in cholesterol evoke neoplasia, but only in the most susceptible members of groups of experimental animals.

Tests of hypotheses (a) and (b) are to be described more fully in a forthcoming paper. Highly purified cholesterol, prepared by Dr. C. W. Shoppee, using bromination, crystallization of acetate and chromatography, has been

injected into 25 mice; the experiment had been in progress for 22 months without result as yet. (b) is being tested by concentrating the impurities in commercial cholesterol 3 times and in a second experiment 10 times, and injecting the products. These experiments have been in progress for 11 months without result as yet. (c) was investigated by mixing oils containing reactive double bonds (i.e. unsaturated) with commercial cholesterol to see if the activity could thereby be increased (Table I, experiment 10b).

Cod liver oil certainly increases the potency of cholesterol dissolved in lard, but since at least one sarcoma has now been obtained with cod liver oil alone (Table I, 10b, series x), it is uncertain whether the oil is a co-carcinogen or a carcinogen.

10c. *Effect of simple crystallization on the carcinogenic potency of commercial cholesterol.*

The batch of cholesterol used in the preceding experiment was crystallized from acetone to give 3 fractions:

- (1) Least soluble.
- (2) Moderately soluble.
- (3) Mother liquors, which is obviously different from (1) and (2), for (3) is softer, yellower, more soluble in lard and smells of candle wax.

The three fractions were injected as usual into 20, 21 and 22 mice respectively; 3 sarcomas appeared in the fraction 1 (least soluble) series and 1 in fraction 2 series in the 14th, 18th, 18th and 15th months.

10d. *Sarcoma production by commercial cholesterol.*

Table II shows the total of sarcomas produced by commercial cholesterol.

10e. *Compounds related to cholesterol.*

Compounds related to cholesterol and particularly those which might be present in samples of the impure sterol were also tested, namely, isocholesterol, isolumisterol, cholesterylene (2 different preparations and also the mixture of compounds present in the mother liquors after crystallizing out the cholesterylene), ergosterol, cholestene, 7-dehydrocholesterol (2 different preparations, one a commercial product containing 60 per cent of the pure dehydrocholesterol), a mixture of epi- and allo-cholesterol, vitamin D₃, and 7-ketocholesterol. Except for those treated with 7-ketocholesterol all the mice have now died without tumours being produced; 90 survived 12 months out of 100 initially, and 56 survived 18 months. The test on 7-ketocholesterol has now been in progress for 17 months, without result.

11. *Pre-neoplastic changes at the site of injection of lipid.*

Since the detection of only a few tumours could be important in deciding the technique of further fractionation, it was essential not to miss any tumours not visible macroscopically. The remaining mice (now 700) are examined post mortem by sections through the fat vesicle, so that any pre-neoplastic changes might be detected. In 15 of the 700, changes were seen suggestive of pre-sarcomatous change at the site of injection.

12. *Steiner's investigations.*

In a recent elaborate report Steiner, Warren Stanger and Bolyard (1947) give further data on sarcoma production by unsaponifiable fractions in three strains of mice, Albino, A, and C57 (Table IV, A):

TABLE IV, A.—*Experiments of Steiner, Warren Stanger and Bolyard (1947):—Sarcoma Production by Unsaponifiable Fraction of Tissues.*

Experi- ment.	Unsaponifiable fraction of—			Number of cases (pooled).	Mice at start.	Mice at 12 months.	Sarcomas.
Cancer cases :							
1 .	Human kidney	.	.	61 .	50 .	39 .	0
2 .	„ spleen	.	.	54 .	50 .	47 .	5
3 .	„ colon	.	.	54 .	49 .	45 .	0
4 .	„ heart	.	.	63 .	43 .	19 .	1
5 .	„ liver	.	.	18 .	101 .	45 .	17
Non-Cancer cases :							
6 .	Human kidney	.	.	58 .	50 .	44 .	1
7 .	„ spleen	.	.	51 .	50 .	40 .	0
8 .	„ colon	.	.	51 .	47 .	45 .	0
9 .	„ heart	.	.	35 .	31 .	7 .	0
10 .	„ liver	.	.	15 .	120 .	39 .	44
11 .	Human liver (stillborn infants)	.	.	40 .	39 .	34 .	3
12 .	Pig liver	.	.	— .	115 .	82 .	4
13 .	Beef liver	.	.	— .	115 .	69 .	0
14 .	Pig heart	.	.	— .	30 .	9 .	0
15 .	Tricaprylin controls	.	.	— .	154 .	116 .	1
Total					1044	680	76

The yield of local sarcomas at the site of injection (7.3 per cent) is of the same order as that obtained in this Institute, namely 4.7 per cent (63 sarcomas in 1298 mice, of which 936 lived 12 months). These 1298 mice are all those injected with unsaponifiable material of animal origin, including commercial cholesterol.

In earlier papers, Steiner (1942, 1943) gives results which, taken in conjunction with the data from his recent paper, can be expressed thus (Table IV, B):

TABLE IV, B.—*Experiments of Steiner, Warren Stanger and Bolyard (1947):—Sarcoma Production: Effect of Pooling of Unsaponifiable Fractions.*

Material.			Ratio: Sarcoma mice	total mice used.
1.	8 cancer livers pooled	.	12/56 = 21%	} mean 21%
2.	7 non-cancer livers pooled	.	5/63 = 8%	
3.	15 " " " "	.	44/120 = 37%	
4.	18 cancer livers pooled	.	17/101 = 17%	
5.	37 " " tested separately (8 in 37 were active)	.	12/456 = 2.4%	}
6.	30 non-cancer livers tested separately (6 in 30 were active)	.	10/440 = 2.3%	

Thus the effect of testing a number of livers (67) separately was to reduce the sarcoma production by a factor of 9, i.e. 9 times less than the potency of 48 livers pooled, which is of the order of expectation if it be noted that Steiner (1942) found 1 in 5 of his livers (cancerous and non-cancerous) to contain the active factor, and if it be assumed that pooling the livers had the effect of distributing the agent from the active livers among the inactive ones. Furthermore, it may be noted that from Steiner's (1942) experiments 1 and 2 one would conclude that

cancer livers are of the order of $2\frac{1}{2}$ times as potent as non-cancer livers, but from experiments 3 and 4 it would appear that the ratio of potencies is reversed, leading to an overall discrepancy as high as 5 to 1. The writer would point out that three experiments (Nos. 10a and 3f, Table I and Experiment 4) carried out here suggest that unexplained differences in the susceptibility of the mice to biological carcinogens can completely alter conclusions as to the carcinogenicity or otherwise of a substance undergoing test.

TABLE V.—*Latent Period* (Months) of Sarcomas Produced in Mice by Injections of Lipoid Substances.*

Latent period (months).	Hieger (1940). Number of tumours.	Hieger (1946). Number of tumours.	Recorded in the present paper. Number of tumours.	Total.
6	—	—	1	1
7	—	—	—	—
8	—	—	1	1
9	—	—	—	—
10	1	—	—	1
11	2	—	1	3
12	2	—	2	4
13	1	—	2	3
14	—	1	6	7
15	1	2	1	4
16	—	1	4	5
17	1	—	3	4
18	2	—	2	4
19	2	1	3	6
20	1	1	3	5
21	2	—	3	5
22	1	—	4	5
23	2	1	3	6
24	—	1	2	3
25	—	—	1	1
26	—	—	—	—
27	—	—	—	—
28	—	—	—	—
29	—	—	1	1
Total				69

* "Latent period" is taken as the interval between the start of the experiment and the grafting of the induced tumour. The actual initiation time for the tumour is, of course, shorter. The interval between detection and grafting is seldom more than a few weeks. Once the primary tumour is well started it grows with great speed.

Note.—The figures in the last column (the overall latent period) show a periodicity remarkable in such data. It is possible that at the large number of mice involved, nearly 2000, statistical laws begin to operate.

DISCUSSION AND CONCLUSIONS.

The experimental data given here and in earlier papers from this Institute, based on 69 sarcomas produced in a total of 1966 animals, present a number of difficult problems.

No completely satisfactory explanation of the following experimental findings is as yet available :

(1) Cod liver oil, lard and sesame oil produced at least 1 sarcoma each in 25, 350 and 20 mice respectively. Other work has shown that some fats can bring about sarcoma development in sensitive mice or rats.

(2) Commercial cholesterol in lard gave sarcomas in about 5 per cent of 213 mice (Table II).

(3) The mixture formed when cholesterol is refluxed with KOH and distilled spirit gave no tumours when injected into 10 C57 mice, but in 10 stock mice there arose 2 sarcomas, and in 4 other mice of the same series changes were detected, post-mortem, suggestive of pre-neoplasia, at the site of injection. A similar preparation where cholesterol was refluxed with KOH and alcohol gave no tumours when injected into 10 stock and into 10 C57 mice.

(4) Fractionation of an active fraction from tissues by simple crystallization does not always lead to any concentration of the active factor.

(5) Although several experiments showed that the addition of cod liver oil to commercial cholesterol increased its potency, yet in another experiment the crystallization of the commercial sterol from acetone, which removed some waxy constituents, *also* increased the potency.

The writer has attempted to reconcile these apparently contradictory findings by putting forward the following suggestions: Firstly, that the operation of little understood factors covered by the expression "individual variations of susceptibility of the experimental animals" is very important in determining the carcinogenic potency of the lipids, and secondly, that lipids and particularly sterols or closely associated compounds are slow-acting carcinogens. It is clear that large series of mice are required in this type of experiment where the agent is of low potency, and where, therefore, the presence or absence of especially sensitive individuals may well be the principal factor in the assessment of the potency of a test preparation, since in small series of mice the sensitive members may die first, leaving those which are comparatively insensitive.

The possibility is not quite excluded that lipids are often contaminated by traces of benzpyrene from air-borne soot.

Thirdly, the existence of about 300 synthetic carcinogens (hydrocarbons, azo compounds, stilbenes, etc.) suggests that carcinogenic potentiality is probably more widely distributed among chemical compounds than was at first anticipated, and thus there is no good reason for excluding compounds of the lipid group.

Fourthly, some of the experiments described here suggest that the carcinogenicity of lipid substances may either be enhanced or be inhibited by other lipids.

Fifthly, should the thesis proposed here, namely, that lipids are slow-acting carcinogens, be conclusively proved, then, "spontaneous" cancer or at least some forms of the disease could be considered as "caused" by body lipids existing locally in some peculiar chemical or spatial or temporal conditions.

The production of sarcoma by commercial cholesterol obviously casts some doubt on the conclusions which have been drawn from all previous experimental work where cholesterol was involved, and where almost all investigations have taken for granted that cholesterol is non-carcinogenic. The weakly carcinogenic potency of a number of other lipid substances and the unexplained variations of sensitivity of the animals used for testing underline the need for much further study in this field. The experiments described here suggest the question, Why has it required 20 years, that is, since the discovery of the carcinogenic hydrocarbons in 1929, to show that commercial cholesterol is carcinogenic? The answer is obvious; investigators in chemical carcinogenesis have become accus-

tomed to the use of powerful, quick-acting carcinogens, and as a result tumour induction by slow-acting carcinogenic substances has been dismissed as "non-specific."

Control experiments have, in the past, not been done on an adequate scale, and moreover, commercial cholesterol, having been unconsciously equated with the cholesterol in the body, leads to the inference that cholesterol is not a carcinogen since it is present universally. It might be permitted to note, however, that inferences cannot always satisfactorily substitute experiments.

SUMMARY.

1. A total of 69 sarcomas has been obtained at the site of injection of lipid substances in approximately 2000 mice of C57 and MRC strains and commercial mixed stock mice. The average latent period was about 18 months.

2. 63 tumours were produced by unsaponifiable fractions derived from the tissues of human subjects (cancerous and non-cancerous) and from cattle; of the remaining 6 tumours, 5 were produced by non-saponified fat from animal sources and 1 from a plant source.

3. Commercial cholesterol gave 25 sarcomas in 436 mice.

4. No evidence was obtained to suggest that the carcinogenic substances in the unsaponifiable fraction is confined to the tissues of human subjects who have died of cancer.

5. The carcinogenic factor has not yet been isolated in a chemically pure state. Fractionation by simple crystallization led in one experiment to an inexplicable loss of potency.

6. The widespread occurrence of a carcinogenic factor in unsaponifiable fractions from many biological sources suggests that it is either cholesterol itself, or a combination of cholesterol and a small proportion of a frequently occurring co-carcinogen.

7. In such experiments the possibility of contamination with benzpyrene from air-borne soot must be borne in mind.

8. The variations of susceptibility in the experimental mice constitutes a most important factor in determining the potency of carcinogens of biological origin.

This investigation has been supported by grants to The Royal Cancer Hospital, from the British Empire Cancer Campaign, the Jane Coffin Childs Memorial Fund for Medical Research, the Anna Fuller Fund and the U.S. Public Health Service.

REFERENCES.

- APTEKMAN, P. M., KING, H. D., AND LEWIS, M. R.—(1943) *Cancer Res.*, **3**, 856.
BURROWS, H., HIEGER, I., AND KENNAWAY, E. L.—(1932) *Amer. J. Cancer*, **16**, 57.—(1936) *J. Path. Bact.*, **419**, 43.
HARRIS, P. N.—(1941) *Cancer Res.*, **1**, 751.
HIEGER, I.—(1940) *Amer. J. Cancer*, **39**, 496.—(1941) *Science*, **93**, 262.—(1946) *Cancer Res.*, **6**, 657.—(1947) *Nature*, **160**, 270.
KLEINENBERG, A. E., NEUFACH, S. A., AND SHABAD, L. M.—(1940) *Amer. J. Cancer*, **39**, 463.
STEINER, P. E.—(1942) *Cancer Res.*, **2**, 425.—(1943) *Ibid*, **3**, 385.
Idem, WARREN STANGER, D., AND BOLYARD, H. N.—(1947) *Ibid*, **7**, 273.