HEAD INJURIES IN CHILDHOOD*

BY

PHILLIP HARRIS

From the Department of Surgical Neurology, Royal Infirmary, Edinburgh

In most respects the mechanisms and effects of head injury in children are the same as in adults, but there are several differences, and these are discussed with particular reference to the 150 children who were seen in the Department of Surgical Neurology, Edinburgh, during the five-year period 1950-54 (Table 1). During this time 450 adults with head

TABLE 1
TOTAL NUMBER OF PATIENTS, 1950-54

Age in Years	Males	Females	Total
0-1	10	6	16
2-5	20	14	34
6-12	50	11	61
13-16	31	8	39
	111	39	150

injuries were also treated. It is realized that these are selected groups of patients with head injury.

Sixty-one per cent. of the children had mild head injuries, requiring only a brief stay in hospital, they usually had no residual neurological abnormalities and did not require special surgical treatment. The remainder had severe injuries. In the adult series 32% had mild and 68% severe head injuries.

Pathology

In the adult a dynamic head injury is more liable to produce severe displacements of the brain and its blood vessels than is the case in children, where a higher proportion of depressed fractures, and on the whole a lower morbidity and mortality are found (Table 2, Fig. 1). Greater re-organization of nervous function appears to be possible in the child, but babies and young infants have delicate skulls, nervous tissue, and blood vessels, and do not withstand head trauma well. In Table 3 the type of injury is analysed.

Basal fractures of the skull were uncommon in children, but linear, diastatic and in particular compound fractures were frequently seen (Table 2).

* Read at the annual meeting of the British Association of Paediatric Surgeons, Edinburgh, June 6, 1957.

TABLE 2
TYPES OF SKULL FRACTURE

Linear	26
Depressed, simple	9
Depressed, compound	52
(1) Involving frontal air sinuses:	
(a) Minimal cerebral damage	10
(b) Severe cerebral damage	2
(2) Other regions:	
(a) Minimal cerebral damage	30
(b) Severe cerebral damage	10
Total	87

FIG. 1.—Radiograph showing depressed fracture in left parietal region.

TABLE 3
TYPE AND SEVERITY OF INJURY

	Mild	Severe	Total
Acceleration (including penetrating injuries)	44	25	69
Deceleration	40	23	63
Crush	7	4	11
	91	52	143

Although accurate comparisons between the childhood and adult series is not possible, 35% of the former category had depressed compound fractures, which were present in only 12% of the adults.

Possibly because of the late formation of the frontal air sinuses (6th-7th year) in children, they were involved less frequently than in adults.

Ingraham and Matson (1954) analysed the records of 1,330 children with head injuries seen during a 13-year period. Twenty-eight per cent. had cerebral concussion, 11% had depressed and 4.5% compound skull fractures. Two per cent. had an extradural haematoma, and 24% subdural haematomas. Regarding the latter, they state that the frequency with which it is found is proportional to the intensity with which it is sought. Although half of their patients with an extradural haematoma were under 2 years of age, this has not been our experience. The adherence of the dura to the skull and its fusion with the fontanelle may partly account for this (Voris, 1950). Seven per cent. of our children had extradural haematomas, 4% subdural haematomas and 6% intracerebral haematomas; the corresponding percentages in the adult series were 4, 13 and 2.

Even a mild head injury in a child who happens to have hydrocephalus can have serious effects, thus a patient with 'arrested' hydrocephalus may show evidence of rapidly increasing intracranial pressure.

Cerebral thrombophlebitis and venous sinus thrombosis account for some of the neurological signs seen in patients with head injuries and appear to be more significant in children than in adults.

Clinical Aspects

Clinical neurological evaluation is often difficult in the young, especially in the first three months of life, and a reliable history and careful, repeated clinical observation are important.

Eleven patients had an extradural haematoma, the ages varied from 4 to 11 years, and on the whole the history was briefer than was obtained in adults and there was often no initial period of unconsciousness. Epilepsy was not a feature. Voris (1950) had no patient below the age of 10 years with an extradural haematoma.

A characteristic syndrome was not found in infants with chronic subdural haematomas, but irritability, restlessness, epilepsy, vomiting, fever, a bulging fontanelle, retinal haemorrhages and drowsiness were sometimes present. The ability of the infant's head to enlarge to accommodate the haematoma probably accounts for the differences in the clinical picture in the child. Bilateral subdural haematomas are more commonly found in children than in adults.

Birth Injuries

In addition to the 150 children under discussion, 64 patients with cerebral birth injuries were treated in the same five-year period. The true incidence

and significance of cerebral birth injuries is unknown, but they appear to be common (Anderson, 1952). Of the 64 patients, 33 had epilepsy, 16 amentia and hemiplegia, 11 hydrocephalus and four had chronic subdural haematomas. Much is to be learnt about these sequelae of birth injury, and it is suggested that closer cooperation amongst the various specialists involved is necessary.

Special Methods of Investigation

Skull radiographs are important to demonstrate the size and location of the fragments of a depressed fracture (Fig. 1), and the presence of radio-opaque foreign bodies, otherwise they are of limited value. Three of the 11 patients with an extradural haematoma had no fracture, five had linear and three compound depressed fractures.

Occasionally a slowly progressive intracranial expanding lesion in a child may cause local enlargement and thinning of part of the skull. This was seen in the left temporal bone and middle cranial fossa of a 9-year-old boy, who had a chronic subdural haematoma (Fig. 2).

FIG. 2.—Radiograph showing enlargement of left middle cranial fossa with thinning of the skull, due to a chronic subdural haematoma.

Lumbar puncture is of very limited value in diagnosis and in fact may be dangerous in the unconscious or in the restless child. Thus, tentorial impaction may be induced if there is an intracranial haematoma.

In the early stages of a head injury in a child, electro-encephalography has been unhelpful in our experience. Even in the later stages, it may be difficult to interpret the findings in an infant.

Associated Injuries

These usually consisted of fracture of a limb bone (femur in seven cases, tibia in three, humerus in

three, ulna in one, clavicle in four. Two patients had maxillo-facial injuries necessitating special treatment. One patient had a vascular and peripheral nerve injury.

Infection

In one patient a scalp laceration became infected, and in another a brain abscess developed as the result of the penetration of the right frontal lobe by a garden fork.

Sequelae

Twenty-six per cent. of the patients had post-traumatic epilepsy (Table 4). In half there were clinical motor or sensory components. Only five of the children with epilepsy had a compound fracture of the skull with direct cerebral contusion and laceration. Twelve had had no initial period of unconsciousness associated with the head injury. As Livingston (1954) observes, children sustain very many bumps and bruises, and the significance of a head injury in a child in causing epilepsy can be difficult to decide. But on the other hand, 'cryptogenic' epilepsy has a definite stigma, and parents may thus prefer to over-stress the history of a previous head injury. Of 82 children with head injuries studied by Rowbotham, Maciver, Dickson and Bousfield (1954), only four had epilepsy.

Thirteen per cent. of our series of patients had psychological changes (Table 4) and in half of these

TABLE 4
SEQUELAE

Epilepsy	39
Persistent psychological changes	20
Dysphasia	12
Other neurological abnormalities	27

they persisted. When there has been profound depression of all higher functions from a severe brain injury the early phase of recovery of normal alertness is probably shorter in children than in adults. The next phase, in which there is normal alertness but lack of normal self-control, is commonly more striking in children, but again is usually of shorter duration and less frequently requires prolonged psychiatric care. The late phase, in which there may be full neurological recovery but in which residual unfavourable psychological reactions are still present, is relatively infrequent in children. Compared with adults (25%), the typical 'post-concussional syndrome' was rare in children (6%). Six of the latter were over 15 years of age. Adult attitudes appeared to be important. However, Rowbotham *et al.* (1954) found that 33% of children who had head injuries developed changes in temperament, and eight showed an intellectual injury of some degree. Bakwin (1949) also discusses behaviour disorders in children as a result of cerebral damage.

Recovery from brain injury is probably more rapid and complete the younger the individual, but adequate assessment is always necessary since children are relatively uncomplaining patients, and quite serious disabilities may go unnoticed by ordinary methods of investigation. On leaving hospital they may then either be kept off all schooling for unjustifiably long periods, or else be subjected to educational pressures which are beyond their capacity. This is perhaps particularly the case with language disabilities, in which difficulty in learning to spell and read may persist long after oral speech appears to have recovered. Oral speech indeed may not have been suspected of abnormality, for frank jargon is rare in children and muteness in the early stages of recovery may have been regarded as due to unwillingness rather than to inability to speak.

Dysphasia occurred in 12 of our patients and persisted to an important degree in four. It is interesting that if the brain injury occurred at an early age (before 2 years) and involved the left cerebral hemisphere, several of these patients became left-handed and developed speech normally. Other neurological abnormalities consisted mainly of varying degrees of hemiparesis, and, especially in those with the more severe dynamic head injuries, extrapyramidal signs.

Fatalities

There were 15 deaths (10%), and this number reflects the type of patient that tends to be referred to a neurosurgical department. Nine of these deaths were inevitable, as severe primary or early secondary brain-stem damage was evident when the patient was first seen. One of the other patients, who had an associated fractured femur, died from fat embolism. The most important group were those patients with an extradural haematoma, four of whom died. In each there was delay in instituting treatment.

The fatality rate in the adult series of patients with head injury was two and a half times that of the series in children.

Treatment

If the clinical features or radiological appearances in a patient with a depressed skull fracture suggest brain damage, surgical exploration of the wound is indicated. The depressed bony fragments are elevated and necrotic brain and blood clot removed. A serious contracting scar must be avoided (Penfield and Erickson, 1941).

The brain grows rapidly in the first years of life and therefore intracranial space-occupying lesions must be promptly removed. If not, cerebral atrophy may result, often with important psychological and physical abnormalities (Fig. 3).

Small skull defects in young children recalcify, but

for protective, cosmetic and personal reasons cranioplasty, using tantalum or acrylic, is necessary for large defects. It was carried out in 13 of the series of patients under discussion. In the initial

FIG. 3.—Encephalogram demonstrating enlargement of the left lateral ventricle as a result of cerebral atrophy secondary to delayed evacuation of a subdural haematoma.

stages of a severe head injury, warmth was previously used to treat shock, but nowadays cooling of these patients has been often found to be life-saving.

Less vigorous cooling is necessary in children than is required in adults. Other recent advances in the surgical treatment of certain types of epilepsy—the cause often being a head injury in infancy—include temporal lobectomy and hemispherectomy.

There appears to be no real evidence to suggest that the prophylactic use of anti-epileptic drugs prevents the development of post-traumatic epilepsy in children.

Causes and Prevention of Accidents

There were 56 traffic accidents, seven of the children being cyclists, nine passengers, and 40 pedestrians. Thirty-seven patients fell, and most of the remainder received direct blows to the head by stones, pickaxe, javelin, etc. Children tend to have poor judgment, are impulsive, and the movements of the younger ones are rather incoordinate, and they are therefore susceptible to accidents. There is a striking sex difference (Table 1) in that males predominate in a ratio of 3 to 1.

Many of the accidents occurred because parents permitted young children to play in the street. Education and example are very important in the prevention of accidents to children; however, warnings can be overdone, and the child may develop a 'deaf ear'. The causation and prevention of traffic accidents are fully discussed in 'Medical Aspects of Traffic Accidents' (The Sun Life Building, Montreal, 1955). Dietrich (1954) also discusses the problem.

A few of the patients under discussion appeared to show accident proneness, and, as Langford (1953) points out, this is not quite the same as is found in adults. The children tended to be of the over-active, impulsive type, often with insecure home conditions.

Conclusions and Summary

The main features of head injuries in childhood are discussed, with reference to 150 consecutive patients, and comparisons are made with head injuries in adults. The greater number of male patients was striking (3 : 1). There was a higher percentage of mild head injuries in children (2 : 1), and a correspondingly higher mortality rate in adults (2.5 : 1). Whereas depressed fractures were common in children (3 : 1), basal fractures were infrequent. Hydrocephalic children may be more severely affected by a head injury than normal children. Clinical neurological assessment can be difficult in young children. Greater cooperation of doctors in different specialities may permit of a better understanding of the incidence, significance and management of birth injuries.

Lumbar puncture and electro-encephalography are of limited use in diagnosis. The incidence of post-traumatic epilepsy was 26%; neither obvious cerebral contusion nor an initial unconscious period were necessary for the appearance of epilepsy. Psychological disorders were not uncommon; their diagnosis and significance are discussed. A medical psychologist is an important member of a head injury team. 'Post-concussional syndromes' are rare in children.

Proper treatment of compound fractures of the skull and the timely evacuation of intracranial haematomas are important.

Accident susceptibility, proneness and prevention in relation to head injuries in childhood are discussed.

I wish to acknowledge with gratitude the valuable help given to me by Dr. J. A. L. Naughton, M.C., medical psychologist, in the management of psychological disorders occurring in these patients, and also for his help in the preparation of this paper. I am indebted to Professor N. M. Dott and Mr. F. J. Gillingham for kindly allowing me to include certain of their cases in this series.

REFERENCES

- Anderson, G. W. (1952). *J. Pediatr.*, **40**, 340.
- Bakwin, H. (1949). *Ibid.*, **34**, 371.
- Dietrich, H. F. (1954). *Pediatric Clinics of North America* [Nov.], p. 759. Philadelphia and London.
- Ingraham, F. D. and Matson, D. D. (1954). *Neurosurgery of Infancy and Childhood*. Springfield, Illinois.
- Langford, W. S., Gilder, R. Jr., Wilking, V. N., Genn, M. M. and Sherrill, H. H. (1953). *Pediatrics*, **11**, 405.
- Livingston, S. (1954). *The Diagnosis and Treatment of Convulsive Disorders in Children*. Springfield, Illinois.
- Penfield, W. and Erickson, T. C. (1941). *Epilepsy and Cerebral Localization: A Study of the Mechanism, Treatment and Prevention of Epileptic Seizures*. Springfield, Illinois.
- Rowbotham, G. F., Maciver, I. N., Dickson, J. and Bousfield, M. D. (1954). *Brit. med. J.*, **1**, 726.
- Voris, H. C. (1950). *Arch. Surg. (Chicago)*, **60**, 906.