Accepted 26 February 2006

Published Online First 23 June 2006

Competing interests: None declared.

References

- Miliauskas JR. Bladder xanthomas. *Histopathology* 1992;**21**:177–8.
- Sebo TJ, Bostwick DG, Farrow GM, et al. Prostatic xanthoma: a mimic of prostatic adenocarcinoma. *Hum Pathol* 1994;**25**:386–9.
- Mai KT, Isotalo PA, Guindi M, et al. Intestinal epithelial lesions associated with signet ring cell carcinoma of the colon and small intestine. *Pathology* 2002;**34**:51–6.
- Suster S, Fletcher CDM. Gastrointestinal stromal tumors with prominent signet-ring cell features. *Mod Pathol* 1996;**9**:609–13.
- Lo CY, Lorentz TG, Poon CSP. Xanthogranulomatous inflammation of the sigmoid colon: a case report. *Aust NZ J Surg* 1996;**66**:643–4.
- Miliauskas JR. Rectosigmoid (colonic) xanthoma: a report of four cases and review of the literature. *Pathology* 2002;**34**:144–7.
- Pieterse AS, Rowland R, Labrooy JT. Gastric xanthomas. *Pathology* 1985;**17**:455–7.
- Weinstock LB, Shatz BA, Salzman RJ, et al. Xanthoma of the colon. *Gastrointest Endosc* 2002;**55**:410.
- Colella U, Sturgill BC. Isolated xanthomatosis of the small bowel. *Hum Pathol* 1985;**16**:422–4.
- Goodman MD. Segmental xanthomatosis of the ileocecal valve with anatomic and functional obstruction. *Arch Pathol Lab Med* 1997;**121**:75–8.
- Ashfaq R, Timmons CF. Xanthomatous pseudotumor of the small intestine following treatment for Burkitt's lymphoma. *Arch Pathol Lab Med* 1992;**116**:299–301.
- Fernández-Alonso J, Medina-Pérez M, Gómez-Izquierdo L. Xanthomatous pseudotumour of the stomach after chemotherapy for gastric lymphoma in a patient with acquired immunodeficiency syndrome. *Histopathology* 1994;**25**:589–91.
- Bejarano PA, Aranda-Michel J, Fenoglio-Preiser C. Histochemical and immunohistochemical characterization of foamy histiocytes (mucinophages and xanthelasma) of the rectum. *Am J Surg Pathol* 2000;**24**:1009–15.
- Beutler SM, Fretzin DF, Wellington J, et al. Xanthomatosis resembling scleroderma in multiple myeloma. *Arch Pathol Lab Med* 1978;**102**:567–71.
- Vimala R, Ananthalakshmi V, Murthy, et al. Xanthelasma of esophagus and stomach. *Ind J Gastroenterol* 2000;**19**:135.
- Kimura K, Hiramoto T, Buncher CR et al. Gastric xanthelasma. *Arch Pathol* 1969;**87**:110–17.
- Herrera-Goeppfert R, Lizano-Soberón M, García-Perales M. Verruciform xanthoma of the esophagus. *Human Pathol* 2003;**34**:814–15.
- Kunze KC, Baum RA, Nasrallah SM. Gastric xanthoma. *Gastrointest Endosc* 1987;**33**:114–115.
- Moretó M, Ojembarrena E, Zaballa M, et al. Retrospective endoscopic analysis of gastric xanthelasma in the non-operated stomach. *Endoscopy* 1985;**17**:210–211.
- Kameya S, Nakamura S, Mizutani K. Xanthoma in the stomach. *Gastrointest Endosc* 1963;**5**:37–41.
- Mast A, Elewaut A, Mortier G, et al. Gastric xanthoma. *Am J Gastroenterol* 1976;**65**:311–17.
- Pope TL, Shaffer H. Small bowel xanthomatosis: radiologic-pathologic correlation. *Am J Radiol* 1985;**144**:1215–16.
- Nakasono M, Hirokawa M, Mugaruma N, et al. Colorectal xanthomas with polypoid lesion: report of 25 cases. *APMIS* 2004;**112**:3–10.
- Gaster M, Golden R, Will D, et al. Xanthomatosis involving the colon and small intestine; report of a case. *Am J Dig Dis* 1961;**6**:312–21.
- Boruchowicz A, Rey C, Fontaine M, et al. Colonic xanthelasma due to glyceride accumulation associated with an adenoma. *Am J Gastroenterol* 1997;**92**:159–61.
- Romeu J, Rybak B. Lipid proctitis. *N Engl J Med* 1979;**301**:1099–100.
- Terruzzi V, Minoli G, Butti GC, et al. Gastric lipid islands in the gastric stump and in non-operated stomach. *Endoscopy* 1980;**12**:58–62.
- Berthrong M, Fajardo LF. Radiation injury in surgical pathology. *Am J Surg Pathol* 1981;**5**:153–75.
- Esparza AR, McKay DB, Cronan JJ, et al. Renal parenchymal malakoplakia. Histologic spectrum and its relationship to megalocytic interstitial nephritis and xanthogranulomatous pyelonephritis. *Am J Surg Pathol* 1989;**13**:225–36.
- Michal M, Chlumska A, Mukensnabl P. Signet-ring cell aggregates simulating carcinoma in colon and gallbladder mucosa. *Pathol Res Pract* 1998;**194**:197–200.
- Schiffman R. Signet-ring cells associated with pseudomembranous colitis. *Am J Surg Pathol* 1996;**20**:599–602.
- Sidhu J, Liu D. Signet-ring cells associated with pseudomembranous colitis. *Am J Surg Pathol* 2001;**25**:542–3.
- Chen C-G, Chen C-L, Liu H-N. Primary papular xanthoma of children. A clinicopathologic, immunohistopathologic and ultrastructural study. *Am J Dermatopathol* 1997;**19**:596–601.
- Kumakiri M, Sudoh M, Miura Y. Xanthoma disseminatum. Report of a case, with histological and ultrastructural studies of skin lesions. *Am Acad Dermatol* 1981;**4**:291–9.
- O'Donnell JR, Tansey P, Chung P, et al. Acute myelomonocytic leukaemia presenting as xanthomatous skin eruption. *J Clin Pathol* 1982;**35**:1200–3.
- Zegarelli DJ, Zegarelli-Schmidt EC, Zegarelli EV. Verruciform xanthoma. Further light and electron microscopic studies, with the addition of a third case. *Oral Surg Oral Med Oral Pathol* 1975;**40**:246–56.

Pure cartilaginous teratoma of the testis: an immunohistochemistry and fluorescence in situ hybridisation study

In the testis pure teratomas are rare, accounting for less than 7% of primary testicular tumours.¹ This contrasts with the ovary, where 95% of germ cell tumours are pure teratomas.² Teratomas do not always contain tissues representing all three germ cell layers. A monodermal teratoma is one that consists of only one germ cell layer, endoderm, mesoderm or ectoderm. The best recognised monodermal testicular teratomas are carcinoid tumours and primitive neuroectodermal tumours.³ Monodermal teratoma composed solely of cartilage is rare. We report a second case of purely cartilaginous teratoma and provide interphase cytogenetic evidence of its germ cell origin.

Case report

A 21-year-old man presented with a mass in his right testis. Blood chemistry tests showed an increased level of β -hCG (641 IU/l; normal range 0–3 IU/l), and a normal level of α -fetoprotein (2.4 ng/l; normal range 0.0–8.0 ng/l). The patient underwent orchiectomy followed by chemotherapy. Three months later, retroperitoneal lymph node dissection was done, removing a 275 g paracaval and inter-aortocaval mass, periaortic lymph nodes, and pericaval mass.

Methods

Immunohistochemistry

Sections 4 μ m thick were cut from the paraffin block and stained with H&E. Immunohistochemistry for OCT4 was accomplished with a polyclonal goat anti-OCT4 antibody (C20, sc 8629; Santa Cruz Biotechnology,

Santa Cruz, CA, USA; 1:500 dilution, 30 min at room temperature) directed towards the –COOH terminus of the protein, as previously described.^{4,5} Antigen retrieval was carried out by heating sections in 1 mmol/l EDTA (pH 8.0). Endogenous peroxidase activity was inactivated by incubation in 3% H₂O₂. Non-specific binding sites were blocked using Protein Block (DAKO Corp., Carpinteria, CA, USA). Immunohistochemistry for S100 protein was also performed (primary antibody 1:1000 dilution; 30 min incubation; DAKO). Antigen retrieval was performed by heating sections in pH 6.0 citrate buffer. The positive and negative controls performed appropriately for each procedure.

Fluorescence in situ hybridisation

Fluorescence in situ hybridisation (FISH) was performed on the testicular tumour as previously described.⁶ Sections (4 μ m) were prepared from buffered formalin fixed, paraffin embedded tissue block. The slides were deparaffinised prior to hybridisation. Dual-colour FISH was performed by using a mixture of Spectrum Orange labelled centromeric α satellite DNA probe (CEP12 located at 12p11.1–q11, locus D12Z3) and Spectrum Green labelled subtelomeric DNA probes (Tel12 located at D12S2071) for chromosome 12p (probes from Vysis, Downers Grove, IL, USA).

The slides were examined using a Zeiss Axioplan 2 microscope (Zeiss, Göttingen, Germany) with these filters: SP-100 DAPI, FITC MF-101 for Spectrum Green (12p) and Gold 31003 for Spectrum Orange (CEP12) from Chroma (Chroma, Brattleboro, VT, USA). The images were acquired with a CCD camera and analysed with MetaSystem Isis software (MetaSystem, Belmont, MA, USA). Five sequential focus stacks with 0.4 μ m interval were acquired and then integrated into a single image in order to reduce thickness-related artefacts. The images were adjusted to reduce background through threshold adjustment; however, the relative signal intensity and signal locations were not changed.

From the tumour section, 100 nuclei were scored for signals from CEP12 (red) and 12p (green) under the fluorescence microscope with 1000 $\times$ magnification, and the ratio between green and red signals was subsequently calculated. We analysed the spatial distribution of the green and red signals to detect the specific patterns of signal aggregation consistent with isochromosome 12p (i(12p)), as previously reported.^{1–11} The quantitative criteria to determine 12p overrepresentation were described previously.^{6–10–12} A classical seminoma specimen was used as a positive control for FISH analyses.

Results

Pathological findings

The cut surface of the testis revealed a 20 $\times$ 14 $\times$ 10 mm well-circumscribed, white, lobulated rubbery mass located within the testicular parenchyma. Histological examination revealed the mass to be composed of mature cartilage (fig 1). The remaining testicular parenchyma showed seminiferous tubules containing only Sertoli cells. No intratubular germ cell neoplasia, unclassified type (IGCNU) was visible. A focus of fibrous scar was present outside the cartilaginous tumour suggestive of a burnt-out germ cell tumour.

The retroperitoneal lymph node dissection specimens showed necrosis and inflammation

Figure 1 Low-power view of the well-circumscribed cartilaginous teratoma and the adjacent testicular parenchyma (A). High-power view of the teratoma reveals mature cartilage (B–C). Fluorescence in situ hybridisation displays isochromosome 12p (pointed by arrow) (D) and chromosome 12p overrepresentation (E).

consistent with chemotherapy effect on metastatic germ cell tumour. No viable germ cell tumour was identified.

Immunohistochemical and interphase cytogenetic findings

The cartilaginous tumour gave a negative reaction with antibodies to OCT4 but a positive reaction for S-100 protein. The reaction for OCT4 and S-100 protein was negative in non-neoplastic seminiferous tubules.

Interphase cytogenetics analysis using FISH revealed that the cartilaginous testicular tumour was positive for i(12p) in 4% of cells and also had 12p overrepresentation with 12p/12 centromeric ratio of 1.63 (fig 1).

Discussion

Presented here is a case of monodermal pure teratoma of the testis composed solely of mature cartilage. Only one similar tumour has previously been described.¹³ While the histological appearance leaves open the possibility of a benign cartilaginous tumour and the lack of intratubular germ cell neoplasia also might seem to point to that possibility, the presence of isochromosome 12p and overrepresentation of chromosome 12p are strong

evidence that the tumour is a teratoma.^{14–15} In a study of 17 testicular teratomas by Cheng *et al*,⁶ 76% showed i(12p) and 29% had 12p overrepresentation (overall 88% of teratomas studied revealed 12p abnormalities). As seen in fig 1, this tumour is positive for both isochromosome 12p and 12p overrepresentation. More than 80% of post-pubertal teratomas are associated with intratubular germ cell neoplasia, and intratubular germ cell neoplasia was present in the testis containing the pure cartilaginous teratoma described by Singh *et al*.¹³ The present case is unusual in having no intratubular germ cell neoplasia.

Postpubertal teratomas, regardless of their appearance, have a 22–37% incidence of metastasis. If a scar from a burnt-out germ cell tumour is also present, as in this case, the chance of metastasis increases to 66%.¹ A study of patients with low-stage primary pure teratomas of the testis revealed the overall risk of lymph node metastasis to be 40% at retroperitoneal lymph node dissection. This was also associated with a 16% relapse rate following retroperitoneal lymph node dissection.¹⁶ The pathologist must keep in mind that in patients with pure testicular teratoma, metastases may consist of other types of germ cell tumours.¹⁷

Ryan Des Jean, John N Eble, Shaobo Zhang, Liang Cheng

Department of Pathology and Laboratory Medicine, Indiana University School of Medicine, Indianapolis, Indiana, USA

Correspondence to: Dr Liang Cheng, Department of Pathology and Laboratory Medicine, Indiana University School of Medicine, 350 West 11th Street, Clarian Pathology Laboratory Room 4010, Indianapolis, IN 46202, USA; lcheng@iupui.edu

doi: 10.1136/jcp.2007.048702

Accepted 21 June 2007

Published Online First 29 June 2007

Competing interests: None declared.

References

- Woodward PJ, Heidenreich A, Looijenga LHJ, *et al*. Germ cell tumors. In: Eble JN, Sauter G, Epstein JI, Sesterhenn IA, eds. *World Health Organization classification of tumours: pathology and genetics of tumours of the urinary system and male genital organs*. Lyon: IARC Press, 2004:221–49.
- Ulbricht TM. Germ cell tumors of the gonads: a selective review emphasizing problems in differential diagnosis, newly appreciated, and controversial issues. *Mod Pathol* 2005;**18**:S61–79.
- Ulbricht TM, Amin MB, Young RH. *Tumors of the testis, adnexa, spermatic cord, and scrotum*. Washington, DC: Armed Forces Institute of Pathology, 1999.
- Jones TD, Ulbricht TM, Eble JN, *et al*. OCT4 staining in testicular tumors: a sensitive and specific marker for seminoma and embryonal carcinoma. *Am J Surg Pathol* 2004;**28**:935–40.
- Cheng L, Sung MT, Cossu-Rocca P, *et al*. OCT4: biological functions and clinical applications as a marker of germ cell neoplasia. *J Pathol* 2007;**211**:1–9.
- Cheng L, Zhang S, MacLennan GT, *et al*. Interphase fluorescence in situ hybridization analysis of chromosome 12p abnormalities is useful for distinguishing epidermoid cysts of the testis from pure mature teratoma. *Clin Cancer Res* 2006;**12**:5668–72.
- Coventry S, Punnett HH, Tomczak EZ, *et al*. Consistency of isochromosome 7q and trisomy 8 in hepatosplenic gamma/delta T-cell lymphoma: detection by fluorescence in situ hybridization of a splenic touch-preparation from a pediatric patient. *Pediatr Dev Pathol* 1999;**2**:478–83.
- Meng FJ, Zhou Y, Giwerzman A, *et al*. Fluorescence in situ hybridization analysis of chromosome 12 anomalies in seminoma cells from patients with carcinoma in situ of the testis. *J Pathol* 1998;**186**:235–9.
- Velagaleti GV, Tapper JK, Rampy BA, *et al*. A rapid and noninvasive method for detecting tissue-limited mosaicism: detection of i(12)(p10) in buccal smear from a child with Pallister-Killian syndrome. *Genet Test* 2003;**7**:219–23.
- Cossu-Rocca P, Zhang S, Roth LM, *et al*. Chromosome 12p abnormalities in dysgerminoma of the ovary: a FISH analysis. *Mod Pathol* 2006;**19**:611–15.
- Poulos C, Cheng L, Zhang S, *et al*. Analysis of ovarian teratomas for isochromosome 12p: evidence supporting a dual histogenetic pathway for teratomatous elements. *Mod Pathol* 2006;**19**:766–71.
- Kernek KM, Brunelli M, Ulbricht TM, *et al*. Fluorescence in situ hybridization analysis of chromosome 12p in paraffin-embedded tissue is useful for establishing germ cell origin of metastatic tumors. *Mod Pathol* 2004;**17**:1309–13.
- Singh N, Cumming J, Theaker JM. Pure cartilaginous teratoma differentiated of the testis. *Histopathology* 1997;**30**:373–4.
- Blough RI, Heerema NA, Ulbricht TM, *et al*. Interphase chromosome painting of paraffin-embedded tissue in the differential diagnosis of possible germ cell tumors. *Mod Pathol* 1998;**11**:634–41.

- 15 Summersgill B, Goker H, Osin P, *et al*. Establishing germ cell origin of undifferentiated tumors by identifying gain of 12p material using comparative genomic hybridization analysis of paraffin-embedded samples. *Diag Mol Pathol* 1998;7:260-6.
- 16 Leibovitch I, Foster RS, Ulbright TU, *et al*. Adult primary pure teratoma of the testis: the Indiana experience. *Cancer* 1995;75:2244-50.
- 17 Simmonds PD, Lee AH, Theaker JM, *et al*. Primary pure teratoma of the testis. *J Urol* 1996;155:939-42.

Cytophagic histiocytic panniculitis with fatal haemophagocytic lymphohistiocytosis in a paediatric patient with perforin gene mutation

Cytophagic histiocytic panniculitis (CHP), or histiocytic cytophagic panniculitis, is a rare form of panniculitis associated with haemorrhagic diathesis and histiocytic lymphohistiocytosis (HLH), initially described in 1980 as a benign lymphoproliferative disease.¹ In 1991 Gonzalez *et al* reported a unique entity of subcutaneous T-cell lymphoma with haemophagocytosis, later designated as subcutaneous panniculitis-like T-cell lymphoma (SPTL).² Marzano *et al* suggested that CHP and SPTL might span a clinicopathological spectrum in which there is a natural progression from CHP to SPTL.³ HLH is a clinical syndrome of immune deregulation with hypercytokinaemia causing dysfunctions of various organs and a high mortality. Familial HLH (FHL) is associated with several hereditary defects. Stepp *et al* first showed that the mutation in perforin gene (*PRF1*) at chromosome 10q21 was responsible for 20–40% of FHL patients.⁴

Case report

An 11-year-old girl with CHP and HLH presented with spiking fever and indurated skin nodules over the left thigh (fig 1A). She had psychomotor retardation and spastic type cerebral palsy at 2 years of age and epilepsy at age 7. Physical examination revealed indurated skin nodules and hepatosplenomegaly without lymphadenopathy. Laboratory tests showed anaemia, thrombocytopenia, impaired liver function, and raised triglyceride level, without coagulopathy. Blood and urine cultures were negative. Her fever and skin lesions responded dramatically to prednisolone and intravenous immunoglobulins. She became prednisolone-dependent despite attempts at tapering off by adding methotrexate and cyclosporine A. Several episodes of HLH and infections led to three additional admissions in three years. Six months later, she developed prominent facial bruising and swelling. Virological surveys revealed prior/remote Epstein-Barr virus (EBV) infection without infection by hepatitis viruses A, B, C, or cytomegalovirus. Whole body CT scans revealed brain atrophy, pericardial effusion, and ascites without lymphadenopathy. Three courses of 13-cis retinoic acid with prednisolone led to regression of skin lesions. Unfortunately, she developed spiking fever and died of coagulopathy with acute abdominal bleeding and hypovolaemic shock, four years after initial presentation.

Pathological and molecular genetic findings

The skin biopsies at presentation and at 3 years revealed significant lobular panniculitis with

Figure 1 (A) Photograph showing plaques and nodules on both thighs. (B) Photomicrograph showing a lymphohistiocytic infiltration involving predominantly subcutaneous fat with one bean bag-histiocyte in the centre (H&E); inset shows a bean bag-histiocyte highlighted by anti-CD68 (immunohistochemical staining). (C) Photomicrograph showing small lymphocytic infiltration in a lace-like fashion rimming fat spaces with high-power in the inset (H&E). The majority of these infiltrating lymphocytes express: (D) T-cell intracellular antigen-1 (immunohistochemical stain for T-cell intracellular antigen-1) and granzyme B; (E) with only a few cells expressing perforin (immunohistochemical stain for perforin). (F,G) Photomicrograph at autopsy showing florid haemophagocytosis in the marrow (F) and late-stage neurodegenerative changes in the frontal grey matter with marked neuronal loss and increased number of reactive astrocytes indicated by arrows (G) (H&E). Parental/guardian informed consent was obtained for publication of this figure.

small lymphocytic infiltration within the lobular septa and around the individual fat cells, with occasional benign histiocytes showing haemophagocytosis (fig 1B). These small lymphocytes exhibited slightly irregular nuclear contours with hyperchromasia (fig 1C). Immunohistochemically, nearly all the infiltrating lymphocytes expressed CD2, CD3, CD5, CD7, CD8, T-cell intracellular antigen-1 (fig 1D), and granzyme B, but not CD4 or CD56. Repeated immunostainings showed that the majority of the lymphocytes were negative for perforin (fig 1E). These infiltrating lymphocytes were negative for EBV by in situ hybridisation. Clonality study using paraffin sections from skin biopsy with PCR for T-cell receptor- γ chain (TCR γ) gene rearrangement was performed in duplicate with appropriate controls; the result was polyclonal. Bone marrow aspirations at presentation and at 3 years revealed histiocytic haemophagocytosis, with the latter showing more significant haemophagocytosis and T-cell infiltration.

At autopsy she was 45.5 kg and 137 cm with moon face, ecchymosis, ulcerated plaques in the extremities, bloody ascites, enlarged liver (1750 g) and spleen (180 g). Sections revealed lobular panniculitis and fulminant haemophagocytosis in liver and bone marrow (fig 1F) without lymphoma. Sections of the brain revealed prominent neuronal loss with gliosis and spongiosis in the cortex of the frontal (fig 1G), temporal, parietal, and occipital lobes, consistent with a late-stage neurodegenerative process. Cloning and sequencing for *PRF1* using DNA extracted from bone marrow at diagnosis as previously described⁵ showed a heterozygous nonsense point mutation with 1168 C>T (Arg390stop) (fig 2). The same assay using peripheral blood from the patient's healthy mother and only sister was negative.

Discussion

To our knowledge, there has been no genetic study on patients with CHP associated with HLH. Recent progress has clarified four types of