Saline suspensions of both samples were *not* agglutinated by one anti-C serum free from incomplete anti-D, five anti-E sera free from incomplete anti-D, five anti-c sera, three anti-e sera, one anti-C^w serum, and three samples of incomplete anti-D. Neither sample of red cells reacted with two incomplete anti-C sera free from incomplete anti-D (from Rh-positive donors). Neither sample reacted with the serum of the original $-D-/-D-$ patient, Mrs. B.: this serum had agglutinated all of more than 1,400 samples of Group O red cells, and was shown to contain anti-C, anti-c, and anti-e (Race, Sanger, and Selwyn, 1950, 1951).

Saline suspensions of red cells of the original $-D-/-D-$ patient, Mrs. B., had been agglutinated by 14 out of 16 incomplete anti-D sera. Dr. Philip Levine, of Ortho Research Foundation, has tested saline suspensions of the red cells of Mrs. V. W. against titrations of 13 incomplete anti-D sera; the results are shown in the Table; the cells were agglutinated by 11 of the 13 sera. The agglutination of these cells, however, is not directly related to the titre of incomplete anti-D, as is indicated by the striking differences observed in the behaviour of the five sera of highest titre. Dr. Levine showed that the absence of C from the red cells of Mrs. V. W. could also be demonstrated with the aid of a complete anti-D+anti-C serum; he found that the anti-D component could be successfully blocked by the addition of selected incomplete anti-D (see Table).

Differences in Behaviour of Various Anti-D Sera. Tests by Dr. Philip Levine with 13 Sera Containing Incomplete Anti-D

Titre	No. of sera	Titration with cells of Mrs. V. W. in saline
1/4	 1	1/2
1/8	 1	1/1
1/16	 1	1/8
1/32	 1	1/2
1/64	 3	1/4, 1/4, 1/64
1/128	 1	1/4
1/512	 5	0, 0, 1/2, 1/128, 1/256

The Serum of Mrs. V. W.

The serum of Mrs. V. W. has reacted with all of approximately 250 unselected samples of Group O red cells so far tested; it does not react with those of her brother J. C. or with those of Mrs. B. The serum reacts only weakly with cells suspended in saline but strongly by the indirect antiglobulin, the albumin, and the trypsin methods.

Titration of the serum of Mrs. V. W. showed values of 1/128 to 1/256 with the indirect antiglobulin test. Titration with bloods of various genotypes failed to reveal direct evidence of a mixture of antibodies. Absorption experiments now being done present considerable difficulties because of the small quantities of serum available and a gradual diminution of titre in more recent bleedings. So far only anti-e has been isolated, after absorption with cDE/cDE red cells. In view of the parallelism with the blood of Mrs. B. it is very likely that the high incidence of positive reactions is due to the presence of antibodies for both C and c and perhaps E as well as e. Under such conditions all bloods would be incompatible except those with the identical genotype $-D-/-D-$. No atypical antibodies could be demonstrated in the serum of the compatible sibling, J. C.

As in the case of Mrs. B., there is a history of cousin marriage: the parents of Mrs. V. W., each of whom must have contributed the chromosome $-D-$, were first cousins. Efforts are being made to trace any consanguineous relationship between the British and American families. Further studies are planned on three additional surviving siblings, five others having died—three from tuberculosis, one from influenza, and one from accidental causes.

It may be that the chromosome $-D-$ is not as rare as it was first thought to be; it may, for example, be no rarer than the chromosome Cde. In the heterozygous state the chromosome $-D-$ would usually escape detection, though it can be recognized if it is specially sought for (Race, Sanger, and Selwyn, 1951).

Summary

Two further examples of blood representing the Rh genotype $-D-/-D-$ are described. The samples come from a white American woman and her brother, offspring of a first-cousin marriage.

The sister has had five blood transfusions. Her serum is incompatible with all of about 250 unselected samples of Group O blood, but it is compatible with her own red cells, her brother's, and with those of the original $-D-/-D-$ British patient. Of the several antibodies which are doubtless present only anti-e has so far been identified. No unusual antibodies could be found in the serum of the brother.

The red cells of the two Americans are the only samples so far found to be compatible with the serum of the British patient, besides her own, though more than 1,400 samples have been tested.

We wish to thank Mrs. V. W.'s physician, Dr. I. N. Hunnicutt, of Virginia, for his active help in this investigation.

REFERENCES

- Race, R. R., Sanger, Ruth, and Selwyn, J. G. (1950). *Nature, Lond.*, 166, 520.
— — — (1951) *Brit. J. exp. Path.*, 32, 124.

Medical Memoranda

Uterine Chorionepithelioma

The possible development of chorionepithelioma following hydatidiform mole is well recognized, although there is a wide variation in estimates of the incidence of the malignant change. In the follow-up of cases of hydatidiform mole the biological tests for the excretion of gonadotrophin in the urine occupy a prominent position, but the interpretation of their results presents problems which are not widely enough understood, despite the excellent summary by Brews (1950). The following case illustrates the difficulty of diagnosis when the Aschheim-Zondek reaction is only weakly positive.

CASE REPORT

The patient, a 1-para aged 32, had had an uneventful pregnancy three years previously. Her last menstrual period began on February 1, 1950, and very early in the pregnancy she attempted to procure a miscarriage by taking a large dose of quinine. At the ninth week of amenorrhoea vaginal bleeding occurred and the case was treated as one of threatened abortion. It was noted at this time that the uterine enlargement corresponded to that of a 12-weeks pregnancy. The diagnosis of hydatidiform mole was made at the eleventh week, when the fundus of the uterus had arisen to midway between the pubic symphysis and the umbilicus. At this stage the early morning urine gave a positive Aschheim-Zondek reaction in a dilution of 1 in 10, but was negative at 1 in 100.

Vaginal evacuation of the uterus was effected on April 22 after dilatation of the cervix by laminaria tents. The tumour removed was a typical hydatidiform mole with characteristic microscopical appearance. There was no evidence of malignancy in the sections. Twenty days later the uterus had involuted satisfactorily and the Aschheim-Zondek reaction had become negative. But further biological tests were made at intervals, and 57 days after evacuation it was found that the Aschheim-Zondek reaction was again positive, though negative in dilution.

The patient was readmitted to hospital and examined under an anaesthetic on July 7. The uterus was bulky, the cavity measuring 3½ in. (9.5 cm.), and the left ovary was

slightly enlarged. With a ring forceps, fragments of apparently typical hydatidiform mole were removed, and then a curettage was performed. Microscopy revealed considerably more trophoblastic activity than in the original specimen, with variations in nuclear size and density, and occasional mitoses. No villi were seen in the curettings. Some bleeding persisted, and one week later the Aschheim-Zondek reaction was still positive, though remaining negative in dilution. X-ray examination of the chest showed no abnormal shadows.

In view of the recurrence of the mole with atypical trophoblastic activity, total hysterectomy with bilateral salpingo-oophorectomy was performed on July 21—that is, 91 days after evacuation of the mole. Both ovaries contained small luteinized and atretic follicles; in the right ovary there was also a recently ruptured small lutein cyst. The uterus was bulky and contained a haemorrhagic tumour approximately $\frac{1}{2}$ in. (1.3 cm.) in diameter near the right cornu. The growth extended from the endometrium through approximately half the thickness of the uterine wall. Microscopically the growth was a typical chorionepithelioma with malignant trophoblastic elements invading the myometrium. Sections of the ovaries and uterine tubes revealed no secondary deposits.

The patient's progress was uneventful; at examinations 7 and 16 weeks after operation there was no clinical evidence of recurrence, and on each occasion the Hogben test was negative. Subsequent direct follow-up has not been possible, as the patient went abroad, but a letter reports that she was well in January, 1952.

COMMENT

In 1939 Brews observed that as a method of absolute diagnosis of active chorionepithelioma quantitative hormone assays had failed to realize initial expectations. By that time cases had been reported in which the biological tests for urinary gonadotrophin had remained negative in the confirmed presence of a tumour. Cuscaden and Bettinger (1940), in describing a case in which certain features closely resembled that now reported, refer to the possible fallacies of the biological tests and maintain an attitude of great caution in accepting reports of cases of chorionepithelioma without a positive Aschheim-Zondek reaction. The possibility of a negative pregnancy test in the presence of chorionepithelioma is mentioned by Baird (1950) and Browne (1950), while Sutherland (1951) has collected 19 reports of such cases from the literature, and Richli (1950) describes another in detail.

In trying to explain the negative and only weakly positive reactions the factors to be considered include haemorrhage, necrosis, thrombosis, and fibrous reaction; all of these may separate the tumour from healthy maternal tissue and so limit the absorption of gonadotrophic hormone. Browne (1950) suggests that the degree of anaplasia present in the cytrophoblast may determine the strength of the Aschheim-Zondek reaction. Sutherland (1951) attempts to relate the very weakly positive reaction to an almost complete absence of syncytium, although he admits that this is not in accordance with the experimental evidence of Jones, Gey, and Gey (1943), who found that the Langhans cells were apparently responsible for the production of chorionic gonadotrophin.

The cases described by Browne (1930), Ehrhardt and Bureck (1939), Scott (1939), and Brews (1950) illustrate the dangers of waiting to establish a firm histological diagnosis in the presence of a persistently positive pregnancy reaction after evacuation of a hydatidiform mole. When the reaction has become negative soon after expulsion of the mole, and later in the absence of a new pregnancy it again becomes positive, radical treatment is indicated. This view, however, is not shared by Siegler, Ravid, and Tobin (1950), who, in presenting a review of 12 cases of chorionepithelioma and acknowledging the fallibility of the biological tests, advise that, "so long as the titre is decreasing or at least remaining constant, watchful conservatism may still be

applied." Such a recommendation appears highly dangerous, bearing in mind the fatal cases of Brews and of Ehrhardt and Bureck, referred to above.

Shively (1950) assesses the reliance to be placed upon quantitative urinary gonadotrophin determinations by saying that, because of the variations encountered, they should be used only as confirmatory evidence in the diagnosis of these neoplasms and should not be regarded as necessary diagnostic criteria.

Publication of this case report has been approved by the Director-General, Army Medical Services.

P. A. TRAFFORD, M.B., B.S., Major R.A.M.C.,

Military Families Hospital, London, S.E.18.

REFERENCES

- Baird, Dugald (1950). *Combined Textbook of Obstetrics and Gynaecology*, 5th ed., p. 353. Livingstone, Edinburgh.
 Brews, A. (1939). *J. Obstet. Gynaec. Brit. Emp.*, **46**, 813.
 — (1950). *Ibid.*, **57**, 317.
 Browne, F. J. (1930). *Proc. roy. Soc. Med.*, **24**, 1628.
 — (1950). *Postgraduate Obstetrics and Gynaecology*, p. 355. Butterworth, London.
 Cuscaden, W. G., and Bettinger, H. F. (1940). *Med. J. Aust.*, **2**, 377.
 Ehrhardt, K., and Bureck, P. (1939). *Z. Geburtsh. Gynäk.*, **119**, 133.
 Jones, G. E. S., Gey, G. O., and Gey, M. K. (1943). *Bull. Johns Hopk. Hosp.*, **72**, 26.
 Richli, R. (1950). *Gynec. Obstet.*, **49**, 299.
 Scott, W. A. (1939). *Amer. J. Obstet. Gynec.*, **38**, 837.
 Shively, J. A. (1950). *U.S. armed Forces med. J.*, **1**, 505.
 Siegler, S. L., Ravid, J. M., and Tobin, S. M. (1950). *Amer. J. Obstet. Gynec.*, **60**, 586.
 Sutherland, A. M. (1951). *J. Obstet. Gynaec. Brit. Emp.*, **58**, 29.

Traumatic Encephalopathy from Boxing

The risk of serious intracranial damage occurring in boxing is well known, and the following account may be of interest to those concerned with the welfare and medical examination of boxers.

CASE REPORT

A miner aged 26 had boxed on numerous occasions since the age of 15 and as a professional from the age of 20. An ample history, obtained from various sources, failed to reveal evidence of previous mental disorder or any other abnormality, except a minor degree of mental dullness indicated by his school work and Army records. His I.Q. was reported to have been 90 on the Stanford-Binet scale (Form L) when he was tested at some unspecified time before leaving school at the age of 14.

Two years ago, after becoming cruiser-weight champion of his area, slight changes in character became apparent, shown in the form of egocentricity and of monopolizing the conversation with accounts of the various opponents whom he had knocked out. Soon he was knocked out himself after only 18 seconds in the first round, and three months later, in an encounter with a boxer thought by all to be inferior, he took so much punishment that the fight was stopped in the fourth round. His friends were now finding him more quarrelsome and garrulous; he often complained of headaches, and he expressed mild paranoid ideas about his wife which eventually resulted in his admission to a mental hospital a year ago, after he had made threats to kill her with the carving-knife.

He was then found to be slightly confused, was alternately excited or unduly reticent, and had also a slight spatial and temporal disorientation, misidentification of persons around him, confabulation with an associated patchy memory defect for events of the preceding four months, and fleeting olfactory hallucinations. Abnormal physical signs were slurred speech, absent abdominal reflexes on the left side, and a slightly ataxic gait of cerebellar type. The cerebrospinal fluid and an x-ray film of the skull were normal.

Gradual improvement took place over a period of six months with sedation and rehabilitation, and he was then