REFRESHER COURSE FOR GENERAL PRACTITIONERS

RECURRENT BOILS

BY

S. T. ANNING, M.A., M.D., M.R.C.P.

Assistant Physician, Dermatological Department, General Infirmary at Leeds

A boil, or furuncle, is a localized staphylococcal infection of a skin appendage, usually a pilo-sebaceous follicle or a sweat gland. An abscess occurs with central necrosis, followed after a few days by discharge of pus and sloughing of the follicle or gland. Healing is usually rapid. In a carbuncle many skin appendages are infected and the inflammation spreads more deeply, affecting the subcutaneous tissues. A local crop of boils results from infection of near-by follicles, but with wide-spread boils we have a contagious disease affecting any part of the skin surface. The staphylococcus causing these infections is ubiquitous and can be found in the noses of half of the adult population and on the skin of a quarter of them.

Except as regards local treatment, the problem of recurrent boils is much the same whether the affection is generalized or localized to one area, such as the neck or axilla. The problem has serious and important facets. The presence of boils means a high population of virulent staphylococci over the whole skin of the patient, who then becomes a danger to himself and others, indeed to the community. It is a general medical problem in its widest sense, but the possibility of a septic focus in the patient and the reasons for his lowered resistance must be investigated. There is also the question of the indication for chemotherapy and antibiotics, which on the face of it would seem to be the specific line of treatment.

Resistance to Infection

The skin forms an efficient barrier to infection, and when its surface is unbroken it can be penetrated by bacteria only through the openings of follicles and glands. The acidity of the secretions and the keratin layer provides an "acid mantle" which controls the bacterial flora on the skin surface. Once organisms have penetrated the skin the humoral and cellular resistance of the body provides further protection; the development of specific immunity is stimulated.

Friction reduces locally the resistance of the skin to infection. Thus rowers using fixed seats and riders who are inexperienced get boils on the buttocks; rough sleeves or collars may provoke boils on wrists or neck; scratching from pruritus ani may be associated with recurrent boils on the perineum. It is as well to think of scabies when there are multiple boils on the buttocks. But any itching skin disease, whether it be eczema, urticaria, pediculosis, or some other, may be complicated by boils, as may any general disease associated with itching (jaundice, Hodgkin's disease, leukaemia, etc.).

The resistance of the skin to infection is reduced by moisture and excessive oiliness, because these disturb the acid mantle. The moist, greasy skin of the seborrhoeic subject is particularly prone to infection, and maceration of the skin brought about by profuse sweating, wet occupations, and moist applications lowers its resistance. "Salt-water boils" are seen on the wrists of sailors and fishermen whose skin, wet with sea-water, is chafed by the stiff cuffs of their oilskins. When work requires prolonged contact of the skin with cutting oils, lubricants, greases, and waxes, oil acne and boils may appear, especially on the hairy

parts such as the forearms and thighs. Tar workers often suffer with recurrent boils. Alkalis which are keratolytic and neutralize the acid mantle also lower the skin's resistance.

The presence of a boil leads to an enormous increase in the bacterial population of the skin, and infection of other pilo-sebaceous follicles becomes more likely. The population is also increased when there is dental or nasal sinus infection or when other septic foci are present. The old dictum should be remembered: look in the mouth, nose, and throat whenever there is skin sepsis. It has been shown that with tinea pedis there is a local increase in the number of staphylococci, and a general increase may follow contact with another individual suffering from boils.

More important, probably, than the organisms on the skin surface is a lowering of general resistance to infection. In an annotation on pyococcal dermatoses (*British Medical Journal*, 1949, 1, 858) it was stated that a psychogenic, endocrine, nutritional, or toxic influence plays a major part in these conditions. This would seem to be true, and the order in which they are placed should be noted.

In many patients with recurrent boils the condition appears during a period of mental strain or emotional distress. The lowering of resistance to infection in this way need cause no surprise and it is generally appreciated by the lay public. In plague epidemics in times past it was recognized that those most fearful of infection were the most likely to fall victims. Endocrine influences play a part which is not well understood, but the frequency of boils in adolescence seems related to the increased activity of the sebaceous glands from hormonal stimulation. Some people are particularly predisposed to boils, which appear as their reaction to stress of any kind. The reason for this is not known, but is probably associated with endocrine disturbances.

Lowering of the resistance of the skin to infection is associated more often with malnutrition than undernutrition, and particularly with excessive carbohydrate intake. Any disorder of metabolism may lower resistance, but patients with diabetes mellitus are thought, though on no very firm evidence, to be especially prone to skin infection. Whether or not this be true, the diagnosis of diabetes mellitus in a patient with skin infection is of the utmost importance because of the greater insulin requirement and the danger of coma. The urine of a patient with boils must always be examined—it may lead to the discovery of diabetes, perhaps previously unsuspected. However, boils and carbuncles may produce a transient toxic glycosuria which is of no importance. Other general debilitating conditions, such as jaundice, anaemia, nephritis, and tuberculosis, also reduce the resistance of the skin to infection.

Complications

Complications have been less common since the advent of chemotherapy and the antibiotics. Mild debility and local complications such as lymphangitis and lymphadenitis are common, but staphylococcal septicaemia or septic embolism causing carbuncle of the kidney and perinephric abscess, osteomyelitis, or cerebral abscess are not often encountered. It is well known that thrombosis of the intracranial venous sinuses may complicate boils on the head, especially on the face, by direct extension. It is less generally appreciated

that it may occur with boils on any part of the body as a result of a septic embolism. If cooks or others who touch food have boils, outbreaks of food-poisoning may occur. The precipitation of coma in a diabetic with boils or carbuncles has already been mentioned.

Diagnosis

In the diagnosis of recurrent boils there is little difficulty except to determine the cause, and this requires a full assessment of the underlying factors. Unless a careful history is taken and the patient is examined fully a blunder will occasionally be made. A bromide eruption with small papulo-pustules developing into sterile boils, pustular ringworm (kerion) with discrete lesions, papulo-pustular syphilide, and papulo-pustular tuberculide may all be mistaken for boils if care is not taken. Cutaneous anthrax, which is seen occasionally in this country in leather workers or others who handle hides, might be mistaken for a single boil but not for recurrent ones. A papule, rapidly becoming a pustule, appears usually on an exposed area of skin. There is intense local inflammation, necrosis of the pustule follows with the formation of a black slough, and secondary vesicles soon form around it.

General Treatment

The treatment of widespread boils clearly demands general measures, but even in grouped or localized boils these must not be neglected. The treatment of any patient with recurrent boils must be considered as a general medical problem and the underlying factors dealt with. We must think in terms of focal sepsis and the resistance of the patient to infection. Chemotherapy and the antibiotic drugs provide specific therapy and are most impressive in the treatment of boils. Those occurring on the face should always be treated in this way. Unfortunately, in suppressing the staphylococcus so effectively they do not allow the process of active immunity to be developed, and a relapse often follows cessation of treatment. Care must be taken that concentration on local treatment combined with injections of penicillin does not lead to neglect of the cause of the problem.

Whether the boils are localized in groups or whether they are widespread, the possibility of focal sepsis should be considered. Infection of the teeth, nose, nasal sinuses, or throat must receive appropriate treatment. Tinea pedis must not be neglected.

Whenever emotional or mental strain plays a part rest will be required, possibly a holiday from work or away from home. Mild sedation is helpful in such patients, and phenobarbitone, $\frac{1}{4}$ gr. (16 mg.) twice daily by mouth, may usefully be given.

The question of employment must be considered. The seborrhoeic type of individual tolerates heat and moisture badly, and skin sepsis is common in miners of this type who work under such conditions. After greasy work the skin should be washed thoroughly with soap and water. Sponging with cetrimide as a 1% solution in water removes grease and dirt from the skin and is actively bactericidal. It is inactive in the presence of soap and occasionally irritates the skin, but otherwise is helpful if the patient can be persuaded to go to the trouble of using it.

A balanced diet is important. Sweet cakes, suet puddings, fried potatoes, and other carbohydrates should be eaten in moderation. Chocolate is best avoided. Sugar in tea and coffee may be replaced by saccharin. Citrus fruits and fresh vegetables may be eaten in abundance. Additional vitamin-B complex is helpful. Yeast is not of much value unless it is fresh and taken in large quantities. Patients with boils are sometimes anaemic, and iron may be given—for example, one or two ferrous sulphate tablets three times a day after meals.

Active immunization by an autogenous staphylococcal vaccine or staphylococcus toxoid is found to be effective by some physicians, toxoid appearing to be the better. It is as well to start a course when there are no active boils. A combination of toxoid and a stock vaccine such as "vac-coid" may be used, 0.1 ml. of the weak preparation being enough for the first dose. The material is injected intradermally and the dose increased by 0.1 ml. each week until 0.5 ml. is reached, when the stronger preparation, containing five times the quantity of toxoid, may be used in a similar manner, beginning once more with a dose of 0.1 ml. Reactions, general or local, are to be avoided, and if such occur the dose for the following injection should not be increased.

If boils do not stimulate the production of active immunity it seems unlikely that vaccines will do so unless they provide a stronger stimulus, whether specific or non-specific. Some authorities believe the action of vaccines and toxoid to be one of desensitization to the staphylococcus and its toxins. However, treatment with vaccines or toxoid, or with manganese or tin, does not provide a cure, and, like antibiotic therapy, will prove disappointing unless the essential cause is found and treated.

Local Treatment

The principles of local treatment are rest, dry heat, and the minimum of interference. Because it is important to prevent, if possible, the infection of other follicles, the surrounding skin should be kept healthy and as sterile as practicable. The skin around a boil should be swabbed with spirit or brilliant green paint (perchloride of mercury 0.5%, brilliant green 0.5%, industrial spirit 99%). The topical application of antibiotics is only of temporary value, and may lead to sensitization of the skin. Hot fomentations and bread or kaolin poultices should be avoided, as these may burn the skin and certainly make it sodden and more liable to infection. Adhesive plaster dressings also make the skin unhealthy. Dry heat is useful; an infra-red lamp or an old-fashioned electric radiator or even a hot-water bottle over dry dressings may be employed. Short-wave diathermy, if available, is helpful. Local fractional doses of superficial x rays seem to abort the condition in the early stage and to prevent crops of boils appearing in the area treated.

The squeezing of boils is best avoided. Surgery has little place in the treatment, but when a boil becomes fluctuant, and not before, it may be incised. Large cruciate incisions and packing are quite unnecessary and do more harm than good. A dry dressing to absorb the pus may be applied and changed frequently, though if preferred magnesium sulphate paste may be used.

Widespread Boils

Once a boil has burst there is a great increase in the bacterial population of the skin near by and over the body as a whole. With lowered resistance this may result in recurrent widespread boils, and the condition must be treated as a contagious disease in which there is superficial external infection and reinfection. Every precaution possible to prevent the continuation of autogenous reinoculation must be taken. The bacterial increase can be controlled to some extent by bathing in or sponging with 1% solution of cetrimide or with 1 in 4,000 potassium permanganate. The permanganate must be dissolved in a little water before addition to the bath, as contact of a crystal with the skin may cause burning. Potassium permanganate discolours the bath, which is cleaned with difficulty, though the brownish stain can be removed with vinegar, oxalic acid, or some of the abrasive cleansers. Antiseptic soaps may be preferred, but they do not have the toughening and tanning effect of permanganate, and they may neutralize the acid mantle. After such antiseptic baths a dusting powder

containing 10% of calomel in zinc oxide is especially useful with widespread furunculosis, provided the patient is not sensitive to mercury. Cotton underclothes, changed daily, should be worn, as they can be boiled. Soiled dressings must not be allowed to touch skin areas which have not been protected by disinfectant solutions, and the hands of the patient and those attending him should be kept scrupulously clean.

The bacterial population of the skin remains high for weeks after a boil has healed, and these efforts to reduce the staphylococcal flora must be continued for some time. The nose of the patient is likely to be harbouring numerous virulent staphylococci, although physical signs are absent. Careful cleansing of the nasal vestibule with soap and water several times a day or a thin smear of ammoniated mercury 10 gr. (0.65 g.) in hydrous ointment 1 oz. (30 g.) reduces the danger from this source. In the presence of sinus infection this measure is, of course, purely palliative, and, as mentioned before, treatment of the nose and sinuses may be required.

Grouped Recurrent Boils

Hidradenitis suppurativa is an example of recurrent boils appearing as a local condition. The apocrine sweat glands develop at the time of puberty in the axillae, on the nipples, and in the region of the anus and genitals. They are subcutaneous and their ducts usually open into hair follicles.

A troublesome inflammation of these glands occurs occasionally and is more common in women. One or both axillae may be affected, and sometimes the perineum and groins. In these areas recurrent abscesses may result. A tender subcutaneous swelling appears, with redness of the overlying skin. Although it may settle down without supuration, it often becomes fluctuant and discharges pus after bursting. Other glands become infected, and the condition may continue for weeks with great discomfort. Irregular scarring with cheloid formation may result.

It is probable that there is sweat retention followed by secondary infection. Although psychological and endocrine factors play an important part, mechanical causes may be present, such as maceration of the skin from sweat, especially when rubber dress protectors are worn. Damage to the duct openings by deodorant applications or depilatories may be a cause.

As with skin infection elsewhere, the cause of the breakdown of the patient's resistance must be determined. Antibiotics are only helpful temporarily, because a relapse often occurs when their use is stopped. The application of brilliant green paint (see above) to the whole of the axilla or other affected area, together with irradiation by x rays in a few fractional doses at intervals, will usually clear the disease rapidly, provided general causes are not neglected. As with other boils, fomentations are best avoided and dry heat should be applied. Short-wave diathermy is helpful. Incision may have to be considered, but it is wise to be conservative in order to avoid scarring and cheloid formation. Many of the swellings settle with treatment, and incision is justified only if there is marked fluctuation over an abscess. Full consideration of the treatment, which may involve endocrine therapy or even excision and grafting, is outside the scope of this article.

Conclusion

The objectives in the treatment of recurrent boils should be to keep the skin dry, clean, and free from irritation, to eradicate foci of infection, and to raise the general resistance of the patient.

Next Refresher Course Article.—"Prolapse," by Mr. A. J. Wrigley.

HEALTH SERVICE COSTS

[FROM A SPECIAL CORRESPONDENT]

A year ago, Mr. Crookshank, then Minister of Health, was being hard put to it to maintain the £400m. limit on the Budget's share of Health Service expenditure set by Labour Chancellors of the Exchequer. He could do it, in fact, only by imposing the charges on drugs, appliances, and dental treatment authorized by last year's National Health Service Act. But no sooner had his Health Service estimates been published in full than the Danckwerts award to general practitioners knocked the limit for six, mainly because of the back payments involved. This year's estimates are freed from this complication of back payments, but the effect of the award on the limit is clearly seen in the following table :

*Health Service Estimates**

Total Net Cost to Taxpayer	£m.			
	1951-2 398	1952-3		1953-4 410
		Original 392	Revised 429	
Grants to local health authorities	18.6	20.6	21.2	21.6
General-practitioner service	48.4	48.5	86.9	58.2
Pharmaceutical service†	43.6	39.9	40.9	45.0
Dental service†	36.2	20.1	20.4	22.0
Supplementary ophthalmic service†	14.6	6.3	6.3	6.4
Grants to hospital boards	274.3	295.6	296.0	297.1
Of which maintenance	260.8	285.7	286.1	287.0

* Excluding civil defence. † Allowing for charges to patients.

Thus, compared with the pre-Danckwerts estimate, the total net cost of the Health Service to the taxpayer has risen by £18m. Of this, more than half is accounted for by the higher payments to general practitioners resulting from the award; but there are also small and, in one case at least, ominous increases in the other main items of expenditure.

*Expenditure of Local Health Authorities**

	£m.		
	1951-2	1952-3	1953-4
Health centres	†	†	†
Care of mothers and young children	9.4	9.6	9.5
Midwifery	5.0	5.4	5.1
Health visiting	2.7	3.1	3.2
Home nursing	4.3	5.4	5.5
Vaccination and immunization	0.5	0.6	0.6
Ambulance service†	7.9	9.3	9.7
Prevention of illness and aftercare	1.7	2.1	2.2
Home helps	3.5	4.3	4.7
Mental health	1.8	2.1	2.2
Miscellaneous	0.2	0.3	0.3
	37.0	42.2	43.0

* Including estimated expenditure from rates.

† Less than £100,000 in England and Wales.

‡ England and Wales only.

Expenditure on the services of the local health authorities has virtually been stabilized. Indeed, it is impossible to see how it could be otherwise. The initiative in this branch of expenditure comes from the local authorities, and about half the total cost falls on the ratepayer. Already local authorities have had to raise their rates for the coming year—increases of as much as 2s. in the pound are common. They may well hesitate before expanding further any of their health services. And the argument that by so doing they might relieve the hospitals, and therefore the taxpayer, will, not unnaturally, carry no weight. Estimated expenditure on health centres in England and Wales is falling from £94,000 to £50,000, which must mean a virtual cessation of new centres. In Scotland, on the other hand, where the cost of health centres is borne on the executive councils' vote, although the original