hospital treatment. In February, 1952, pulmonary tuberculosis was diagnosed following a two-months history of cough, sputum, and loss of weight. She was admitted to the diabetic unit at Romsley Hill Sanatorium on March 28. Her general condition was good, she was afebrile, her E.S.R. was 70 mm. in one hour (Westergren method), her sputum was tubercle-positive, and her chest x-ray film showed acute disease confined to the right upper lobe with cavitation in the right upper zone.

Her diabetes before admission had been treated with unrestricted diet and soluble insulin (S.I.) before each meal (36 units t.d.s.). Blood sugars ranged between 172 and 354 mg. per 100 ml. and glycosuria was persistent. She was stabilized on a 2,500-calorie diet and 40 units of S.I. together with 30 units of protamine zinc insulin (P.Z.I.) in the mornings, and 30 units of S.I. in the evenings. Her tuberculosis was treated with rest in bed and a three-months course of isoniazid (200 mg. daily) from April 15 to July 15 with considerable clinical and radiological improvement. Symptoms subsided, she gained 20 lb. (9.1 kg.) in weight, the E.S.R. dropped to 16 mm. in one hour, and her chest x-ray film showed considerable clearing of the disease in the right upper lobe, but there was residual cavitation for which surgery will be required. Her insulin dosage had to be raised to 44 units of S.I. and 40 of P.Z.I. in the mornings and 30 units of S.I. in the evenings in order to maintain satisfactory control of her diabetes during the period of isoniazid administration. Three weeks after isoniazid was discontinued she had her first insulin reaction and there was less glycosuria throughout the day. She developed a second insulin reaction two weeks later, and the dosage had to be reduced to 34 units of S.I. and 40 units of P.Z.I. in the mornings and 30 units of S.I. in the evenings. Control of the diabetes was more difficult to achieve during the period of isoniazid administration, and insulin dosage had to be raised. Three weeks after isoniazid was discontinued control of the diabetes was again more satisfactory, and the development of insulin reactions necessitated reduction of the insulin dosage.

Case 2

A woman aged 36 enjoyed good health until February, 1950, when diabetes was diagnosed. This was treated with 15 units of globin insulin daily. Following a haemoptysis on January 8, 1952, pulmonary tuberculosis was diagnosed, and she was admitted to the diabetic unit at Romsley Hill Sanatorium on January 19. Her general condition was fair; she was febrile (temperature 100.6° F.—38.1° C.), her E.S.R. was 46 mm. in one hour (Westergren method), her sputum was tubercle-positive, and her chest x-ray film showed extensive bilateral disease with cavitation in both lungs. Her diabetes was poorly controlled and she was stabilized on a mixed dose of 24 units of S.I. and 30 units of P.Z.I. in the mornings and 10 units of S.I. in the evenings, with a 2,500-calorie diet. Her tuberculosis was treated with a three-months' course of streptomycin (1 g. daily) and P.A.S. (20 g. daily) from January 20 to April 20, with moderate clinical and radiological improvement. Symptoms subsided, she became afebrile, the E.S.R. dropped to 15 mm. in one hour, and there was moderate clearing of the disease in both lungs. A three-months course of isoniazid (200 mg. daily) was begun on June 30, and glycosuria became more pronounced and fasting blood sugars were in the region of 200 mg. per 100 ml., so that insulin dosage had to be increased to 32 units of S.I. and 32 units of P.Z.I. in the mornings and 16 units of S.I. in the evenings. So far she has had isoniazid for two months, with some clinical and radiological improvement, but her diabetes has been more difficult to control and her insulin requirements have increased.

Summary

After isoniazid the oral glucose-tolerance test shows raised blood-sugar levels in both diabetic and non-diabetic tuberculous patients.

The insulin requirements of diabetic patients receiving isoniazid may be increased.

We thank Dr. T. Russell Fraser, Department of Medicine, Postgraduate Medical School of London, for valuable advice and suggestions; Dr. W. Whitelaw, pathologist to Dudley Road Hospital, in whose department most of the laboratory investigations were done; Mr. R. W. Richardson for help in the biochemical studies; Dr. H. Bickel for the sugar chromatograms; Mr. Peter Hilton for statistical advice; Mr. L. Barratt for the diagrams; Mrs. B. I. Roe for her assistance with the secretarial work; and the patients who co-operated in this investigation.

REFERENCES

- American Trudeau Society (1952). *Amer. Rev. Tuberc.*, **65**, 649.
 Benson, W. M., Steffko, P. L., and Roe, M. D. (1952). *Ibid.*, **65**, 376.
 Bueding, E., Stein, M. H., and Wortis, H. (1941). *J. biol. Chem.*, **140**, 697.
 Dunn, J. S., Sheehan, H. L., and McLetchie, N. G. B. (1943). *Lancet*, **1**, 484.
 Elmendorf, D. F., Cawthon, W. U., Muschenheim, C., and McDermott, W. (1952). *Amer. Rev. Tuberc.*, **65**, 429.
 Elsom, K. O., Lukens, F. D. W., Montgomery, E. H., and Jonas, L. (1940). *J. clin. Invest.*, **19**, 153.
 Izume, S., and Lewis, H. B. (1926). *J. biol. Chem.*, **71**, 51.
 Koref, O., Vargas, L., jun., Rodriguez, F. H., and Telchi, A. (1944). *Endocrinology*, **35**, 391.
 Lepkovsky, S., Wood, C., and Evans, H. M. (1930). *J. biol. Chem.*, **87**, 239.
 Medical Research Council (1952). *British Medical Journal*, **2**, 736.
 Mueller, A., and Fox, S. H. (1951). *J. Amer. Pharm. Ass. (Sci. ed.)*, **40**, 513.
 Pachman, D. J. (1941). *Amer. J. Physiol.*, **133**, 43.
 Preziosi, P. (1952). *Rif. Med.*, **66**, 706.
 Robitzek, E. H., and Selikoff, I. J. (1952). *Amer. Rev. Tuberc.*, **65**, 402.
 Rubin, B., Lee Hassert, G., Thomas, B. G. H., and Burke, J. C. (1952). *Ibid.*, **65**, 392.
 Selikoff, I. J., and Robitzek, E. H. (1952). *Dis. Chest.*, **21**, 385.
 Solimann, T. (1948). *A Manual of Pharmacology*, 7th ed., p. 241. Saunders, Philadelphia

SURGICAL TREATMENT OF INTRACTABLE PHANTOM-LIMB PAIN

BY

MURRAY A. FALCONER, M.Ch., F.R.C.S.

Director, Guy's-Maudsley Neurosurgical Unit

Painful sensations in phantom limbs are common, but they vary in pattern and probably also in their mode of production. Indeed, Riddoch (1941) says that about half the people who undergo major amputations feel pain in their phantoms at some period or other. There are, for instance, the bothersome pains in the stump and phantom associated with tender and irritable stump neuromas, the pains felt in the missing limb at the site of wounding, and other fugitive pre-amputation pains. Many of these pains are transient, and if they do not disappear spontaneously are benefited by simple measures, such as resection of painful neuromas (Bailey and Moersch, 1941; Riddoch, 1941; Henderson and Smyth, 1948), procaine block of the sympathetic chain (Livingston, 1938), or even repeated percussion of stump neuromas (Russell, 1949; Russell and Spalding, 1950). But in a small minority, possibly less than 1% of amputees, the phantom pain is intense and intractable, has features which resemble causalgia, and is not relieved by these measures. It is this type of case that concerns us here.

In 1945 Falconer and Lindsay reported two cases of severe pain in a phantom upper limb which had responded to a high cervical chordotomy, after previous treatments, including resection of neuromas, sympathetic-chain block, and brachial-plexus block, had failed to give relief. Subsequently my experience has increased to 12 patients, all referred to me at the Dunedin Neurosurgical Unit in New Zealand between 1944 and 1949, and all submitted to antero-lateral chordotomy after

simpler measures had proved unsuccessful. These patients have now been followed up for periods of between two and seven years, thereby affording an indication of the permanence of the procedure. Nine patients have been greatly benefited as regards their pain, one is only slightly improved, while in two pain has recurred. Some of those benefited, however, have since developed sequelae of the operation which must be weighed against its benefits.

Phantom Upper Limbs

Six patients had upper-limb phantoms, and their salient clinical features are listed in Table I. Five were amputees, including two Service men of the recent war (Cases 1 and 2),

TABLE I.—*Painful Phantom Upper Limbs Treated by High Cervical Chordotomy*

Case No.	Sex and Age	Reason for Amputation	Site of Amputation	Duration of Pain (Years)	Post-operative Follow-up (Years)	Present State
1	M 29	Gun-shot wound	Upper arm	2	7	Phantom pain relieved
2	M 33	"	"	3	7	Phantom pain relieved, but chronic cervical subluxation present
3	M 48	Gangrene following fractures	Elbow	12	5	Phantom pain relieved, but chronic lumbar-disk syndrome present
4	F 44	Gangrene due to vascular lesion	Forearm	6	3	Phantom pain slightly improved
5	M 58	No amputation; brachial-plexus lesion		20, slightly; 1½, severely	2	Phantom pain relieved
6	M 38	Crushing injury	Forearm		½	Pain recurred within one month. Later to pectomy. Suicide

while the remaining patient had sustained a complete tear of the brachial plexus. Each described his pain as of burning or aching quality, located mainly in the thumb, fingers, and occasionally also the wrist of the missing hand, and associated with feelings of involuntary clenching or squirming movements of the fingers. In Cases 1, 2, and 6 pain had been present right from the time of amputation, while in Cases 3 and 4 its appearance had been delayed for six and four years respectively. The patient with the brachial-plexus tear had experienced mild pain for over 20 years, dating back to the time of injury, but pain had been distressing only for the last one and a half years. None of the amputees had any serious pain in his stump, but one suffered from frequent jactitations of the stump (Case 4).

Each patient had previously undergone other operative procedures with little or no benefit. All the amputees had had their stumps trimmed and their neuromas resected. Cases 4 and 5 had undergone some form of sympathectomy. A diagnostic sympathetic block was performed in the remaining four patients, but with beneficial effect in only two of them. In Case 3, in which the block did seem to afford relief, I subsequently carried out a Smithwick sympathectomy, but the patient's pain recurred within a month. Each patient was also tested by a brachial-plexus block, and in four there was no relief of pain. Case 4, in which the test seemed to benefit, subsequently underwent a posterior rhizotomy of C 4-T 1 inclusive, but the pain returned within three weeks. The two patients seen most recently (Cases 5 and 6) were also treated by repeated percussion of neuromas, but with only partial and fleeting benefit.

All six patients were then submitted to a high cervical chordotomy performed under local analgesia, so that the sensory loss resulting from any incision in the cord could be checked immediately. In each patient the antero-lateral

column of the cord on the side opposite to the stump was incised between the C 1 and C 2 roots. The incision was made in stages, the initial cuts usually numbing only the lower limb and the lower part of the trunk. Thereafter the incision was deepened and extended until pain was relieved in the phantom. We found this did not occur until the level of analgesia had risen to include the stump. By this time we usually found that the antero-lateral column had been incised to a depth of 5 to 6 mm. at a point immediately in front of the ligamentum denticulatum, and that the

FIG. 1.—Diagram illustrating extent of incision usually necessary in high cervical chordotomy.

incision had been carried forwards to emerge on the surface about 3 mm. anterior to the line of the anterior nerve roots (Fig. 1). Such a depth of incision correlates well with Walker's (1940) observations on the position and extent of the pain and temperature fibres in the cervical cord (Fig. 2).

When tested during the convalescent period each patient had a complete loss of pain and temperature sensibility below the C 4 and C 5 segmental levels. Touch was spared, and there was usually a contralateral Horner's syndrome. Often the patient remarked that he felt he had regained some control over the movements of the phantom digits. As regards the later results, continued relief of pain has been noted in four patients, but in Cases 4 and 6 there has been some return of pain. Case 4, however, is no longer so disturbed as formerly. In Case 6 severe phantom pain returned within three to four months of operation, and subsequently was eased only slightly by a topectomy; ultimately this patient took his own life.

However, the four patients whose phantom pains were permanently relieved have each developed other symptoms, although on balance each is better than he was before operation. The first patient remained free of complaints for six years, but wrote recently saying that his stump is now "very sensitive and objects strongly to any pressure," although he has not yet mentioned this to his doctor. When I personally examined him five years after operation the loss of pain sensibility was still complete, whereas in the re-

FIG. 2.—Arrangement of the spinothalamic and other tracts in the lower cervical region of the spinal cord, according to A. Earl Walker (1940). The heavy dots represent fibres concerned with temperature; medium dots, fibres mediating pain; and fine dots, fibres carrying touch and pressure impulses. Note the lamination of the tracts. S represents fibres conveying impulses from the sacral segments, L lumbar, T thoracic, and C cervical.

maintaining three patients some return of pain sensibility in the leg and trunk was noted after a year, and continued or rapidly repetitive painful stimuli applied to these parts gave rise to irritative and burning sensations. Two patients developed even more serious complications. One of them, about seven months after operation, returned to a clerical occupation which involved much flexion of his neck, and shortly afterwards developed a chronic subluxation of C2 vertebra on C3, and of C3 vertebra on C4. The other, several months later, developed a chronic fourth-lumbar-disk syndrome which did not respond to prolonged conservative treatment, and in the end was treated by operation with some improvement. It is possible that this last complication was provoked by a lumbar puncture.

Phantom Lower Limbs

Six of my patients had lower-limb phantoms, and their particulars are set out in Table II. All were amputees, three of them being Service men (Case 7, Norwegian seaman ;

TABLE II.—*Painful Phantom Lower Limbs Treated by High Thoracic Chordotomy*

Case No.	Sex and Age	Reason for Amputation	Site of Amputation	Duration of Pain (Years)	Post-operative Follow-up (Years)	Present State
7	M 36	Infected G.S.W.	Mid-thigh	3½	3	Phantom pain relieved
8	M 56	Infected compound fracture	"	7	5	"
9	M 29	" "	"	4	3	Pain relieved, but paraesthesiae in foot
10	F 33	Suppurative arthritis of knee	Lower thigh	3	3	Phantom pain relieved
11	M 68	Infected compound fracture	Mid-leg	8	2	Pain recurred within two months
12	M 61	Infected G.S.W.	Upper thigh	28, intermittently ; 2, severely	2	Phantom pain relieved

Case 9, 1939–45 war ; Case 12, 1914–18 war). Each described his pain as either burning or aching, and situated in the toes and forefoot of the missing limb, and perhaps also in the heel and ankle. In three the phantom digits were either squirming or rigidly flexed, but in the others the motor state of the missing limbs was not recorded. Two of the patients had tender hyperaesthetic stumps, and three showed marked jactitation of the stump. In four patients the pain had appeared immediately after a surgical operation, which in two was the primary amputation (Cases 8 and 11), in the third was a reamputation occasioned by failure of the first amputation to heal (Case 7), and in the fourth was a resection of stump neuromas (Case 9). In Cases 10 and 12 pain was delayed in appearance for two months and one month respectively.

Each patient had previously undergone a resection of neuromas, and Case 7 had even had his limb reamputated three times at successively higher levels. Only Case 12 was benefited by these procedures, and in this particular patient resection of neuromas had been performed three times in the course of 28 years, on each occasion conferring benefit for from three to ten years. The neuromas were no longer accessible and percussion treatment failed to benefit him. Each patient was also treated by a lumbar sympathetic block and by amylocaine hydrochloride spinal analgesia. In each instance the sympathetic block subjectively warmed the phantom, but in only Case 10 did it relieve the pain, and then only temporarily. In Cases 8, 10, and 11 amylocaine was reported as easing pain, but in the others it had no effect.

All six patients were submitted to a high thoracic chordotomy performed under general anaesthesia at the level of the T2 and T3 vertebra. In each patient the antero-lateral

column on the side opposite the stump was incised to a depth of 3 to 4 mm. immediately in front of the ligamentum denticulatum, and the incision carried forwards to emerge on the surface 2 to 3 mm. in front of the line of anterior nerve roots. In my experience there is no need to perform thoracic chordotomy under local analgesia, as such an incision constantly produces a satisfactory degree and extent of analgesia.

Following operation, in each patient loss of pain and temperature sensibility was apparent below the T5 to T7 segmental levels on the side of the stump, while the sense of pain in the phantom disappeared, although, as with the upper limbs, the phantom itself remained. In Case 11, however, pain returned within two months, while in Case 3 a continuous feeling of pins-and-needles appeared after six months in the phantom foot. In the remaining four patients the relief of pain has continued for from two to five years. One of these (Case 7) when last heard of had epigastric pain suggestive of a peptic ulcer ; this presumably was unrelated to chordotomy. Case 10 had undergone an amputation of her second leg, also for chronic suppurative arthritis, but 18 months later had not developed pain in the new phantom.

Discussion

Thus antero-lateral chordotomy would appear to offer prolonged, if not permanent, relief from severe pain in a high proportion of phantom limbs, both upper and lower, but not in all. It is interesting to note how it compares with other methods of treatment, and what light it throws on the mechanisms by which phantom phenomena are produced.

Phantom-limb pain of the type considered in this report is rare, occurring in possibly less than 1% of cases. Ewalt, Randall, and Morris (1947) found only eight such cases in a group of 2,284 American Service men, and Henderson and Smyth (1948) only two or three among some 300 amputees in a prisoner-of-war camp. My own experience is probably similar. I cannot tell what proportion of civilian amputees were represented by my patients, but three of them were New Zealand Service men injured in the recent war, and, including these, the New Zealand War Pensions Office has knowledge of only four or five such cases occurring in a total of 496 surviving amputees (D. Macdonald Wilson, 1951, personal communication).

The rarity of the condition and the fact that the sufferers often show evidence of a disturbed personality have led some clinicians to regard the pain as largely a psychological disturbance (Bailey and Moersch, 1941 ; Ewalt, Randall, and Morris, 1947 ; Henderson and Smyth, 1948). Guillaume, Froideveaux, and Mazars (1947) and Kolb (1950) have even described cases which were benefited by psychotherapy, while Pisetsky (1946) reported a case cured by electric convulsion therapy. Certainly some of my patients were greatly demoralized by pain, and were perhaps unstable individuals, but so are many sufferers from other organic painful conditions, such as trigeminal neuralgia. Most of my patients had previously been seen by psychiatric colleagues, who reported that they could not find any significant psychogenic features, neither could they relieve these patients by psychotherapy. I find it difficult to believe that in my hands antero-lateral chordotomy acted as a psychotherapeutic procedure, when previous operative procedures on the stump had failed to give relief. Instead, I am impressed by the fact that these patients all described their pain in much the same terms, locating the pain mainly in the missing digits and associating it with clenching or squirming of those digits. As their symptoms thus conform to a fairly distinctive pattern, and as there was no possibility of collusion between individual patients, I feel that such painful phantom-limb syndromes must be regarded as organically determined.

Comparison of Various Therapeutic Procedures

It is generally agreed that pain of the severity and intractability shown by my patients is not relieved by any destructive procedure which involves the primary sensory

neurone, such as reamputation of the limb, resection of neuromas, section of nerve trunks or plexuses, or posterior rhizotomy (Leriche, 1939; Riddoch, 1941; White, 1950); neither is it relieved by sympathectomy (White, 1950). My own experiences bear out this view. However, this type of pain is often relieved by such procedures as percussion of stump neuromas, antero-lateral chordotomy, posterior chordotomy, ablation of the post-central gyrus, and, finally, leucotomy, but the relief afforded is not always of the same quality.

Repeated Percussion of Stump Neuromas

Repeated percussion of stump neuromas either with a mallet or with a vibrator is the basis of a simple but often effective treatment introduced by Russell (1949). This procedure generally gives relief from pain for only a few hours at a time, so that repeated and self-administered treatments are necessary, the phantom persisting during these periods of relief. This treatment seems to be most effective for cases with tender neuromas of the type that would be eased by nerve-blocks, but occasionally it will numb the pain of phantom lower limbs even when the pain has not been influenced by deep spinal analgesia (Russell, 1952, personal communication). The treatment, however, is not always effective, and Russell and Spalding (1950) mention that in one group of 18 cases there were eight failures. As most of my cases were operated on before the advent of Russell's papers, I tried his method in only three patients: in Cases 5 and 12 it was unsuccessful, possibly because the neuromas were inaccessible, and in Case 6 it produced some relaxation of pain and posture in the clenched phantom hand, even though a plexus block had been completely ineffective, but the degree of relief it afforded did not satisfy the patient.

The mechanism of pain relief underlying these observations is not certain. Russell felt that by maintaining the neuromas in a "chronically concussed state" he rendered them insensitive, while the repeated trauma caused the nerve endings to degenerate gradually and be replaced by fibrous tissue. This explanation is possibly faulty, for more certain ways of cutting off abnormal impulses arising in neuromas, such as procaine blocks or even section of the nerve, consistently fail to give relief in the type of case under consideration. Rather, I think, the explanation lies in the fact that by traumatizing the neuroma a barrage of afferent impulses is sent up to the spinal cord, as is postulated in the hypothesis which I shall later develop. Bartlett (1951) has already put forward this explanation, which possibly also accounts for the observations of Bailey and Moersch (1941) that slight pressure on the nerve trunk above the amputation may cause the phantom to disappear, whereas section of the nerve does not.

Antero-lateral Chordotomy

Although Bailey and Moersch (1941), de Guitérrez-Mahoney (1944), and Coleman (1946) were unsuccessful with this method, Riddoch (1941) and White (1950) had experiences similar to mine, while Bromley (1930) reported a successful outcome in a painful phantom of the upper limb which was treated by high cervical chordotomy. The general finding has been that following this operation the pain in the phantom disappears, although the phantom itself persists. This observation, taken in conjunction with the observation that posterior rhizotomy has no effect on such pain, suggests that in the type of case we are considering the structures responsible for the pain are the secondary sensory neurones which have their cell bodies in the substantia gelatinosa of Rolando. In the very few cases in which the operation fails, neurones at still higher levels in the pain-pathways of the central nervous system are presumably perpetuating the pain.

Posterior Chordotomy

Pool (1946) and also Browder and Gallagher (1946, 1948), working independently, investigated the effect of section of the posterior column of the spinal cord. Their hypothesis was that, if the proprioceptive pathways from the whole limb were severed, the cramped posture of the phantom limb

might be relieved, and consequently the pain in it, and even the phantom itself might disappear. Pool operated on three cases, all phantoms of the upper limb, dividing the lateral part of the posterior column at the level of the fifth cervical lamina. In at least one patient, operated on under local analgesia, he found that pain was not relieved until the incision had been carried laterally across the line of the posterior nerve roots, and so into the substantia gelatinosa of Rolando. Post-operatively all three patients seemed benefited, the phantom hand being free of pain, but still retaining its sense of position. In each patient diminished sensibility to pinprick associated with an exaggerated response to temperature was observed over the upper thoracic segments. Possibly the explanation of these sensory changes in both the thorax and the phantom lay in the involvement of the secondary sensory neurones of the substantia gelatinosa. The benefit seen in Pool's cases, however, was not permanent (Pool, 1952, personal communication).

Browder and Gallagher operated on six cases, comprising four lower-limb phantoms and two upper-limb phantoms. For the lower-limb cases they divided the whole of the dorsal column on the appropriate side at the mid-thoracic level. For the upper limb they operated at the C2 level, sectioning the lateral two-thirds of the dorsal column without disturbing gait or stance. After dorsal chordotomy, complete loss of the phantom foot as well as the cramp-like pain occurred in three patients; in the other pain was relieved, but the phantom persisted. In none of these four patients was there any disturbance of motor power in the stump or of touch, although vibration sensibility in the stump was diminished in three. The results in the two cases of phantom upper limb were difficult to evaluate, and neither case could really be regarded as successful. Browder (1952, personal communication), however, said that the benefit observed in the lower-limb cases has been permanent.

The rationale of these operations is difficult to understand, for the fibres in the posterior column are the proximal axons of cells in the dorsal root ganglia, and yet posterior rhizotomy which severs the same axons but at another site does not relieve pain.

Ablation of the Post-central Gyrus

In 1944 de Guitérrez-Mahoney reported having successfully relieved pain in a phantom upper limb by excision of the appropriate area of the post-central gyrus of the opposite cerebral hemisphere. Since then other surgeons, also operating under local analgesia, have confirmed that stimulation of the appropriate area with an electrical current prior to excision will intensify the pain, while resection of the area will abolish the awareness of the phantom in addition to the pain. There is, however, some doubt about the permanence of pain-relief after this operation. Thus Echols and Colclough (1947) and Stone (1950), who, like de Guitérrez-Mahoney in his first paper, reported their experiences shortly after operation, claimed good results, but Horrax (1946, 1948) and Sunderland and Kelly (1948) stated that in their cases pain has usually returned within a few weeks or months. Evidently this has been de Guitérrez-Mahoney's own experience, for in 1948 he stated that in three out of his four personal cases there had been some return of pain after several weeks.

The fact that stimulation of the post-central gyrus intensifies the pain while excision of the gyrus destroys the phantom as well as the pain suggests that the sensations connected with the phantom enter into consciousness, at least partly, in this area. In discussing his long-term findings, de Guitérrez-Mahoney drew attention to the second sensory cortical area described by Adrian (1940), by Woolsey and Fairman (1946), and by Penfield and Rasmussen (1950), which is situated along the upper bank of the Sylvian fissure in the parietal region, and which receives sensory impulses from both sides of the body. It would seem possible that the return of pain some time after post-central gyrectomy may be due to increased functioning of this second sensory area.

Frontal Lobotomy

Several surgeons, notably Van Wagenen (1944), Watts and Freeman (1948), and Waugh, Cameron, Howarth, and Matas (1949), have reported successful results in painful phantoms after prefrontal leucotomy, while Le Beau (1950) has observed benefit following topectomy. The relief produced by these procedures is presumably similar to that observed following these same operations for pain due to malignant disease and other organic pain-states, in that the pain persists but no longer worries the patient as formerly. Some degree of emotional blunting is probably inseparable from a beneficial response. Therefore such operations should not be tried or contemplated for painful phantom limbs until other measures, such as percussion of neuromas and antero-lateral chordotomy, have been tried and have failed.

Mechanism of Pain-production

Weir Mitchell (1872), in his classical paper, postulated that the phenomena of phantom limbs were due to an "ascending neuritis" spreading up from the severed ends of peripheral nerves. No one, however, has ever demonstrated the histological changes of inflammation within the nerves, but variations of the theory substituting local irritation of nerve fibres within the neuromas or at the cut ends of nerves can readily be advanced to explain why some painful phantoms, principally those of milder degree, are eased by resection of stump neuromas. The "peripheral" theory, however, will not explain why phantom phenomena persist when the peripheral nerve is blocked or sectioned, nor will it explain the method of pain relief in the intractable cases which have here been considered. Hence some have argued that the mechanism must be a central one, situated in the cortex (Henderson and Smyth, 1948).

Certainly, as I have already mentioned, the evidence afforded by stimulation of the post-central gyrus in conscious patients and the effects of its excision suggests that in some way phantom sensations enter into consciousness in this cortical area. But these sensations may still arise from the activity of nerve cells intermediate between the cortex and the peripheral neurone. In an earlier paper (Falconer and Lindsay, 1945) it was pointed out that, in cases with intractable pain, antero-lateral chordotomy renders a phantom painless, whereas posterior rhizotomy does not, and consequently that the neurones presumably responsible for the pain are the cells of the substantia gelatinosa of Rolando. I would like to develop this hypothesis further in the light of more recent evidence, both theoretical and experimental.

When a peripheral nerve is sectioned the stream of afferent impulses, which used to pass up it continuously to the spinal cord, either ceases completely or is grossly diminished and distorted. The observations of Weddell, Sinclair, and Feindel (1948) on the anatomical basis underlying pain in painful scars are germane in this connexion. They compared the distribution of nerve fibres in painful and painless scars with that in normal skin, and found that in painful scars the nerve endings were arranged in isolated fashion, and not interwoven as in normal skin and in sensitive but painless scars. They argued that the pattern of sensory impulses relayed to the spinal cord from these painful areas was probably an incomplete one, and that it was this distortion of the pattern rather than the diminution in the total number of impulses which resulted in the sensorium discerning the diffuse and aching pain that is characteristic of hyperpathia. Possibly the same principles can be used to explain painful phantoms, for, because of the section of several peripheral nerves inevitable in an amputation, a distorted as well as a diminished pattern of impulses must reach the cells of the substantia gelatinosa, and the harmonics of this pattern may be such that unusual painful impulses are evoked. If the stump neuromas are then percussed, a barrage of afferent impulses is conveyed up the hitherto relatively silent peripheral nerve, and perhaps restores both the pattern and the intensity of the afferent impulses to more normal limits.

Eccles and McIntyre (1951) have described experiments which may help to explain the actual changes that may occur in the synapses around the cells of the substantia gelatinosa. In their experiments in cats they sectioned several posterior nerve roots serving one hind limb, dividing the roots just distal to their ganglia, and then allowed the animals to recover. Several weeks later they stimulated these same posterior roots singly at sites proximal to the ganglia, and observed the changes which had taken place in the mono-synaptic (two-neurone) reflex arcs that form the basis of the ankle-jerk and other simple myotatic stretch-reflexes. To control their work they stimulated the nerve roots adjacent to the operated area, as well as the nerve roots on the unoperated side. They recorded several important findings. First, the operative section greatly impaired the reflex effectiveness of stimulating the dorsal root. For instance, a single momentary electrical stimulus applied to the root would no longer evoke a contraction in the muscle. Secondly, if the reflex arc was first conditioned by a tetanizing stimulus—for example, 6,000 volleys at 400 volleys a second—the muscle would respond to a single momentary stimulus (single volley) given even several minutes after the tetanizing stimulus had ceased. The influence exerted by this tetanizing stimulus lasted longer on the operated than on the control side, although the muscle response was less on the operated side. Thirdly, they found that if the dorsal root immediately rostral to those operatively divided was stimulated, its reflex effectiveness was greater than that of its companion on the unoperated side. Eccles and McIntyre thought that these observations indicated that usage leads to increased functional efficiency of synapses and disuse to defective function, and also that, when habitual afferent impulses are cut off, stimulation along other pathways may sometimes result in increased potentiation of motor neurones. It seems not unreasonable to apply these observations to the state of the cells of the substantia gelatinosa following an amputation by conceiving that when these cells are cut off from their normal pattern of afferent impulses they may become stimulated by impulses from other pathways. Further, percussion of the stump neuromas may act like a tetanizing stimulus and help to condition the nerve cells to react in a more normal manner.

I therefore postulate that the explanation of severe pain in phantom limbs, of the type here surveyed, is that the cells of the substantia gelatinosa, cut off from their normal-patterned stream of stimuli, discharge spontaneously or react to unusual sources of stimuli, and so provoke pain. This explanation would account for many features of the syndrome, and is in keeping with the neuro-anatomical and neuro-physiological observations just considered.

Summary

Experiences gained in treating 12 patients with intractable phantom-limb pain are recorded, six of whom had involvement of an upper limb and six of a lower limb. Each patient had suffered severe pain for some considerable time, mostly a matter of years, and many of them had previously been treated by such measures as resection of stump neuromas, sympathectomy, or re-amputation without benefit. Each patient was then submitted to an antero-lateral chordotomy at an appropriate level and subsequently a pronounced relief of pain was observed in nine of them, slight benefit in one, and a return of pain in the remaining two.

Pain of the type and severity shown by these patients is rare after amputation. Several other measures, including percussion of stump neuromas, antero-lateral chordotomy, posterior chordotomy, post-central gyrectomy, and leucotomy, may relieve such pain, and their respective merits are discussed.

It is felt that in the type of case described antero-lateral chordotomy gives the best results. The mechanism of pain production and its relief by the various treatments is discussed.

Mr. Anthony James and Dr. D. Macdonald Wilson, both of New Zealand, have given me much assistance in following up my patients. I am also grateful to Professor J. C. Eccles and to Dr. T. McLardy for helpful discussions regarding the mechanism of phantom-limb pain.

REFERENCES

- Adrian, E. D. (1940). *J. Physiol.*, **98**, 16P.
 Bailey, A. A., and Moersch, F. P. (1941). *Canad. med. Ass. J.*, **45**, 37.
 Bartlett, J. E. A. (1951). *Brain*, **74**, 363.
 Bromley, L. (1930). *Guy's Hosp. Rep.*, **80**, 234.
 Browder, J., and Gallagher, J. P. (1946). *N.Y. St. J. Med.*, **46**, 2403.
 — (1948). *Ann. Surg.*, **128**, 456.
 Coleman, C. C. (1946). In *Surgical Treatment of the Nervous System*, edited by F. W. Bancroft and C. Pilcher, p. 449. Philadelphia.
 Eccles, J. C., and McIntyre, A. K. (1951). *Nature, Lond.*, **167**, 466.
 Echols, D. H., and Colclough, J. A. (1947). *J. Amer. med. Ass.*, **134**, 1476.
 Ewalt, J. R., Randall, G. C., and Morris, H. (1947). *Psychosom. Med.*, **9**, 118.
 Falconer, M. A., and Lindsay, J. S. B. (1945). *Brit. J. Surg.*, **33**, 301.
 Guillaume, J., Froideveaux, and Mazars (1947). *Rev. neurol., Paris*, **79**, 213.
 de Guiterrez-Mahoney, C. G. (1944). *J. Neurosurg.*, **1**, 156.
 — (1948). *Surg. Clin. N. Amer.*, **28**, 481.
 Henderson, W. R., and Smyth, G. E. (1948). *J. Neurol. Neurosurg. Psychiat.*, **11**, 88.
 Horrax, G. (1946). *Surgery*, **20**, 593.
 — (1948). *Ann. Surg.*, **128**, 469.
 Kolb, L. C. (1950). *Proc. Mayo Clin.*, **25**, 467.
 Le Beau, J. (1950). *J. Neurosurg.*, **7**, 79.
 Leriche, R. (1939). *The Surgery of Pain*, translated by A. Young. London.
 Livingston, W. K. (1938). *Arch. Surg.*, **37**, 353.
 Mitchell, S. W. (1872). *Injuries of Nerves and Their Consequences*. Philadelphia.
 Penfield, W., and Rasmussen, T. (1950). *The Cerebral Cortex of Man*. New York.
 Pisetsky, J. E. (1946). *Amer. J. Psychiat.*, **102**, 599.
 Pool, J. L. (1946). *Ann. Surg.*, **124**, 386.
 Riddoch, G. (1941). *Brain*, **64**, 197.
 Russell, W. R. (1949). *British Medical Journal*, **1**, 1024.
 — and Spalding, J. M. K. (1950). *Ibid.*, **2**, 68.
 Stone, T. T. (1950). *Arch. Neurol. Psychiat.*, Chicago, **68**, 739.
 Sunderland, S., and Kelly, M. (1948). *Aust. N.Z. J. Surg.*, **18**, 75.
 Van Wagenen, W. P. (1944). Quoted by J. C. White, *J. Amer. med. Ass.*, **124**, 1030.
 Walker, A. E. (1940). *Arch. Neurol. Psychiat.*, Chicago, **43**, 284.
 Watts, J. W., and Freeman, W. (1948). *Proc. Ass. Res. nerv. ment. Dis.*, **27**, 715.
 Waugh, O. S., Cameron, H. F., Howarth, J. C., and Matas, J. (1949). *Canad. med. Ass. J.*, **61**, 364.
 Weddell, G., Sinclair, D. C., and Feindel, W. H. (1948). *J. Neurophysiol.*, **11**, 99.
 White, J. C. (1950). *Lancet*, **2**, 161.
 Woolsey, C. N., and Fairman, D. (1946). *Surgery*, **19**, 684.

The annual report on the medical services of Nigeria for 1950-1 contains an account of a scheme to eradicate mosquitoes at Ilaro in Southern Nigeria. The scheme was based on the suggestions of the Expert Committee on Malaria of the World Health Organization, and the preliminary malaria survey began in 1949. Ilaro has some 12,000 inhabitants and 2,300 houses, and is a typical mixed urban and rural community situated within a geographical zone where hyperendemic malarial conditions exist. *P. falciparum* is the main parasite species but there is also *P. malariae*. *A. gambiae* and *A. funestus* are the main vectors, with *A. funestus* persisting throughout the year, while *A. gambiae* has a pronounced seasonal importance during the rains. Residual spraying began in March, 1950. The insecticide used was the benzene hexachloride wettable powder "P.520," containing 6.5% of the gamma isomer. The spraying of the 2,300 houses, containing over 11,000 rooms, is being repeated every three months and the results are now being assessed by the study of entomological and malarimetric data. According to the report it is still too early to assess the results of this scheme, which must last for three full years before any valid information is obtained. Nevertheless the preliminary results have shown that there has been a spectacular decrease of adults of *A. funestus*. The decrease of malarimetric indices in the local population has been slow, but this was not unexpected in an area previously "saturated" with malaria. The most pronounced reduction of the parasite rate occurred during the last half of 1950 in very young children. There was also a considerable reduction of malaria morbidity recorded at the Ilaro dispensary. The scheme is costing about £3,000 per year, or approximately 5s. per head of the local population.

DETECTION OF TUBERCULOSIS IN GENERAL PRACTICE BY PHOTOFLUOROGRAPHY

BY

F. A. NASH, M.R.C.S., L.R.C.P.

Medical Director, South-west London Mass X-ray Service

B. BENJAMIN, B.Sc., F.I.A.

Statistician

AND

T. J. LEE, M.B., B.S., M.R.C.P.

General Practitioner

Stradling (1948) has shown that the average apparent diagnostic delay in cases of pulmonary tuberculosis referred to his chest clinic at Hammersmith amounted to six months. Procrastination for two months occurred before the patient sought medical advice. The general practitioner waited an average of four months before he referred the patient to the clinic for special chest investigation. For years Toussaint (1946) and others have stressed that, if we are to find our cases earlier, minor respiratory and other symptoms in people who are not ill must be regarded as indications for a chest x-ray examination. Stradling lists the following symptoms or conditions based on Toussaint (1946) as indications for a chest x-ray film: cough, haemoptysis, dyspnoea, laryngitis, lassitude, bronchial catarrh, pneumonia, influenza out of season, recurring febrile chills, P.U.O., pleurisy, pleurodynia, or pain in chest, loss of weight, anorexia, anaemia and debility, amenorrhoea (unexplained), ischio-rectal abscess, erythema nodosum, phlyctenular conjunctivitis, contact.

X-ray Film Requirements

It was decided to ascertain the incidence of these symptoms, to which reflex dyspepsia was added, in a general practice of 3,500 in Wandsworth. In the single month of March, 1949, 155 patients of one of us (T. J. L.) complained of or presented with one or more of the above symptoms. Of the 155, 106 attended the surgery with these symptoms for the first time in March, 1949; 30 had previously attended with the same or allied symptoms in January or February, 1949; and 19 had come with the same or allied symptoms in October to December, 1948. All the 155 patients were referred for photofluorography. Had the scheme been in operation before March, 1949, presumably at least 49 of the 155 patients would already have been x-rayed. Later, over a period of two years, it was found that, using clinical judgment, about 20 films a month were required for the practice of 3,500. *This is just over five films a month per thousand patients* in the practice. The general practitioner ordering the x-ray examination had spent many years in general practice and had also had considerable experience of modern sanatorium treatment. This practice is probably representative of the area, and therefore a provisional estimate of the number of films required to be taken for general practitioners, using the indications previously mentioned, in a chest clinic area with a population of 200,000 would be 1,000 chest x-ray films a month, or 12,000 a year. Hunt (1951) stated that "during the last three years" 624 chests had been x-rayed—apparently in a practice of about 3,800. This gives an average of about 4.5 films per 1,000 patients a month, which is very similar to our figure of 5 per 1,000 a month.

Trenchard and Grenville-Mathers (1952) have said: "Instead of throwing the diagnostic net of the clinics ever