the organism being a normal inhabitant of the intestinal tract of man. The customary brief skin "cleansing" with alcohol would not be sufficient to kill spores. Alternatively, the organism may have been present in any of the injection solutions or apparatus; proof of this is impossible to obtain.

Summary

Myositis due to *Clostridium welchii* infection was found in the region of injections of sodium sulphathiazole at the post-mortem examination of a man who died of acute coronary thrombosis. It is believed that the very alkaline solution of sulphathiazole was responsible for muscle necrosis, and that accidentally introduced infection of a necrotic focus led to gas-gangrene. The possible source of the infection is mentioned, and some simple experiments in the culture of *Cl. welchii* at varying pH values are described.

We are grateful to Professor H. W. C. Vines for valuable advice and to Mr. H. Neville Stafford, coroner for Westminster, for permission to publish.

REFERENCES

- Banks, H. S. (1940). *Proc. roy. Soc. Med.*, 33, 555.
 Gale, E. F. (1941). *Biochem. J.*, 35, 66.
 — and van Heyningen, W. E. (1942). *Ibid.*, 36, 624.
 McIntosh, J., and Selbie, F. R. (1943). *Lancet*, 1, 793.
 Selbie, F. R., and McIntosh, J. (1943). *J. Path. Bact.*, 55, 477.

PETHIDINE AND THE TRIPLE RESPONSE

BY

B. L. FINER, M.B., B.S.

AND

M. W. PARTINGTON, M.B., B.S.

(From the Department of Physiology, London Hospital Medical College)

Zuck (1951) and Pearce (1951) observed that the intravenous injection of pethidine was followed by the appearance of a wheal extending proximally along the line of the vein. Nasmyth and Stewart (1950) observed that the injection of pethidine into human skin produced a triple response which was reduced in extent by antihistamines, and suggested that pethidine was a histamine liberator. We have made similar observations. In view of the widespread use of pethidine and reports of unexpected hypotension and collapse following its administration (Giff, 1947; Butler, 1951; Zuck, 1951), it was decided to investigate the production of this triple response further. The results are reported below.

Methods

Intradermal injections (0.1 ml.) were made into the skin of the forearm with a Mantoux syringe.

The effects of intravenous injections were studied under two different sets of conditions. (1) *Venous congestion*:—A sphygmomanometer cuff was placed round the upper forearm and pumped to 70 mm. Hg. The injection (in a volume less than 0.5 ml.) was made into a small wrist vein. Immediately the needle was removed, pressure was applied with a pad of cotton-wool over the site where the vein had been punctured to

prevent blood leaking out of the vein. The pressure in the cuff was maintained for three to four minutes after the injection and then released. (2) *Circulatory occlusion*:—A sphygmomanometer cuff was placed round the upper forearm and pumped to 70 mm. Hg. This pressure was maintained for one minute to congest the vessels of the forearm and then raised to 200 mm. Hg. One minute later the injection was made into a wrist vein, and after a further five minutes the pressure was released.

The Intravenous Injection of Pethidine and Histamine

A triple response is seen over the veins in the forearm if pethidine (25–100 mg.) is injected into a wrist vein with an unrestricted circulation in the arm. The reaction is more conveniently studied if smaller doses (5–15 mg.) are used with some obstruction of the venous return as described above.

Within 30 seconds of injection under these conditions an intense sensation of itching spreads up the forearm. This is often followed by a change in the colour of the vein to a characteristic mauve or plum colour. At the same time a flare appears in the skin round the site of injection and then on each side of the vein for a distance of up to 3 cm., progressively following the vein up the forearm. Within five minutes of injection a wheal appears, localized at first to a thin strip over the vein, but later spreading a few millimetres into the skin on each side. If the vein divides, the flare and wheal follow the branches. Sometimes an unaffected area of skin is seen between the site of injection and the changes higher up in the forearm. Whealing and flaring may occur distal to the site of injection and occasionally discrete wheals form in the skin. Massive wheals are sometimes seen. The typical appearance following an injection of pethidine under these conditions is shown in Fig. 1.

Within two or three minutes of the intravenous injection of pethidine with the circulation *occluded*, a red reaction is seen in the skin. A number of discrete irregular red areas appear, some distal and some proximal to the site of injection. The areas lie over and on each side of the veins in the forearm. They increase in size slowly and may coalesce. When the circulation is re-established and the reactive hyperaemia has subsided the forearm is seen to be extensively whealed, and the margins of the wheals roughly correspond with the margins of the previously observed red reactions. Surrounding all the whealed areas there is a bright red arteriolar flare. On four occasions after the pressure in the cuff was released, the subject's face was seen to flush, and a characteristic metallic taste and a transitory headache were experienced, indistinguishable from those produced by injections of histamine.

All these changes, except the very large wheals, may be reproduced exactly if histamine (15–30 µg. in terms of the base) is substituted for pethidine in the above experiments.

The lymphatic vessels play no part in this response, as the following experiment shows: if pethidine or histamine

FIG. 1.—Changes in the forearm after 15 mg. of pethidine injected intravenously. Sphygmomanometer cuff (at 70 mm. Hg) at the elbow for four minutes after injection.

is injected intravenously with the veins congested, and immediately after the injection a tourniquet is applied to the forearm just proximal to the site of injection, the changes in the forearm occur in their usual sequence.

These observations can be most readily interpreted by assuming that the vein wall is permeable to both histamine and pethidine.

Effect of "Histamine Liberators" on Skin Over Vein

MacIntosh and Paton (1949) have shown that, although the injection of a histamine liberator compound into the skin is followed by a triple response, a second injection of the same substance into the same site within 24 hours produces either no response or one that is much reduced. They suggested that the original injection had reduced the normal complement of histamine in the skin, and that this had not been restored by the time the second injection was given. We have found that the skin behaves similarly after intradermal injections of pethidine (50 μ g.), which supports the conclusion that pethidine is a histamine liberator.

Pethidine (10 mg.) was injected intravenously while the circulation was occluded and the areas of red reaction were outlined in ink. Twenty-four hours later pethidine (50 μ g.) was injected intradermally into an area of skin which had previously shown a red reaction. The response was grossly reduced compared with an intradermal injection of the same amount into a comparable site on the other forearm. An intradermal injection of histamine had its usual effect. It seems likely, therefore, that the triple response seen after an intravenous injection of pethidine is due to the liberation of histamine from the skin.

Effects of Antihistamines

Treatment with an antihistamine drug modifies the triple response to a subsequent intradermal injection of histamine. Experiments were therefore carried out to investigate the

Effect of Pre-treatment with Diphenhydramine Hydrochloride (100 mg.) on Skin Response in Forearm after Intravenous Injections of Pethidine (100 mg.)

Subject	Effect of Pethidine	
	Without Diphenhydramine Hydrochloride	With Diphenhydramine Hydrochloride
1		0
2	++	++
3		++
4	++	
5		+
6	++	+
7	++	
8	++	
9	++	
10		0
11		+
12	++	
13	++	
14		+
15		+
16	++	

0 = No response.

+ = Mauve vein coloration with or without flare.

++ = Includes whealing of all grades.

Subject 6 was seen twice, with three-weeks interval between observations.

results of (a) the oral administration of an antihistamine drug (diphenhydramine hydrochloride) on the response of the skin to an intravenous injection of pethidine given two hours later; (b) the injection of mepyramine alone; and (c) the intravenous injection of mixtures of mepyramine and pethidine and of mepyramine and histamine.

(a) In a series of 16 cases for operation, it was arranged that half the patients received diphenhydramine hydrochloride (100 mg.) by mouth two hours before the anaesthetic was given. The patients were induced with thiopentone, and then pethidine (100 mg.) was injected into a wrist vein. The reactions in the forearm were noted and graded on a rough quantitative scale (see Table) before the observer knew which patients had received diphen-

hydramine hydrochloride. All except one of the subjects who received this drug showed either no changes in the skin or a much reduced response compared with the controls.

(b) The intradermal injection of mepyramine is followed by the appearance of a red reaction in the skin (Halpern, 1952). This reaction may be distinguished from the red reaction produced by histamine by the following characteristics: (1) the colour is a dull rather than a bright red; (2) when the red area fades there is no "speckling" at its margin; and (3) the intradermal injection of mepyramine into an arm with the circulation occluded does not give rise to a red reaction until the circulation is re-established. When mepyramine (5-10 mg.) is injected intravenously under the conditions of *venous congestion* a red reaction is seen in the forearm which is in every way similar to that described above as following the intradermal injection of mepyramine. This red reaction is distributed along the course of the veins in a manner similar to that of the triple response after injections of pethidine. If mepyramine is injected intravenously when the circulation is *occluded* no red reaction appears until the circulation is re-established. These results may best be explained by assuming that the vein wall is permeable to mepyramine.

(c) The intravenous injection of mixtures of mepyramine (10 mg.) and pethidine (5 mg.) or of mepyramine (10 mg.) and histamine (20 μ g.) under the conditions of *venous congestion* described above produced no whealing in the forearm, but only a red reaction similar in character to that following the injection of mepyramine alone. Further, the intravenous injection of such mixtures when the circulation was occluded produced no red reaction until the circulation was re-established.

Since mepyramine does not prevent the liberation of histamine but modifies its effects by competing with it for tissue receptors (Paton and Schachter, 1951), the inhibition of the triple response in the above experiments supports the conclusion that the vein wall is permeable to mepyramine.

Effects of Pethidine on Cat's Blood Pressure

The effects of pethidine on the cat's blood pressure were found to be substantially the same as those described by Duguid and Heathcote (1940). The fall in blood pressure in cats under chloralose following the intravenous injection of pethidine (3 mg./kg.) lasted from 45 to 90 minutes (Fig. 2). Recovery from the hypotension from pethidine was not hastened by mepyramine (0.8-1.6 mg./kg.), nor did pre-treatment with mepyramine in doses up to 3 mg./kg. modify the blood-pressure response.

It follows that the hypotensive effects of pethidine in the cat cannot be explained solely on the basis of histamine liberation.

FIG. 2.—Cat, 2.9 kg.; under chloralose. Effects of pethidine (3 mg./kg.) on the blood pressure. Time of injection marked with an arrow. B, C, D, and E are successive tracings at 15-minute intervals.

Discussion

Lewis (1927) presented evidence that when a substance, other than histamine, was injected into human skin and produced a triple response, it did so by virtue of the liberation of the histamine already present in the skin but held in an inactive form. The demonstration that a triple response is inhibited by treatment with one of the antihistamine drugs supports Lewis's conclusions. MacIntosh and Paton's (1949) observations in man that the second of two injections of a histamine-liberating compound into the same site within 24 hours fails to produce a full triple response, coupled with the work of Feldberg and Talesnik (1952) on the slow rate of recovery of skin histamine following its liberation in animals, also supports Lewis's hypothesis. It follows that if any substance (1) produces a triple response on intradermal injection in man, (2) produces a reduced triple response when the subject is treated with an antihistamine drug, and (3) fails to produce the same response when a second injection is made into the same site within 24 hours, it is likely that the substance is a histamine liberator.

The present observations show that pethidine fulfils these criteria, and so support the conclusions of Nasmyth and Stewart (1950). They also provide an explanation for the observations of Zuck (1951) and Pearce (1951), and show that the walls of the veins in the forearm are permeable to pethidine, histamine, and mepyramine. No other reference has been found to the passage of substances through the walls of veins of comparable size apart from the observation of Magendie, who in the early part of the nineteenth century applied a "spirituous extract of nux vomica" to the outside of the jugular vein of a dog and observed signs of poisoning within four minutes (Franklin, 1937).

In man, the present observations show that the appearance of a triple response in the skin after the intravenous injection of pethidine is to be expected if the injection is made into a superficial vein. Although a fall in blood pressure is common after giving pethidine intravenously (Branwood, 1943), sustained hypotension and collapse are rare. The experiments on the cat indicate that, at least in this animal, the hypotensive effect of pethidine is not due solely to the liberation of histamine. It seems likely that in man some factor other than histamine liberation is involved in the production by pethidine of severe hypotension and collapse.

Conclusions

The triple response in the skin after an intravenous injection of pethidine is due to the permeation of pethidine through the vein wall and the liberation of histamine from the skin overlying it. The walls of the superficial veins in the forearm are also permeable to histamine and mepyramine.

It is unlikely that the occasional case of vascular collapse and marked hypotension following an injection of pethidine occurs as the result of the liberation of histamine.

We wish to thank Professor J. L. D'Silva for continued advice and encouragement. Miss P. Archer kindly drew the illustration for Fig. 1.

REFERENCES

- Branwood, A. W. (1943). *Edinb. med. J.*, **50**, 177.
Butler, E. B. (1951). *British Medical Journal*, **2**, 715.
Duguid, A. M. E., and Heathcote, R. St. A. (1940). *Quart. J. Pharm.*, **13**, 318.
Feldberg, W., and Talesnik, J. (1952). *J. Physiol.*, **117**, 3P.
Franklin, K. J. (1937). *A Monograph on Veins*, p. 116. Baillière, Tindall and Cox, London.
Giff, J. H. P. (1947). *British Medical Journal*, **1**, 901.
Halpern, B. N. (1952). *Ibid.*, **1**, 380.
Lewis, T. (1927). *The Blood Vessels of the Human Skin and their Responses*. Shaw, London.
MacIntosh, F. C., and Paton, W. D. M. (1949). *J. Physiol.*, **109**, 190.
Nasmyth, P. A., and Stewart, H. C. (1950). *Ibid.*, **111**, 19P.
Paton, W. D. M., and Schachter, M. (1951). *Brit. J. Pharmacol.*, **6**, 509.
Pearce, C. (1951). *Brit. J. Anaesth.*, **23**, 205.
Zuck, D. (1951). *British Medical Journal*, **1**, 125.

THE "REFLUX" FACTOR IN ERYTHROCYTE SEDIMENTATION TESTS

BY

W. D. ALEXANDER, M.B., Ch.B.

(From the Gardiner Institute of Medicine,
University of Glasgow)

The most popular methods of determining the erythrocyte sedimentation rate are those of Wintrobe and Westergren. The internal measurements of Wintrobe and Westergren sedimentation tubes are similar except that the length of the former is half that of the latter (2.5 mm. diameter; 100 and 200 mm. long). It was therefore surprising to find that the sedimentation rate of a specimen of oxalated blood determined in a Wintrobe tube differed from the determination in a half-filled Westergren tube. It was noticed that the period of aggregation before fall of the cells started was much longer in the Wintrobe than in the Westergren tube. The technique used to fill the tubes would seem to be the cause, since the columns of blood are identical qualitatively and quantitatively.

One way in which the method of filling the tubes differs is that in the Westergren method blood is drawn past the zero mark and then allowed to fall to it, so that a thin film of blood is left on the inside of the tube above the mark; whereas the Wintrobe tube is filled to the zero mark and the pipette withdrawn, leaving the inside of the tube above the mark free of film. In the first few minutes after a Westergren tube is set up there is therefore a reflux of cells and plasma on to the blood level at the zero mark as the blood adherent to the inside of the tube above the mark runs down to the meniscus. This reflux of blood is of course absent in the Wintrobe tube.

Since the presence or absence of this reflux of blood seemed to exert some effect on the sedimentation of oxalated blood, the effect of this factor on the classical Wintrobe and Westergren methods was investigated. The objects of this paper are: (1) to show that reflux of blood on to the meniscus is a factor frequently affecting the half-hour reading in the Wintrobe method of determining the E.S.R.; and (2) to demonstrate, in four cases in which the readings of the two classical methods were in disagreement, that if reflux of blood on to the meniscus was made to occur in the Wintrobe tube the discrepancy disappeared.

Methods

Comparisons between the readings of the standard Wintrobe and Westergren methods were made on blood from 40 unselected patients. By the Westergren method 7 showed a sedimentation rate between 0 and 10 mm., 14 between 10 and 30 mm., and 19 over 30 mm.

In four cases there was disagreement between the one-hour readings of the two methods, a raised Westergren reading being associated with a normal Wintrobe reading. Each of these patients had active organic disease, and a raised E.S.R. was expected. Blood from these four cases, and from a further 24 in which the readings of the methods were in agreement, was examined to determine the effect of reflux of blood on to the meniscus in each of the two techniques.

In the group of 24 patients in whom the standard methods gave similar readings two pairs of tubes were set up: (1) a pair of Wintrobe tubes (containing oxalated blood), and (2) a pair of Westergren tubes (containing citrated blood). In one of each pair reflux of blood on to the meniscus was allowed, in the other prevented.