REFRESHER COURSE FOR GENERAL PRACTITIONERS

PROLAPSE

BY

A. J. WRIGLEY, M.D., F.R.C.S., F.R.C.O.G.

Obstetric Physician and Lecturer in Midwifery and Diseases of Women, St. Thomas's Hospital, London

Prolapse signifies the descent through the vaginal orifice of any structure lying immediately above the pelvic floor. Thus the term may imply descent, singly or in combination, of the uterus, bladder and urethra, anterior wall of the rectum, or pouch of Douglas. Prolapse is usually found in parous women, and therefore it is assumed that weakening of the pelvic floor muscles and fascia and damage to the transverse cervical ligaments are produced by the birth of a baby. Prolapse is rarely seen in nulliparous women; when this does occur it may be associated with a general weakness on the part of the patient as the result of overenthusiastic fasting, or with some lesion in the central nervous system, usually in the lower part of the spinal cord. Prolapse to the laity means descent of the womb, and this may explain the general expectation, which is present even with a few doctors, that its logical treatment is the removal of this offending organ.

Symptoms and Signs

Nearly all women who suffer from prolapse notice the same symptom and phrase their complaint in the words that they "feel something coming down." It is as rare to find prolapse without this symptom as it is to hear of the symptom and discover no prolapse. Concurrently the patient may state that she easily tires and that she has acquired a low backache that becomes progressively more pronounced as the day wears on and is relieved immediately she lies down. Difficulties with micturition may or may not be present. Quite often a patient in whom there exists a large cystocele states that micturition is normal, whereas another with relatively slight descent of the bladder and urethra may be greatly troubled with stress incontinence of urine or frequency of micturition. Occasionally the patient discovers that emptying of the bladder can be completed only by the digital reduction of a large cystocele. A similar trouble with defaecation is dealt with in a like manner in the case of a large rectocele. When a severe degree of prolapse occurs, the patient's discomfort is great, and it is rare for complete procidentia to be tolerated for any length of time.

Although prolapse may occur directly after childbirth, in the majority of patients it develops at about the time of the menopause. This may be due to a general diminution in muscle tone at about this time in a woman's life or to a considerable increase in weight. Rarely, prolapse is suddenly produced following some severe and perhaps unexpected muscular effort such as the lifting of a heavy weight or an attempt to recover from a slip or fall. Prolapse always becomes worse if the patient becomes pregnant.

The diagnosis of prolapse is easy. The patient should be examined lying on her back with the knees drawn up and well separated. She is then asked to give a violent cough or to bear down hard. Prolapse will then be visible, and it will be possible to discover its exact nature and extent. A faulty diagnosis is rarely made either by the patient or by her doctor. Very occasionally a cyst beneath the anterior vaginal wall suggests a cystocele, or a fibroid polyp may simulate uterine descent.

Prevention

Theoretically it should be possible to do much to prevent prolapse, and, indeed, over the last two or three decades

much has been tried. It must be noted, however, that in spite of conscientious painstaking work on the part of the doctor and midwife, and the enthusiastic co-operation of the patient, the condition seems to be as commonly seen as ever it was. Prenatal physiotherapy by education in muscular relaxation and contraction, the prevention of any expulsive efforts during the first stage of labour, the use of the forceps only during the second stage of labour, and renewed physiotherapy after delivery directed toward re-education of the pelvic floor muscles might with reason have been expected to produce good results. There is no evidence that this is so. Those who advocate that early rising after childbirth will lessen the likelihood of prolapse argue with others who believe that the uterine supports will function better if they are allowed a few days' rest and peace and quiet in which to recover. Of all these various methods, the art of midwifery in the conduct of the actual labour is undoubtedly by far the most important.

Treatment of Prolapse

Unless contraindicated, the treatment of prolapse is by operation. Operation is contraindicated if the condition of the patient is such as to make its performance dangerous. It is not advised if the patient is advanced in years, though recent advances in methods of anaesthesia and improved surgical technique have greatly extended the age of safety. Each patient will be judged individually. Operation is not advised if the patient suffers from some chronic illness, and especially if, following a pelvic floor repair, she is likely to be affected by a persistent or troublesome cough. Operation would not be advised if the prolapse was noticed directly after childbirth, for a considerable degree of improvement should be expected in the course of normal involution aided by symptomatic treatment. Operation would not be advised if the patient was young and intended to become pregnant, and never, of course, during pregnancy. A severe degree of prolapse, however, in a woman in the child-bearing period of her life merits operative repair; such an operation is no contraindication to further pregnancy or to the expectation subsequently of a normal delivery. A recurrence of the prolapse in such a patient is very far from inevitable, and the small risk of such a recurrence is justified because the operative safety of a further pelvic floor repair is very much greater than that of caesarean section, which is the only method of avoiding the risk.

Palliative Treatment

This treatment is now reserved for those patients in whom any operative procedure is inadvisable or who refuse operation. A quarter of a century ago many more patients suffering from prolapse were treated palliatively, and operation was often reserved for those few in whom palliative treatment for some reason was ineffective. The change of outlook that has occurred is probably due to the appreciation by doctors and by their patients of the great benefits that follow a correctly performed pelvic floor repair.

Palliative treatment consists in the insertion into the vagina of a ring pessary of such a size that it is larger than the hernial orifice through which the prolapse occurs. The pessary is usually circular and may be made of rubber or of vulcanite. The rubber pessary is usually employed because it is more easily inserted and removed. Rubber is

much more irritant to the tissues of the vaginal wall than vulcanite, and is therefore more likely to produce an offensive purulent discharge. The correct size of pessary to be used in any particular patient can be determined only by trial and error. It is usually advised that the smallest pessary should be inserted which appears to control the prolapse. Such advice inevitably leads to many errors and many trials, and it is much better in the first instance to insert the largest pessary of whose presence the patient is not aware. The patient who for the first time has been so treated for prolapse must be instructed upon the necessity of regular douching to try to prevent infection, and of regular visits to her doctor or nurse for the removal and cleaning, or changing, of her pessary. Most women hate a pessary life, and with good reason.

Operative Treatment

Many and varied have been the operative procedures suggested for the cure of prolapse. Only one has stood the test of time, and that is the operation first evolved at the beginning of the century by the surgeons in Manchester, notable among whom were Donald and Fothergill. The various parts of this operation are designated anterior colporrhaphy, amputation of the cervix, posterior colporrhaphy, and perineorrhaphy. The use of the word colporrhaphy is unfortunate because it implies the use of the vaginal wall in the cure of the condition. The part played by the vaginal wall is exactly analogous to that of the skin overlying the umbilical hernia. It becomes stretched by the hernial protrusion and redundant skin has to be removed. The essential procedure consists in an exposure of the separated inner margins of the two levatores ani muscles with their adjacent fibromuscular fascia. This necessitates dissection beneath the anterior and posterior vaginal walls and exposure of the transverse cervical ligaments by amputation of the cervix. Thus the reconstruction of the pelvic floor and of the perineum is performed. There is no necessity to open the abdomen. Vaginal hysterectomy can be combined with pelvic floor repair and is especially indicated in the presence of some pathological condition of the uterus. It is undoubtedly a more severe proceeding, but the results are excellent.

Aftercare

The care of a patient on whom the Manchester operation has been performed may be divided into the initial period of her stay in hospital and the following two months when she is at home. In the latter period she will certainly be in the hands of her family doctor, and very often, especially in the smaller hospitals, he is also responsible during the first three weeks. For this reason a brief account will be given of the management of these patients until they are well enough to go home. The operation is followed by a definite, and sometimes considerable, amount of pain and discomfort, and morphine should not be withheld during the first 24-36 hours. Thereafter tab. codein. co. or its equivalent should be sufficient. The patient may be returned to the ward with a self-retaining catheter inserted into the bladder. This should be released by the nursing staff at regular intervals and is usually removed on the third or fourth day. By this time much of the initial and local discomfort will have abated and the patient in consequence is far less likely to have difficulty in micturition. If she is able to pass her water naturally the nursing staff must be instructed to pass a catheter regularly on at least two occasions in each 24 hours after the patient has passed water, in order to discover the presence of, and to remove, any residual urine. When by this means less than 2 oz. (60 ml.) is found to remain in the bladder, catheterization may be discontinued. Cystitis is a frequent and troublesome complication of any pelvic operation. The belief that it is most likely to result from repeated catheterization is probably incorrect. A far more probable explanation is that it is due to temporary imperfect emptying of the bladder and

the infection of a considerable quantity of unsuspected residual and stagnant urine. The routine administration of urinary antiseptics should be left to the discretion of the doctor in charge of the case. Aperients should be ordered as soon as the patient has recovered from the immediate ill-effects of the operation. There is no need to purge the patient, nor is it necessary to order an enema. The Manchester operation is not a dangerous proceeding, but one of the few alarming complications is due to occasional phlebo-thrombosis or even thrombophlebitis. Instruction in post-operative deep breathing exercises and early movements on the part of the patient is of great and proved importance in its prevention. Indeed, no harm seems to result from early ambulation, though this must not prejudice the patient's stay in hospital, which should be from two to three weeks, depending upon the severity and extent of the operative repair. Reactionary bleeding is rare, but secondary haemorrhage following amputation of the cervix does occasionally occur at about the end of the first week. It may be severe and even dangerous. An initial injection of morphine $\frac{1}{4}$ gr. (16 mg.) should be given. The patient should be turned on to her left side into the Sims position and a Sims speculum gently inserted into the vagina. A good light is essential. With dry gauze and a pair of dissecting forceps all blood and clot is removed, and the vagina should be carefully, systematically, and firmly packed with dry sterile gauze. In nearly every case this will deal with the condition and no further bleeding occurs after the removal of the gauze plug at the end of 24 hours. Very occasionally, and if the blood soaks past the gauze, it is necessary to take the patient back to the operating theatre and to deal with the bleeding point or area by the insertion of a catgut stitch. The wounds in the anterior and posterior walls of the vagina and in the vaginal vault rarely heal entirely by first intention, and in consequence many patients following a Manchester repair operation acquire for a few days a noticeable and offensive vaginal discharge. When, and if, this occurs the patient should be ordered a twice daily vaginal irrigation with a dilute solution of some non-irritant antiseptic given at body temperature. A solid douche nozzle must not be used, but the fluid should be run into the vagina through the largest size of rubber catheter. Before the patient leaves hospital a digital examination of the vagina must be made. Occasionally the wound on the anterior wall becomes adherent to the wound on the posterior wall. Two or three weeks after the operation the adhesions will be slight and easily separated by the examining finger.

On return home the patient should be told to use her common sense. Prolonged periods of standing, lifting of heavy weights, straining at stool, and similar exertions will all tend to stretch an insecure wound and to undo the repair. Her return to normal activities should be gradual and she should be able to lead a full and normal life once more about three months from the time of her operation. A final examination made about this time is most advisable in every case, and not until then should a return to married life be contemplated.

Conclusion

Prolapse nearly always occurs in women who have borne children, and often first becomes troublesome about the time of the menopause.

The treatment of prolapse is by operative repair unless contraindications exist. The contraindications to the operation are obvious.

The success of the repair operation depends not only on the surgeon but also on all those who take part in the aftercare of the patient up to the end of the third month.

Next Refresher Course Article.—"The Management of Late and Latent Syphilis," by Dr. G. L. M. McElligott.