improvement in most of the cases treated by pneumoperitoneum, which warrants its use in a larger series of patients.

A pneumoperitoneum may be continued indefinitely, refills being given at weekly or, more usually, fortnightly intervals, although in Reich's experience the high position of the diaphragm was maintained for long periods after the pneumoperitoneum was discontinued. This he attributed to the fact that during treatment the patient performed an unconscious respiratory exercise, relaxing the diaphragm and contracting its antagonists, the abdominal muscles, and continued to do so without the aid of a pneumoperitoneum.

Summary

A brief review of pulmonary emphysema and methods of accepted treatment is given.

The progressive nature of the disease and the misery which is the lot of those severely affected warrant exploration of all new avenues of treatment.

Nine severe cases of pulmonary emphysema treated by pneumoperitoneum are presented, with results encouraging enough to justify further study. The rationale of pneumoperitoneum is to increase the intra-abdominal pressure sufficiently to cause elevation of the diaphragm during expiration, and yet not to impede its contraction during inspiration. Increased excursion of the diaphragm is thus obtained, with consequent improvement in pulmonary ventilation.

The vital capacity was increased in seven cases and the excursion of the diaphragm, measured radiologically, was improved in all cases. No improvement was evident in two cases with a history of congestive cardiac failure and in one case with asthma.

REFERENCES

- Alexander, H. L., and Kounz, W. B. (1934). *Amer. J. med. Sci.*, **187**, 687.
 Banyai, A. L. (1946). *Pneumoperitoneum Treatment*. Mosby, St. Louis.
 Christie, R. V. (1934). *J. clin. Invest.*, **13**, 295.
 — (1944). *British Medical Journal*, **1**, 105.
 Furman, R. H., and Callaway, J. J. (1950). *Dis. Chest.*, **18**, 232.
 Gordon, B. (1934). *Amer. J. med. Sci.*, **187**, 692.
 Katz, L. N. (1946). *Electrocardiography*, 2nd ed. Kimpton, London.
 McMichael, J., and Sharpey-Schafer, E. P. (1944). *Quart. J. Med.*, n.s. **13**, 123.
 Meakins, J., and Christie, R. V. (1934). *J. Amer. med. Ass.*, **103**, 384.
 Monaldi, E. (1937). *Rev. argent. Tuberc.*, **3**, 161.
 Mounsey, J. P. D., Ritzmann, L. W., Selverstone, N. J., Briscoe, W. A., and McLemore, G. A. (1952). *Brit. Heart J.*, **14**, 153.
 Piaggio-Blanco, R. A., and Sayagues, C. (1938). *Rev. Tuberc. Urug.*, **7**, 103.
 Reich, L. (1924). *Wien. Arch. inn. Med.*, **8**, 245.
 Rubin, J. H., and Gass, G. D. (1948). *Canad. med. Ass. J.*, **59**, 162.
 Sisti, M. A. (1936). *Policlinico, Sez. prat.*, **43**, 2099.
 Whitfield, A. G. W., Arnott, W. M., and Waterhouse, J. A. H. (1950). *Quart. J. Med.*, n.s. **19**, 319.
 — Smith, O. E., Richards, D. G. B., Waterhouse, J. A. H., and Arnott, W. M. (1951). *Ibid.*, n.s. **20**, 247.

In his preface to the annual statistical report of the Middlesex Hospital for 1952, Colonel the Honourable J. J. Astor, chairman of the board of governors, points out that, while in the past the main function of the report was to persuade the public to subscribe to the hospital's funds, its present task "is to say whether public money, assigned by Parliament for the upkeep of their hospitals, has been used to the best advantage." During the year under review there were 633,000 out-patient attendances at the hospital and its constituent institutions, compared with 582,000 in 1951. The physiotherapy clinic accounted for 219,000, of whom 10,000 were new patients; 117,000 attended the rheumatism clinic, and 40,000 attended for radiotherapy. In-patients numbered 14,800, a slight increase on the previous year. Good progress is reported in converting Latimer House, acquired in 1951 to serve as an institute of clinical research and experimental medicine, and the institute will be in action later this year.

SENSITIVITY STUDIES ON SHIGELLA SONNEI

BY

G. B. FORBES, M.D.

Pathologist, Kent and Canterbury Hospital

This report concerns a study of the antibacterial activity of streptomycin, chloramphenicol, aureomycin, and terramycin against *Shigella sonnei*, together with observations on the suitability of these antibiotics for the treatment and control of Sonne dysentery.

Seventeen representative strains of *Sh. sonnei* were collected from separate outbreaks and isolated cases of Sonne dysentery which occurred in East Kent and Hertfordshire. The identity of the strains was confirmed by biochemical tests and agglutination to titre with specific typing serum. In addition the strains were grouped according to their bacteriophage reactions (Hammarström, 1947) and fermentative behaviour towards xylose (Bojlen, 1934) to ensure that the collection included a variety of types. Dr. Mayr-Harting, of Bristol, kindly performed the bacteriophage tests, and found that most strains belonged to phage type 3, a type commonly encountered in this country (Cooper and Mayr-Harting, 1951); five other phage types were represented (Table I). One of the strains fermented xylose; the others were xylose-negative and gave the usual biochemical reactions of strains isolated in this country (Carter, 1937).

TABLE I.—Xylose Fermentation and Bacteriophage Reactions

Strain	Xylose Fermentation	Phage Type
1, 2, 4, 7, 8, 9, 10, 11, 14, 15, 16, 17	—	3
3	—	85
5	—	32
6	—	66
12	—	23
13	+	? 17b

In-vitro Experiments

I. Tube Sensitivity Tests

Method.—Graded concentrations of each antibiotic were prepared in nutrient broth and tubed in 1-ml. amounts; 0.02 ml. of 1:100 dilution of an overnight broth culture of each strain was added to the appropriate set of tubes, which were then incubated at 37° C. The lowest concentrations of antibiotic required for growth suppression as judged by complete absence of turbidity and bacterial deposit were noted after 12, 18, 24, 36, and 42 hours' incubation.

Results.—The results of these tests, expressed as the average sensitivity of the 17 strains to each antibiotic, are given in Table II. It will be seen that the end-points increased progressively with prolongation of the incubation period beyond 12 hours. With the more stable antibiotics (streptomycin, chloramphenicol, and terramycin) the inhibiting concentrations rose slightly, to give readings at 42 hours approximately twice as high as those obtained at 12 hours.

TABLE II.—Mean Sensitivity of 17 Strains After Varying Periods of Incubation

Antibiotic	Hours' Incubation				
	12	18	24	36	42
Streptomycin	4.94*	6.23	7.29	7.76	8.47
Chloramphenicol ..	5.52	7.41	7.88	8.47	8.58
Aureomycin	1.01	2.00	4.00	7.53	9.06
Terramycin	2.06	2.94	3.76	3.88	4.35

* Minimal inhibiting concentration in µg./ml.

In the case of aureomycin, which is notoriously unstable at 37° C., there was a ninefold increase during the period of observation. In fairness to aureomycin, the readings obtained after the shortest workable incubation period—namely, 12 hours—were regarded as being most suitable for comparative purposes. These results are set out in Table III,

TABLE III.—Range of Sensitivity of 17 Strains After 12 Hours' Incubation

M.I.C. at 12 Hours ($\mu\text{g./ml.}$)	Strepto- mycin	Chloram- phenicol	Aureo- mycin	Terra- mycin
8	4	7		1
4	13	9		15
2		1	1	1
1			12	
0.8			4	
Range	4-8	2-8	0.8-2	1-4

and may be summarized as follows: (1) all strains were naturally sensitive to all four antibiotics, being inhibited by 8 $\mu\text{g./ml.}$ or less; (2) the strains exhibited a narrow range of tolerance to each of the four antibiotics; (3) on a weight-for-weight basis aureomycin and terramycin were slightly more effective as growth inhibitors than streptomycin and chloramphenicol.

II. Mode of Action of Antibiotics

Tests were carried out in order to compare the speed and mode of antibacterial action of the four antibiotics on *Sh. sonnei* in the presence of faecal matter.

Method.—A gramme of faeces from a healthy subject was suspended in physiological saline and the suspension distributed in four equal amounts. The specimens were seeded with approximately 100,000 *Sonne* organisms, and, after thorough shaking with beads to ensure an even distribution of the organisms, 0.02 ml. was withdrawn from each specimen and subcultured on plates of desoxycholate medium. The antibiotics under test were then added to the inoculated specimens, so that each specimen contained one of four antibiotics in a strength of 10 $\mu\text{g./ml.}$ in one experiment and 500 $\mu\text{g./ml.}$ in a second experiment. The specimens were incubated at 37° C., and at hourly intervals up to eight hours, and again at 24 hours, subcultures were made as described above. After overnight incubation the subcultures were examined and colonies of *Sh. sonnei* were identified and counted.

Results.—The results of these experiments are given in Tables IV and V. At a concentration of 10 $\mu\text{g./ml.}$ streptomycin exerted a bactericidal action on *Sh. sonnei*, killing all organisms in less than five hours, while the other antibiotics were essentially bacteriostatic in their mode of action. At

TABLE IV.—Survival Time of *Sh. sonnei* in Presence of 10 $\mu\text{g.}$ of Antibiotic per Gramme of Faeces (pH 7.4)

Antibiotic	Hours' Exposure									
	0	1	2	3	4	5	6	7	8	24
Streptomycin	56*	50	23	15	2	0	0	0	0	0
Chloramphenicol	57	46	60	39	50	56	37	30	33	10
Aureomycin	60	52	48	45	60	60	25	46	25	3
Terramycin	50	40	43	53	50	30	30	33	26	2

* Number of colonies of *Sh. sonnei* on subculture.

TABLE V.—Survival Time of *Sh. sonnei* in the Presence of 500 $\mu\text{g.}$ of Antibiotic per Gramme of Faeces (pH 8)

Antibiotic	Hours' Exposure									
	0	1	2	3	4	5	6	7	8	24
Streptomycin	500*	0	0	0	0	0	0	0	0	0
Chloramphenicol	500	400	128	18	4	10	16	15	9	0
Aureomycin	500	0	0	0	0	0	0	0	0	0
Terramycin	500	0	0	0	0	0	0	0	0	0

* Approximate number of colonies of *Sh. sonnei* on subculture.

a concentration of 500 $\mu\text{g./ml.}$ streptomycin, aureomycin, and terramycin were highly bactericidal, killing all *Sonne* organisms within one hour. Chloramphenicol effected a reduction in the population of *Sonne* organisms, but a number remained viable while in contact with 500 $\mu\text{g./ml.}$ for as long as eight hours. These experiments were repeated several times with close reproduction of results. The tests were also carried out using nutrient broth in place of the faecal suspension, and again the results were comparable, although terramycin in this medium was slightly less bactericidal at the higher concentration.

III. Acquisition of Resistance

Method.—Sensitivity tests were carried out on three strains by the conventional tube-dilution technique as described in experiment I. After 18-24 hours' incubation at 37° C. the highest concentration of antibiotic which permitted distinct visible growth was noted, and the culture at this end-point was used to inoculate a fresh set of concentrations on a slightly higher range. This procedure was repeated daily for eight consecutive days.

Typical patterns of acquired resistance *in vitro*.

Results.—*Sh. sonnei* acquired resistance to streptomycin more readily than to the other antibiotics. The rate of acquisition of resistance, however, varied not only with different strains but also when the tests were carried out in duplicate on any one strain. But at least it can be stated that the lowest increase in resistance to streptomycin (a 64-fold increase) was equal to the highest rate of increase acquired to the other antibiotics. The strains developed an eightfold increase in resistance to both aureomycin and terramycin, and the results of repeat tests were quite consistent. Chloramphenicol did not favour the development of resistance to the same extent as streptomycin, the average increase on testing three strains being $\times 35$. Typical patterns of the development of tolerance to each of the four antibiotics are given in the Chart.

Clinical Experience with Streptomycin

Of the four antibiotics, only streptomycin has been used in enough cases of *Sonne* dysentery to allow an assessment of its clinical value.

Sixteen acute cases and one carrier were treated with this antibiotic in a dose of 0.5 g. twice daily for two to three days. The drug was given orally, mixed with flavouring agents.

Clinical cure was achieved within 24 hours of beginning treatment in all acute cases, and bacteriological cure, as judged by repeated negative stool cultures (an average of 4.3 per patient), was prompt and complete in all but one case. A sample of faeces from this patient yielded four colonies of *Sh. sonnei* on the sixth day after treatment,

although two earlier post-treatment specimens had been negative. Three subsequent specimens from this patient gave negative results. No case relapsed clinically or bacteriologically during the follow-up period, which in some cases was as long as six months.

In addition, two cases of Sonne dysentery which relapsed one week after treatment with a sulphonamide were promptly cured clinically and bacteriologically with streptomycin. Another patient was treated with chloramphenicol for five days, with good clinical response, but a very heavy growth of *Sh. sonnei* was obtained from specimens of faeces sent during treatment and on the day after treatment. Streptomycin was then given, and the infecting organism disappeared from the faeces within 48 hours.

A female member of the staff of a residential nursery was found to be a symptomless carrier of *Sh. sonnei* during investigations following an outbreak of Sonne dysentery which had occurred in the nursery two months earlier. She was sent on leave, and a specimen of faeces submitted to this laboratory confirmed that Sonne organisms were being excreted in very large numbers. Streptomycin was given orally for two days, and three post-treatment specimens of faeces received at weekly intervals were negative for *Sh. sonnei*. The patient returned to work, and a specimen submitted five months later was again free of Sonne organisms.

Discussion

Specific treatment in Sonne dysentery should aim not only at effecting a prompt clinical cure but also at rapidly clearing the intestinal tract of *Sh. sonnei* in order to reduce the period of infectivity to a minimum. As the disease is usually mild and of short duration, it is difficult to assess the value of specific remedies on the basis of clinical response alone; the rate and completeness of clearance is probably a more reliable index of therapeutic efficiency and adequate control. The sulphonamides, particularly sulphadiazine and the poorly absorbed sulphonamide preparations, are of undoubted value in the management of bacillary dysentery, but Sonne organisms often persist or reappear in the faeces during convalescence (Fairbrother, 1944). Moreover, in certain parts of the world, for example Japan, the majority of strains of *Sh. sonnei* are naturally resistant to sulphonamides (Tateno, 1950).

Oral therapy with antibiotics is an obvious alternative to sulphonamide medication in Sonne dysentery, and the experiments described above help to predict which antibiotics are likely to be most useful in this disease.

Welch (1950) assayed the faecal excretion of chloramphenicol, aureomycin, and terramycin after the oral intake of 1 g., and obtained levels of approximately 500 µg. per gramme of faeces with each antibiotic. Streptomycin, which is not absorbed from the bowel, was found in even higher concentrations in the faeces (at least 9,000 µg./g.) by Zintel *et al.* (1945). These levels greatly exceed the concentrations which were required to inhibit the growth of all 17 strains of *Sh. sonnei* used in this study, and from this it is apparent that all four antibiotics are potentially effective in the treatment of Sonne dysentery. However, streptomycin, aureomycin, and terramycin were found by experiment to exert a rapid bactericidal action at concentrations obtainable in the faeces following oral administration, and in the case of streptomycin the lethal effect was evident when the concentration in the faeces was reduced to 10 µg./g. It is likely, then, that these three antibiotics, especially streptomycin, would effect a complete and permanent eradication of Sonne organisms from the faeces of clinical cases. On the other hand, chloramphenicol exerted only a weak bactericidal action at a concentration of 500 µg., and in theory it should be less efficient at ridding the faeces of Sonne organisms.

The efficacy of streptomycin in Sonne dysentery has been confirmed by clinical experience, both in the present study and by others, notably Ross *et al.* (1949). In fairness to chloramphenicol it should be added that Ross *et al.* (1950) obtained equally good results with this antibiotic, and it

would seem that the theoretical objection to chloramphenicol mentioned above is probably not of significance in clinical practice.

Certain other factors would influence one in favour of streptomycin for the routine treatment of Sonne dysentery. It is relatively cheap and is readily available. It is non-toxic when taken by mouth, and, unlike chloramphenicol and aureomycin, does not irritate the gastric and intestinal mucosa—a point of some importance when treating an infection which itself produces gastro-intestinal upset. Moreover, it is palatable if flavoured, and is well tolerated by children. There are, however, two minor disadvantages peculiar to streptomycin: (1) the results of experiment III indicate that *Sh. sonnei* develops resistance slightly more readily to streptomycin than to the other three antibiotics, but it is unlikely that this disadvantage will prove of any consequence in clinical practice, since the duration of treatment will not normally exceed two or three days; and (2) an acute fulminating type of Sonne dysentery is occasionally met with in young children (Lewis and Claireaux, 1951), and Tatham *et al.* (1951) succeeded in isolating *Sh. sonnei* from the blood of such a case. Obviously an antibiotic which is absorbed from the bowel is to be preferred to streptomycin for the treatment of the septicæmic type of infection, or, if streptomycin is chosen, it should be given parenterally as well as orally.

Summary

Sh. sonnei was highly and almost equally sensitive to streptomycin, chloramphenicol, aureomycin, and terramycin *in vitro*, all test strains being inhibited by 8 µg./ml. or less.

All four antibiotics exerted a bactericidal action on *Sh. sonnei* in a concentration of 500 µg./ml., but the lethal effect of chloramphenicol was weak compared with the other antibiotics. Streptomycin alone was bactericidal at both low and high concentrations.

Sh. sonnei acquired resistance *in vitro* slightly more readily to streptomycin than to the other antibiotics.

Streptomycin, given orally, has proved a highly effective therapeutic agent in the management of Sonne dysentery.

My thanks are due to Dr. B. A. Thompson, Watford, for providing me with a few of the test strains, to Dr. A. Mayr-Harting for typing the strains, to Miss P. Reeves for technical assistance, and to the Wright-Fleming Institute of Microbiology for the supply of terramycin.

REFERENCES

- Boijén, K. (1934). *Dysentery in Denmark*. Copenhagen.
Carter, H. S. (1937). *J. Path. Bact.*, **45**, 447.
Cooper, K. E., and Mayr-Harting, A. (1951). *British Medical Journal*, **2**, 271.
Fairbrother, R. W. (1944). *Ibid.*, **2**, 489.
Hammarström, E. (1947). *Lancet*, **1**, 102.
Lewis, I. C., and Claireaux, A. E. (1951). *Ibid.*, **1**, 769.
Ross, S., Burke, F. G., Rice, E. C., Bischoff, H., and Washington, J. A. (1949). *J. Amer. med. Ass.*, **141**, 183.
— Washington, J. A., and Stevens, S. (1950). *Ibid.*, **143**, 1459.
Tateno, I. (1950). *Jap. J. exp. Med.*, **20**, 795.
Tatham, P., Williams, T. P., and Stewart, G. T. (1951). *Lancet*, **1**, 997.
Welch, H. (1950). *Ann. N.Y. Acad. Sci.*, **53**, 253.
Zintel, H. A., Flippin, H. F., Nichols, A. C., Wiley, M. M., and Rhoads, J. E. (1945). *Amer. J. med. Sci.*, **210**, 421.

Owing to fatalities amongst children, a meeting of branch representatives of the Pharmaceutical Society has requested the Ministry of Health to give publicity to the importance of tablets and other medicines being kept out of reach of children. It was also suggested that, in addition to advertisements, the Ministry should approach the B.B.C. and television authorities to produce a documentary to educate the public in the extreme potency of many of the medicines being dispensed nowadays.