REFRESHER COURSE FOR GENERAL PRACTITIONERS

INTERMITTENT CLAUDICATION

BY

J. B. KINMONTH, M.S., F.R.C.S.

Assistant Surgeon and Assistant Director of Professorial Unit, St. Bartholomew's Hospital

The term "intermittent claudication" is derived from the latin *claudicare*, to limp or halt, not as is sometimes thought from *claudere*, to shut, a derivation which would suggest an erroneous explanation for the symptom by implying that the blood vessels to the limb were subject to intermittent episodes of closure. In fact the symptom is caused in most cases by thrombosis in a segment of a main artery. The collateral circulation past the thrombosis supplies enough blood for muscles at rest but too little for the increased needs during exercise. The thrombosis may originate from injury to the vessel by violence or it may start upon an area of disease in the intima such as an atheromatous plaque. Atheroma is by far the commonest cause of arterial thrombosis at all ages. It is a much more serious lesion than fibrosis or calcification of the tunica media, which may make vessels feel hard to palpation and opaque to x rays but do not themselves cause thrombosis.

Claudication is a painful symptom and therefore alarming to the patient. Usually it does not itself prove a great handicap, and its importance lies more in the other changes in the limb or cardiovascular system that may accompany it. Its treatment, therefore, is almost always incidental to measures planned to improve the circulation in the limb as a whole.

Diagnosis

A carefully taken history is the most important element in diagnosis. No condition other than arterial insufficiency exactly reproduces the sequence of pain after a definite period of exercise relieved by resting and reproduced constantly by the same amount of exercise. Even when the patient also has pain at rest it is usually described as being different in nature from the pain brought on by exercise, and is usually situated in the foot or ankle rather than the leg. Patients with osteoarthritis may complain of pain on exercise, but it is usually worse on starting and wears off with continued movement. Muscular rheumatism or fibrositis may sometimes cause confusion, but the history is never exactly the same, particularly in that it comes in attacks or episodes and is variable, unlike true intermittent claudication. Other symptoms of peripheral ischaemia such as coldness, numbness, or paraesthesiae in the limb and even ulceration or necrosis may coexist and give confirmatory evidence in the diagnosis.

The most valuable part of the clinical examination is palpation of the pulses. In normal people femoral, popliteal, and two ankle pulses can be felt, but in about one subject in ten the dorsalis pedis pulse is absent. Some difficulty may be experienced in feeling the popliteal pulse in fat patients. Should it be impalpable by the usual method with the knee flexed to a right-angle an alternative method can be used with the knee hyperextended on the couch.

Most patients with claudication have no palpable pulses below the femoral. Very rarely a patient is found with normal pulses and no physical signs of disease in the limbs despite a definite history of intermittent claudication. Usually he is a young man who can walk a quarter of a mile (half a kilometre) or more before pain occurs. Arteriography is necessary in such a case to confirm the diagnosis, and mural thrombosis or a very short segment of complete obliteration is the usual finding. Apart from such exceptional cases

no special methods are necessary in the diagnosis of peripheral arterial disease other than those of history-taking and clinical examination.

Conservative Treatment

Thrombosis may occur in a main peripheral artery and give rise to acute symptoms resembling those of embolism, with pain, pallor, and coldness in the foot, but usually it occurs insidiously with few if any symptoms other than the onset of claudication. After the initial stabilization of the circulation a much slower hypertrophy of the collateral vessels takes place, leading to an improvement in the walking distance which may occur over weeks or months. This later spontaneous improvement in blood flow occurs with conservative treatment in many cases, and is the probable explanation of the claims made for various forms of treatment of doubtful efficacy. It is therefore a good rule to allow ample time for this natural improvement to take place, aided by conservative measures, before contemplating more radical ones. In cases in which claudication is the chief or only symptom operation should not be considered until the symptoms have become constant and stationary for three to six months.

Cardiovascular Assessment.—Peripheral arterial obstruction other than that due to trauma is almost always part of a general disease affecting other parts of the body. The heart may be affected, and so its condition must be investigated and any necessary treatment given. If operation is considered it must be decided not only whether the patient is fit for it but whether it will be of benefit for him to walk further. It would be an unhappy sequence for a patient to be cured of claudication only to experience angina pectoris because of his ability to walk further. Evidence of cerebral arteriosclerosis must also be taken into account in deciding treatment.

Diet.—There is evidence that in both man and animals a high intake of fat and lipoids in the diet tends to cause the development of atheromatous lesions in the arteries. It has not, however, been shown that omitting fat from the diet will check further progress of the disease once it has started. If the patient is obese he should be put on a low-calorie diet with the immediately practical object of giving his ischaemic legs less weight to carry rather than in any more ambitious hope of stopping further development of the disease.

Vasodilator Drugs.—The most generally useful is tolazoline hydrochloride ("prisol"). It is given orally in doses of 50 mg. thrice daily. If side-effects such as giddiness or nausea occur the dose is reduced to 25 mg. three times a day.

Smoking and Alcohol.—There have been many attempts to prove that some of the acute forms of arterial obliterative disease are caused by sensitivity to tobacco, but they have not met with general acceptance. There is, however, ample proof that smoking has a constrictor effect on peripheral vessels, leading to temporarily diminished blood flow even in normal subjects. This is due both to overbreathing and to the effect of the tobacco itself. For this reason alone patients with peripheral arterial disease should be advised to give up smoking. Alcohol, on the other hand, is a good peripheral vasodilator and a moderate regular intake is helpful. The patient who is disconsolate at the loss of the one minor vice may be cheered by the retention of the other.

Buerger's exercises. In which the patient elevates the limb for half a minute, holds it horizontal for a minute, and then dependent for a minute, should be done three times and repeated three times a day. They are designed to stimulate the collateral circulation.

General Care of the Feet.—Patients with intermittent claudication very often have deficient circulation in the skin, so that the feet need to be particularly cared for. Minor injuries may lead to indolent ulcers or infection which may further injure the circula-

tion and cause necrosis or gangrene. Shoes must be comfortable and kept in good repair so that friction and blistering or injury by projecting nails never occur. If a chiropodist is available regular visits are advisable; if not the patient himself must be careful when cutting his nails not to injure the soft tissues. The toe-nails are best cut immediately after a bath, when they are soft and unlikely to splinter or leave sharp projections. Interdigital fungus infections are dangerous because they make excoriation

FIG. 1.—Palpation of the popliteal pulse by the flexed knee method.

FIG. 2.—Palpation of the posterior tibial pulse behind the medial malleolus.

and infection likely. A good prophylactic regime is the routine use, after bathing, of 1% salicylic acid in surgical spirit followed by the application of zinc oxide and salicylic acid dusting powder (N.F., 1952), paying particular attention to the interdigital clefts. In order to maintain as much circulation in the feet as possible the patient must at all times keep himself and his feet warmly clad and particularly avoid getting his feet wet and cold. Hot baths are good because they produce reflex vasodilatation. The feet should be kept out of the hot water at first until reflex vasodilatation is caused by the rest of the body warming up: direct application of heat to tissues may increase their metabolic demands beyond the capabilities of supply by a sluggish circulation. The same principle applies in bed. The bed should be warmed and bed-socks worn, but a hot-water bottle is better applied to the trunk than to the feet.

Exercise.—In those cases in which thrombosis has occurred as an acute episode complete rest in bed will be necessary until the circulation is stabilized. Later on in these cases, and in others in which intermittent claudication has followed insidious thrombosis, advice will be needed on how much or how far the patient should walk. Moderate exercise is beneficial, as it provides some stimulus to the collateral circulation, but the patient should stop before or on the onset of the least discomfort. He should not persist until the onset of actual pain, for this may cause tissue damage by ischaemia.

The details of conservative treatment conscientiously followed often prevent the deterioration or loss of a limb. They should be carefully explained to the patient, who should know that successful treatment and perhaps the fate of the limb depend largely upon his own efforts. In cases in which operation is necessary the same care must continue afterwards. Few, if any, surgical measures can restore the circulation to normal.

A number of methods of treatment have been advocated which have been popular from time to time but which need complicated apparatus or an amount of skill, time, and trouble out of proportion to the benefit achieved; some of them carry a disproportionate risk. In this group or in its boundaries lie such methods as intermittent venous occlusion, repeated spinal anaesthetics, caudal blocks and paravertebral blocks, and repeated intra-arterial injections of vasodilators. So-called chemical sympathectomy is of value in certain cases. Old or decrepit patients unfit for general anaesthesia may benefit, but more often when the chief symptom is pain at rest or impending necrosis rather than intermittent claudication. It is done by injecting a few millilitres of 5% phenol around the lumbar sympathetic chain.

FIG. 3.—Palpation of the dorsalis pedis pulse.

Sympathectomy

Sympathectomy is the most generally useful operation for patients with peripheral arterial disease. It improves the blood supply by dilating collateral vessels. The effect is more pronounced in the skin than in the muscles of the limb, and usually only those patients with the less severe forms of claudication have their symptoms abolished. The best results are achieved in patients whose pain does not halt them completely but allows further progress at a reduced pace. Many patients with claudication also have impaired circulation in the skin, which is more important because of its influence on the fate of the limb. This is considerably and permanently helped by sympathectomy and is often the primary indication for operation. Pain in the foot at rest, provided there is no area of necrosis or gangrene, responds well to sympathectomy and is not a contraindication to the operation.

The sympathetic chain and ganglia lying on the bodies of the second, third, and fourth lumbar vertebrae are removed. Excision of the first lumbar ganglion is unnecessary in most cases: it may have the disadvantage of hindering sexual function by interfering with ejaculation. The old operation of periarterial sympathectomy has been abandoned, as has also arteriotomy of the thrombosed segment, except in those cases in which the artery is superficially placed and causing pain.

Tenotomy

It has been found that inactivation of some of the leg muscles often enables a patient to walk further before the onset of pain. The most probable explanation for this is that the demand for blood during exercise is reduced. Division of the Achilles tendon inactivates the gastrocnemius

and soleus muscles, leaving the remaining calf muscles for the patient to walk with. Naturally there is some loss of power and the patient is less agile in ascending stairs or in any action requiring vigorous plantar flexion. For this price a marked improvement in walking distance may be gained. The strongest indication for tenotomy is claudication in one limb after walking a short distance, thirty to forty yards (30 metres) perhaps, in an older patient with relatively good skin nutrition. Many patients require sympathectomy either before or at the same time to improve skin nutrition, but this is not always so and tenotomy may be done as an isolated procedure. A good pre-operative test of the possible effect of tenotomy on a patient with one-sided pain is to raise the heel $\frac{1}{4}$ in. (2 cm.) by placing a felt pad under it. If this increases the walking distance tenotomy is likely to improve it further still. Sometimes the improvement after the heel has been raised is so pronounced that it may be permanently raised, thereby making operation unnecessary.

Bilateral tenotomy may be needed in patients who have pain in both legs after walking a very short distance. The operation can be done under local analgesia as an open or closed procedure. In either case pre- and post-operative ankle exercises must be taught and carried out, and the patient made to start walking within 24 hours to avoid the risk of thrombosis in the veins of the calf muscles. Other means of inactivating muscles in the leg, such as neurotomies, have been tried, but they are less certain, less effective, and often more extensive procedures than tenotomy. They have now been given up.

Reconstructive and Other Operations

In theory the ideal way of treating arterial obliteration is by replacement with some form of tube to conduct the blood. It is possible in practice to treat a small number of patients in this way. The problem has two aspects: that of the material to be used, and that of the choice of patient who may be helped.

Material for Grafting.—At present the material of choice is an arterial homograft. This is a vessel which has been taken under sterile precautions from the body of another person recently dead. Death must have been due to some cause other than infection, and the best material comes from young subjects who have had fatal accidents. The grafts are rapidly frozen to -79° C. and banked at this temperature, which preserves them for long periods in a state ready for use. The second choice is a vein autograft—that is, a vein taken at operation from the patient himself. The great saphenous is the vein usually used. This is a good method for use in war, as no extra organization for storage or supply is needed. It is not so useful in replacing defective arteries larger than the popliteal because of the discrepancy in size, and it is more difficult to anastomose to a thickened arteriosclerotic vessel than an arterial graft would be. Other materials are unsuitable for various reasons. Arteries from the patient himself cannot be spared, nor can veins larger than the great saphenous be used without causing disability. There is still no way of proved safety for preparing heterografts—that is, arteries from different animal species—and the same applies to the various plastic and cloth tubes which have been tried under experimental conditions.

Choice of Patient.—The small group of patients suitable for treatment by grafting consists of those in whom the obstruction is enough to produce marked claudication or necrosis, but in whom the disease is relatively localized in the arterial tree. A number of patients are excluded on general clinical grounds because of cardiac, cerebral, or diffuse disease elsewhere in the body. Others are found to be unsuitable on arteriographic evidence of extensive disease in the arteries of the limb. In others, particularly those with a short block, good collaterals, and mild symptoms, grafting is unnecessary because they do well with less radical treatment, such as sympathectomy.

Thrombo-endarterectomy.—Another method of restoring patency to thrombosed arteries which is occasionally useful is that known as "thrombo-endarterectomy" or "désobstruction." Here a longitudinal incision is made in the vessel wall and the clot removed together with the intima and part of the media along a plane of cleavage which is

usually found without difficulty. A surprisingly smooth surface is left. The opening in the vessel is then sutured. This method has a place in removing very short areas of clot from large arteries such as the iliac in which there is a wide lumen and a vigorous stream to discourage thrombosis. The operation is not applicable to arteries of narrower calibre in which, as might be expected, clotting recurs.

Special Investigations

The state of the peripheral circulation may be studied in several different ways. Venous occlusion plethysmography enables the actual flow of blood in the limb to be measured quantitatively, and is therefore particularly valuable. Skin temperature, and other tests, measure the flow qualitatively. These studies are valuable in research, and in the more unusual and difficult cases of arterial obliteration, but for routine work a careful clinical examination is quite adequate. Arteriography is an exception to this rule. It is essential if arterial grafting is to be considered, as the exact extent of the clot, and the state of the arterial tree, must be known. Percutaneous puncture allows repeated examinations, and is usually done at the level of the common femoral artery. When it is suspected, because of absence of the femoral pulses or because of pain in the buttocks or thighs on walking, that the iliac vessels may be involved the blocked areas must be visualized by injection at a higher level. This can be done by means of translumbar aortography.

Paravertebral sympathetic block has been tried as a test to predict the effect of sympathectomy, but its effect upon claudication has not been found to correspond at all closely with that of operation. It has therefore little place as a pre-operative test in patients with claudication, but is sometimes useful in patients with ischaemia of the skin.

Next Refresher Course Article.—"Myxoedema," by Dr. A. W. Spence.

BRITISH EMPIRE LEPROSY RELIEF ASSOCIATION

The annual meeting of the British Empire Leprosy Relief Association, which was held in London on May 21, was to have been addressed by the Rt. Hon. Oliver Lyttelton, but his absence in Kenya prevented his attendance. In his place an address was given by the Rev. P. T. B. CLAYTON, founder of Toc H. Mr. Clayton said that a new era of hope had dawned with the sulphone treatment of leprosy, but, although the increase in services and child adoptions had been considerable, the areas where treatment was available to the majority of sufferers were still few, and elsewhere only one sufferer in thirty, and until recently often only one in a hundred, were within reach of cure. In Nigeria, leprosy control had been gained by establishing clinics and segregation villages around a central leprosarium, even before Dr. John Lowe's experiments at the association's research unit with volunteer patients had brought still more effective relief. The child adoption scheme had also contributed to the wane of the disease in parts of Nigeria. The demand was for fresh centres in the more remote hotbeds of infection. Although new homes were being built, they existed only in tens where hundreds were wanted, and little advance had as yet been made towards the establishment of the East Africa Research Centre. Thanks to the work of voluntary helpers, said Mr. Clayton, and the increasing aid from firms and trade unions, the receipts of the association had risen from £11,450 in 1939 to £69,876 in 1952.

Sir LEONARD ROGERS, surveying twenty years of investigation and treatment of the disease, spoke of the difficulties which arose when fear of segregation drove victims to hide the disease until it was far advanced, so that doctors could not find early cases to treat. In the Nigerian