tive procedure proved too much for the patient, although it was the only possible measure. There is no certain way of forecasting the outcome of an operation on a patient on the brink of death, and we have been surprised that some have survived and returned to a normal life. The commoner degenerative diseases—renal failure, vascular degeneration, coronary heart disease, cerebral arteriosclerosis—all combined to bring about the fatal issue in these cases.

As might be expected, there were a number of purely vascular causes of death (cerebral thrombosis two cases, coronary disease four cases, iliac-vein thrombosis one case). These usually occurred suddenly and without warning in a patient whose progress had hitherto been satisfactory.

Conclusion

We believe that even major surgical procedures are often justifiable in old age. It is necessary, however, that one's surgical valour should be tempered with discretion and that the operation should be the minimum consistent with cure or symptomatic relief. A colostomy, for example, may be all that is justifiable in large-bowel obstruction, and sometimes, though less often, permanent suprapubic cystostomy is all that is justifiable in prostatic urinary obstruction. We also believe that unless cure or symptomatic relief is likely to be achieved it is better not to operate. It is unjustifiable and unkind to attempt desperate surgical ventures that would be fitting in younger patients.

A better selection of cases would improve the mortality rate, but as usual the problems of selection are insuperable. It is as important to give relief to all those who would survive the surgical ordeal as to withhold operation from those who would not stand it. With experience we have found that an accurate assessment of the result of surgery can often be made from the general clinical condition of the patient, and this is often the best guide to fitness for operation.

With continued improvement in surgical technique, the use of new antibiotics, further knowledge of the biochemistry of disease, and further improvements in anaesthesia we are looking forward to the time when only patients over 80 will be discussed in the surgery of old age.

Summary

A series of 537 operations were performed on patients over the age of 70, of whom 119 died, giving a total mortality rate of 22.2%.

Of these patients, 434 underwent major operations, and 118 of them died, giving a mortality rate of 27.2%. The causes of death are analysed and discussed.

We believe that surgery in old age is rewarding more often than is generally believed, and we think this will prove even truer in the future.

REFERENCES

- Bostick, W. L. (1945). *Amer. J. Path.*, **21**, 1143.
Carp, L. (1950). *British Medical Journal*, **2**, 1198.
Childs, P., and Mason, S. A. (1949). *Ibid.*, **2**, 1389.
Cutler, C. W. (1950). *N.Y. St. J. Med.*, **50**, 541.
Ziffren, S. E., Zager, L. L., and Cullen, S. C. (1950). *Geriatrics*, **5**, 252.

Last year at the Conference on Military Medicine, held at Vichy under the auspices of the Office International de Documentation de Médecine Militaire, delegates discussed the problem of the rôle of medicine in biological warfare. A questionnaire was sent out, and replies have been received from more than 20 countries. A tentative answer to the problem, reported at the conference held in Brussels last month, is to separate curative medicine from experimental research work. The conference voted in favour of referring the whole question of international medical law to international authorities, particularly W.H.O., in order to clarify the position of doctors in peace and in war. Other questions discussed included the identification of medical and religious personnel in the armed Forces, and the minimum numbers of doctors, nurses, chaplains, etc., essential in prisoner-of-war camps.

PSITTACOSIS IN A FAMILY

BY

B. K. ELLENBOGEN, M.D., M.R.C.P.

Senior Medical Registrar, Sefton General Hospital, Liverpool

AND

C. M. MILLER, M.D., M.R.C.P.

Consultant Physician, Sefton General Hospital, Liverpool

Psittacosis in man was for long regarded as a serious disease with a high mortality rate and as being transmitted by parrots. It has, however, become increasingly recognized that a milder form exists, and the virus has been isolated from other birds, notably pigeons (Andrewes and Mills, 1943), domestic fowl, and fulmar petrels. Frequent references to the milder form have been made by American workers: Brown (1943), quoting a series of 45 cases of atypical pneumonia, stated that the psittacosis virus had been isolated in 10. Cases of this type seem to have been rarely described in this country (Thomson, 1930, 1943). The following two cases occurred in one family.

Case 1

A general labourer aged 16 was admitted as a case of pyelitis on May 15, 1951. He gave a history of general malaise, anorexia, headache, nausea, and severe lower back pain for the previous three days and had fainted on the day of onset. When seen these symptoms were still present, and in addition he complained of pain in the eyes on attempting to read. There was no cough or recent upper respiratory infection.

On examination he was flushed, the tongue coated, and the temperature 102.6° F. (39.2° C.). There were no palpable glands and no renal tenderness. Nothing abnormal was found in the abdomen or in the respiratory, cardiovascular, and nervous systems. The urine was normal on ward testing. This clinical picture suggested the possibility of influenza or virus pneumonia.

Meanwhile, "sulphatriad," 1 g. six-hourly, had been started by the admitting officer, and this was continued until May 18, a total of 11 g. having been given. On that day the patient was still febrile and complained of frontal and occipital headache. Examination then revealed a small area of distant bronchial breathing, bronchophony, and whispering pectoriloquy about 2 in. (5 cm.) below the inferior angle of the left scapula, with some inspiratory rales. An x-ray film showed partial consolidation of the left lower zone of "ground glass" appearance. At this stage the diagnosis of virus pneumonia seemed highly probable and Q fever was considered. Chloramphenicol, four capsules (1 g.) thrice daily, was given and the temperature fell to normal within 24 hours, the patient's general condition improving simultaneously. After three days no clinical abnormalities were detected in the lungs and the drug was discontinued (total 11 g.), the temperature remaining normal. An x-ray film on May 25 showed partial clearing, and a repeat film on June 14 showed resolution to be complete.

Laboratory Investigations.—Urine (May 16): no pus, blood, or casts in centrifuged specimen. White-cell count (May 17): 8,850 (polymorphs 72%, lymphocytes 23%, monocytes 4%, eosinophils 1%). Serology (May 18): no cold agglutinins detected in the serum. Virus complement-fixation test:

	May 22	June 5
Psittacosis/lymphogranuloma venereum ..	<1 in 10	1 in 160
Influenza A and B	<1 in 16	<1 in 16
Q fever	<1 in 16	<1 in 16
Streptococcus MG	<1 in 10	<1 in 10

Skin tests (June 23): Frei antigen, positive; psittacosis antigen, negative; control, negative.

Case 2

A factory worker aged 43 (the father of Case 1) was admitted on May 30, 1951, giving a five-days history of general malaise, nausea, headache, sweating, and shivering. He also complained of pain in the right lumbar region associated with slight frequency and scalding of micturition; these symptoms led the admitting officer to regard him as possibly suffering from acute pyelitis. Physical examination at this stage was negative except for some vague tenderness in the right lumbar region, the temperature being 99.4° F. (37.4° C.), and the urine containing no abnormalities on ward testing.

He was given 2 g. of sulphatriad at once and then 1 g. six-hourly. The next day he was still complaining of headache but had no urinary symptoms. By June 1 the morning temperature had risen to 101° F. (38.3° C.) with deterioration in his general condition. Examination revealed a small area of persistent crepitations in the left axilla. In view of his son's recent illness the question of virus pneumonia (possibly Q fever) was seriously considered, but sulphonamide therapy was continued. A chest x-ray film at this stage revealed left mid-zone consolidation, but absence of cough or other respiratory symptoms was noteworthy. On June 3 the temperature was normal, although crepitations were still audible in the left axillary region. By June 8 the chest was clinically clear, sulphatriad was discontinued (total 39 g.), and the patient allowed up. A chest x-ray film on June 4 showed clear lung fields.

Laboratory Investigations—Urine (May 31): Nothing abnormal on chemical testing or in deposit; culture sterile. White-cell count (May 31): 11,300; the pathologist reported that the film showed slight polymorphonuclear leucocytosis. Serology (June 2): Cold agglutinins detectable in patient's serum to titre of 1 in 4 (negative). Virus complement-fixation test:

	June 5	June 23
Psittacosis/lymphogranuloma venereum ..	< 1 in 10	1 in 80
Influenza A and B	< 1 in 16	< 1 in 16
Q fever	< 1 in 16	< 1 in 16
Streptococcus MG	< 1 in 10	< 1 in 10

Skin tests (July 26): Frei antigen, negative; psittacosis antigen, negative; control, negative.

Discussion

The complement-fixation tests leave no doubt that these patients were suffering from psittacosis, for, as Smadel (1943) has pointed out, a rising titre of antibody is essential for the diagnosis, and this was obtained. In Case 1 the titre rose from less than 1 in 10 to 1 in 160, and in Case 2 from less than 1 in 10 to 1 in 80, the antibody response to other agents remaining constant over the same period. The viruses of psittacosis and lymphogranuloma venereum are related antigenically; this explains the positive Frei test in Case 1. The negative skin reactions to psittacosis antigen do not vitiate the diagnosis in view of the other evidence. Inquiries were made for possible sources of infection, and it was disclosed that both patients had been in recent and frequent contact with domestic pigeons kept by a relative. Furthermore, two weeks before the onset of the son's illness it had been noticed that one of the birds was sick. Although proof that the pigeons were the immediate source of infection has not been obtained, this is a reasonable assumption. It is therefore worth re-emphasizing that in cases of obscure pyrexia, especially those associated with minimal pneumonic signs, inquiry for contact with birds should not be confined to those of the psittacine group, but should be extended to include domestic pigeons.

From the clinical point of view the absence of cough and sputum throughout the course of the illnesses was noteworthy; thus without careful attention to the chest the possibility of virus pneumonia might not have been considered and the condition might have been overlooked. This complete

absence of respiratory symptoms is unlike both the psittacine form of the disease (Sturdee and Scott, 1930) and atypical pneumonia of other cause, the latter usually being accompanied by a "distressing unproductive cough" (Coope, 1948), but no inferences need be drawn from these features.

In their work with experimental animals Hurst *et al.* (1950) showed the psittacosis virus to be sensitive to terramycin, aureomycin, chloramphenicol, and penicillin in that order; sulphonamides are known to have little or no effect. It is not easy to be certain when spontaneous remission occurs or when recovery results from treatment, but in Case 1 sulphonamides had had no effect on the temperature or general course of the illness for 72 hours, whereas the administration of chloramphenicol was associated with much general improvement and a fall in temperature within 24 hours. In our view this was probably not fortuitous but a specific effect of the antibiotic. Case 2, which was treated with sulphonamides alone, showed an increase of temperature over the first 48 hours, with deterioration of the general condition followed by a gradual decrease of temperature, but the normal was not attained until the fifth day of administration (total 18 g.), by which time he was symptomless. It is questionable whether sulphonamides accelerated his recovery or merely acted prophylactically against possible secondary invaders. However, it would be unwise to draw conclusions from the experience of two cases, and we simply record our impressions.

Summary

Psittacosis of a milder type is reported as occurring in a father and son, the disease being contracted from contact with presumed infected domestic pigeons. Absence of respiratory symptoms was an unusual feature. Chloramphenicol administered to one of the patients appeared to have an immediate beneficial effect. It is re-emphasized that in the diagnosis of virus pneumonia it is important to inquire whether there has been contact with birds other than those of the psittacine group.

Our thanks are due to Dr. J. C. Brundret for the pathology reports and to the Virus Reference Laboratory, Colindale, London, for the complement-fixation tests and the provision of the antigens for skin testing.

ADDENDUM.—Since submitting this article proof of psittacosis infection amongst the pigeons has been obtained. Samples of blood taken from three adult pigeons chosen at random from the stock were submitted for complement-fixation tests against psittacosis/lymphogranuloma venereum. This proved negative at a dilution of less than 1 in 10 in two of the samples, but the third was positive at 1 in 40.

REFERENCES

- Andrewes, C. H., and Mills, K. C. (1943). *Lancet*, 1, 292.
Brown, J. W. (1943). *Ibid.*, 1, 431.
Coope, R. (1948). *Diseases of the Chest*, 2nd ed. Livingstone, Edinburgh.
Hurst, E. W., Peters, J. M., and Melvin, P. (1950). *Brit. J. Pharmacol.*, 5, 611.
Smadel, J. E. (1943). *J. clin. Invest.*, 22, 57.
Sturdee, E. L., and Scott, W. M. (1930). Ministry of Health Report No. 61. A Disease of Parrots Communicable to Man (Psittacosis). H.M.S.O. London.
Thomson, A. P. (1930). *Lancet*, 1, 396.
— (1943). *Ibid.*, 1, 384.

The general condition of schoolchildren remains satisfactory, says Dr. J. F. Galloway, Wolverhampton school medical officer, in his annual report. There were 25,750 pupils on the school register, and one medical officer to 8,583 pupils, one nurse to 3,218 pupils, and 6,163 pupils per dental operator. He states that 42.3% of the pupils had school meals during the year and 83.2% school milk. Clothing, he says, was generally satisfactory, but what was not satisfactory was that about 8.6% of all children, mostly girls, at some time during the year showed signs of infestation in their hair. It was difficult to improve matters, since the nurses had so much to do.