The results of examining bacteriologically the stools of 234 home contacts are given: 78 (33.3%) were found to be symptomless carriers.

Spread of infection within the schools is described. Apart from the original infant school affected, direct spread within the classroom did not appear to have been the usual mode. The home played a greater part in spreading infection.

Results did not suggest that the density of infection was greater in dirty than in clean homes.

I thank Dr. J. F. Warin, medical officer of health, City of Oxford, for much help and advice during the investigation, and for criticism of this report. My thanks are also due to the staff of the Public Health Laboratory, Oxford, who carried out the bacteriological examinations.

REFERENCES

- Bloch, E. (1938). *British Medical Journal*, 1, 836.
Charles, J. A., and Warren, S. H. (1929). *Lancet*, 2, 626.
Laws, J. J. (1936). *Ibid.*, 1, 192.
Manson-Bahr, P. (1943). *The Dysenteric Disorders*, 2nd ed. London.

AN OUTBREAK OF TRYPANOSOMIASIS

BY

R. G. DAVIES, M.D., D.P.H.

*Deputy County Medical Officer of Health, Denbighshire;
Late Medical Officer, Kenya*

In December, 1948, a male African was admitted to the Native Hospital, Kisumu, in the Nyanza Province of Kenya, complaining of fever, headache, and general malaise. His temperature was 102° F. (38.9° C.), and in the posterior triangle of the neck several glands were enlarged to about the size of a hazel-nut. On examination of a blood slide I saw what appeared to be typical trypanosomes. The man was a railway employee from Kibigori, a village some 20 miles to the east of Kisumu. With the exception of three days in August, 1948, when he had returned to his home some 50 miles to the west of Kisumu, where sleeping sickness was known to exist, he had been continuously resident in Kibigori for nine years.

Although human trypanosomiasis was not known to exist at Kibigori, the tsetse fly *Glossina palpalis* was present along the whole course of the Nyando River, on which Kibigori stands, and its tributaries, from a point about 15 miles east of Kisumu. The Nyando River is about 75 miles long and enters Lake Victoria near Kisumu.

The Veterinary Department was at the time making a tsetse fly survey of the area in connexion with trypanosomiasis in cattle. Because of this case and the presence of *G. palpalis* at Kibigori, it was decided to investigate the area for further cases of human trypanosomiasis.

Present Investigation

The initial investigation was made on the station platform at Kibigori. Africans in the vicinity were persuaded to come to the station, where they were examined for the presence of enlarged axillary or cervical glands. When enlarged glands were found they were punctured and the gland fluid was examined under the microscope for trypanosomes. In this way some 15 cases were discovered in about a week. It became evident, therefore, that it would be necessary to extend the investigation over the whole area of the Nyando basin; this included two native reserves, European- and Indian-owned farms, and some newly established gold mines.

The investigating team consisted of two African laboratory attendants, an African dresser, and myself. The co-operation of the local chiefs in the native reserves was readily obtained, and with their assistance their people were persuaded to come to prearranged points for examination. The position as regards the examination of African farm labour was more difficult. It was necessary to address several farmers' meetings in order to explain the purpose of the investigation and to obtain their co-operation so that their native labour would come forward in sufficient numbers and at such times as to make the investigation practicable. A prearranged schedule was arranged with the farmers' representatives, and the team went from farm to farm examining the units of labour on each.

In all cases the method of examination and diagnosis was by palpation of cervical and axillary glands, gland puncture, and examination of the gland fluid directly under the $\frac{1}{4}$ -in. (4-mm.) objective of the microscope. The trypanosomes were recognized by their movement. This was found to be the simplest and quickest method, although it became tiring if a large number of enlarged glands were found. While the gland puncture was being done a blood slide was taken, numbered, and later examined—usually in the evenings. Staining a smear from the glands was one of the other methods tried, but this was found to take too long and to be too difficult to carry out in the dusty places where it was necessary to work. Lumbar puncture for C.S.F. cell counts and Pandy's test was also carried out, but the difficulties of doing this in the field, and in particular the adverse propaganda which it provided, soon ruled it out as a diagnostic measure.

At the beginning of the survey, if a patient was found with a positive blood, a sample of the blood was taken and injected intraperitoneally into a rat. In the laboratory the trypanosome was found to cause chronic infection in white rats, with short stumpy forms in the peripheral blood. These blood samples were discontinued after the trypanosome had been propagated in the laboratory. In each positive case the patient was questioned closely about his symptoms, which were often described with a characteristic vagueness. Some complained of no symptoms whatsoever, whilst others complained of many symptoms—lassitude, fever, headache, loss of appetite, vague skeletal pains, and general malaise. Most of them, however, had only transient attacks of fever and malaise. On general examination about one-third showed moderately severe enlargement of the spleen. This could not be considered unusual in an area where malaria was hyperendemic. About one-quarter of the cases showed some rise in temperature, and at least one-half had some form of rash, usually scabies.

In an area such as this, where many tropical diseases—malaria, bilharziasis, worm infestation, etc.—were endemic, no great reliance could be placed on ordinary signs and symptoms to form a single diagnosis. The character of the axillary glands, and especially the posterior cervical glands, was the most distinctive point. These glands had a characteristic rubbery consistency; they were single, were about the size of a hazel-nut, were mobile, and could be grasped between the fingers and thumb. After a little experience diagnosis could almost be made by the character of the glands alone, but confirmation was always made by gland puncture and demonstration of the trypanosome under the microscope. A few cases were found which gave a negative gland puncture but showed a positive blood film. These people were traced and told to report for treatment.

As a result of taking careful histories and obtaining details regarding movements with such accuracy as it was possible to obtain from Africans, it became evident that the main focus of infection was an area around Kibigori, with a small isolated subsidiary focus some 15 miles away.

Treatment

Treatment of cases was started at once, and throughout the survey weekly sessions of treatment were held every Tuesday at Kibigori, Tuesday being market day. As each

case was diagnosed 0.5 g. of "antrypol" was given intravenously and the patient told to report for treatment at Kibigori on the Tuesday following. After the initial dose, weekly injections of 1 g. were given up to a total of 9.5 g. for the adult male, and proportionate doses for women and children. It was hoped to follow this course of antrypol with a course of tryparsamide, but the adverse propaganda effect of telling people they were required to attend for a further six weeks was so great that it was decided to rely on the antrypol alone. Little trouble was experienced in getting the Africans to attend for their first three or four injections, but after that, as their symptoms cleared up, it became increasingly difficult to persuade them to complete the course of treatment. This problem was not made any easier by the attitude of certain employers who withheld rations and pay on the days the employees attended for treatment. All cases living far from Kibigori were persuaded to come as in-patients at the Native Hospital, Kisumu.

Results

The investigation was completed just after the rains started in the middle of April, by which time all the farms had been visited as well as the native reserve areas. We examined 362 Europeans and Asians, and 11,533 Africans; 549 of the latter were suspect clinically and were further tested, with the results shown in the table.

	No. Found Positive			Total
	By Gland Only	By Gland and Blood	By Blood Only	
African males ..	73	6	4	83
" females ..	46	2	—	48
" children ..	27	—	—	27
Total ..	146	8	4	158

After the end of the rains a brief re-survey was carried out at Kibigori and at Songhor, the European farming district adjacent to Kibigori. Bush clearing had already begun at Kibigori, but reports from the Tsetse Control personnel seemed to show that the rains had tended to increase and spread the fly in certain areas. This re-survey showed that the disease was still active in the Kibigori area, and indicated the possible existence of a new small focus of infection. The figures for the re-survey were:

No. of Africans examined	1,893
No. of gland punctures done	81
No. of cases of trypanosomiasis found	43

The distribution of all the cases found was as follows:

Kibigori	134
Central Nyanza Reserve	24
Nandi Reserve	6
Farm areas	37

At the end of their course of treatment all cases were re-examined. Patients were regarded as cured if their glands had disappeared, or were fibrosed and negative on puncture, and if they also had a negative blood slide. Of the 201 cases of trypanosomiasis, 105 were treated and cured at Kibigori, and 18 at the Native Hospital, Kisumu; 6 patients died during their course of treatment, probably owing to intercurrent infection. Of the remaining 72 cases, 51 were defaulters after receiving anything from one to six injections, whilst the remainder were still under treatment when I handed over the control of treatment and investigations, which had by then become standardized.

Comment

During the course of the survey, through conversation with some of the very old Africans, the fact came to light that Kibigori had been the centre of a serious epidemic of human trypanosomiasis before the 1914-18 war. Owing to the suppressive measures taken before 1914 or to unknown natural factors, the disease had completely died out between that epidemic and the present outbreak, although the tsetse fly had apparently remained throughout the entire district.

The fly had apparently not been reinfected in spite of much movement of people through the area from regions where sleeping sickness was, and still is, endemic.

Although there was continual movement of the local population within the area itself, especially to and from Kibigori, which is the hub of the district, the fly in the Nyando basin seemed to be infected in only two places, Kibigori itself and at a small site some 15 miles away.

Nearly all cases seemed to be at an early stage. In no instance were glands found which exceeded the size of a hazel-nut. In only two cases was there any suggestion of involvement of the nervous system, and this consisted merely in some marked lethargy and some debility without any definite neurological signs. Both these cases were treated at the Native Hospital, Kisumu; they showed much improvement with antrypol; and after tryparsamide a complete cure resulted, both clinically and from examination of the C.S.F.

Trypanosomes were seldom found in the blood when gland puncture was positive, and even when found were few in number. This, combined with the result of the tests on rats, seemed to show that the infection was caused by *Trypanosoma gambiense*.

No cases were found in Europeans or Asians. The youngest African patient was about 6 years old. Compared with the numbers examined there was a higher proportion of cases among women than among men, perhaps because women went more often to the river to draw water.

With one exception there was no severe reaction to antrypol. The first case of all was over-treated, receiving 10 g. in less than five weeks. A severe exfoliative dermatitis and an acute nephritis developed, but subsequently a complete recovery was made. One case also developed an antecubital cellulitis following an injection of antrypol; recovery followed the application of moist heat.

I am indebted to the Honorary Director of Medical Services, Kenya, for his help and encouragement, and to Dr. W. W. Wilkinson, lately S.M.O., Kenya, for his most valuable advice during the survey.

DIAGNOSIS OF HUMAN TRYPANOSOMIASIS

BY

G. F. T. SAUNDERS, M.D.

Specialist, Colonial Medical Service, Gold Coast

While Cates and McIlroy (1951) showed clinical alertness in suspecting trypanosomiasis in a rather atypical case seen in England, their article indicates that routine methods of laboratory diagnosis are not well known there. Details of the methods developed over a number of years by the Gold Coast medical field units might therefore be of use to home practitioners. These units include mobile teams who examine entire populations and fixed centres at which cases attend. Cases found by the mobile teams are usually less advanced than those who come to the centres.

Blood Films

Trypanosomes may or may not be found at any stage of the disease. About 33% of the cases diagnosed by the mobile teams are found positive in one thick drop, but only 16% of those who attend at the fixed centres. Trypanosomes are most easily found in fresh blood examined with a 1/6-in. (4-mm.) objective; but if this method is inconvenient a thick film stained with 1:10 Giemsa for twenty minutes and examined with the oil-immersion objective will show them. It is best to begin at one corner and work through the entire film line by line on the mechanical stage.