2. Treatment is instituted within approximately four hours of wounding, but clinical judgment may extend this arbitrary period.

3. Operative conditions are adequate and the correct instruments and suture materials available.

4. An atraumatic technique is employed.

When these conditions are satisfied primary suture may be employed for all extensor tendon divisions.

Primary suture may be used for division of the flexor profundus beyond the level of the sublimis insertion, the

FIG. 3.—Types of flexor tendon division showing suitable methods of repair. A, Primary suture is permissible if general conditions are suitable; otherwise suture skin only and perform tendon graft later. B, Unsuitable for primary suture; suture skin only and perform tendon graft later. C, Primary suture is permissible if general conditions are suitable; otherwise suture skin only and perform secondary suture, tendon transfer, or free graft later. D, Primary tendon suture is always advisable if conditions permit. Perform nerve suture later.

flexor pollicis longus in its distal part, and the tendons in the palm, and immediate repair is obligatory for divisions at the wrist level. It is not advisable to perform primary or secondary suture of tendon divisions between the distal palmar crease and the sublimis insertion, for replacement by a free tendon graft at a later stage gives more uniformly satisfactory results.

As in all surgery of the hand, fine instruments and a delicate technique in a bloodless field are essential requirements in tendon surgery. The suture material most commonly used is stainless steel wire in single strand, gauge 40, or the twisted form originating in

Sweden. The wire is attached to a small straight or curved cutting needle. Stainless steel wire causes less reaction than any other suture material and may remain in the tissues indefinitely without ill effect. Satisfactory haemostasis must be obtained before the wound is closed, as haematoma is one of the causes of deep fibrosis and stiffening of the hand. If adequate haemostasis is secured there is no need to use any form of wound drainage.

Bone and Joints

There is much to be said in favour of the present tendency to use internal fixation, whenever possible, for metacarpal and phalangeal fractures associated with severe lacerations. The need for difficult and restrictive external splintage is avoided, and more attention can be given to the careful repair by grafts or flaps of any accompanying skin loss. Also there is not the same risk of late loss of position requiring further manipulation and disturbance of the healing soft tissues.

Internal fixation may be achieved by the use of carefully fashioned intramedullary bone-pegs taken from the tibia, ileum, or ulna, or by transfixion of the fragments by fine Kirschner wires.

Amputations

The necessity to amputate digits does unhappily arise in cases of extensive lacerations, but it is better to err upon the side of conservative rather than radical surgery. It needs considerable experience to decide whether a severely damaged finger can be restored to usefulness or whether it will eventually become an embarrassment. In our desire to preserve all living tissue in the hand we must not lose our sense of proportion and insist upon preserving digits which cannot be of any use.

When several fingers or the thumb are injured it is wise always to be conservative, because even a part of a digit may be of value to the mutilated hand.

Splintage

After any injury, by accident or by surgery, the tissues must be allowed time to rest and recover. The hand is supported by a compression bandage composed of gauze waste, wool, and crêpe bandage carefully and deliberately applied to give even pressure. The tips of the digits are always left exposed for inspection.

In many instances this bandage will be sufficient splintage, but it is often helpful to apply a simple plaster cover to hold the wrist in slight dorsiflexion. The fingers and thumb, unless there is some special contrary reason, should always be in the position of rest, with the thumb in moderate opposition and the fingers semi-flexed.

The hand, after operation, should be supported in an elevated position for 24–48 hours to assist in the vascular drainage and in the avoidance of oedema.

Next Refresher Course Article.—"Rheumatic Fever," by Professor C. B. Perry.

BICENTENARY OF THE MANCHESTER ROYAL INFIRMARY

BY

WILLIAM BROCKBANK, T.D., M.D., F.R.C.P.

The Manchester Royal Infirmary first opened its doors on July 27, 1752, 200 years ago, in a little house in Garden Street, off Withy Grove, on the site now occupied by the Co-operative Wholesale Society's building. The house held 12 beds and was staffed by a matron, three physicians, three surgeons, and a secretary. It was called the Publick Infirmary. The government was placed in the hands of trustees. These were persons who had given to the hospital 20 guineas or upwards at one time or were subscribers of 2 guineas or upwards annually.

On the opening day one out-patient presented himself for treatment. A week later the number had increased to five. The first in-patient, a boy of 12 with ulcers on his legs, was admitted on August 3. The number of patients rapidly increased, so that by the middle of October the matron had to be provided with a nurse and a servant. In April, 1753, a room was set aside for an apothecary's shop. In this way began the dispensary, which proved to be a most important feature of the hospital. At the end of a year's work 75 persons had been admitted and 249 out-patients treated, at a total cost of a little more than £400.

At the start of the second year the adjoining house was taken over and equipped with 10 beds. More important was the decision to build a hospital elsewhere, on land now known as the Piccadilly site, situated in open country at the upper end of Market Street Lane. Plans were drawn up for the erection of a building to contain 80 beds for patients, "with proper convenience for nurses and servants and including a hot bath." This "New Infirmary" was brought into use in June, 1755, with 50 beds. The two houses in Garden Street had served their purpose for nearly three years, during which time 374 in-patients and 1,629 out-patients had been handled.

The Manchester Infirmary in 1829, with the adjoining lunatic asylum, on the Piccadilly site.

Having triumphantly launched the infirmary, the board flanked it with a lunatic asylum in 1766 and a complete suite of public hot, cold, and vapour baths in 1780. Persons disordered in their senses were promised "that humane treatment and judicious advice that they had no encouragement to expect in private madhouses." It was also necessary to make a substantial extension to the hospital. It is impossible to stress too greatly the effect of the infirmary on the life of the town at the end of the eighteenth century, due to such far-reaching ideas as the provision of public baths, the introduction of a public health service, the prevention of smallpox, the condemnation of slums, the plea for control of lodging-houses, the investigation of cases of cancer, and the concern about the prevention of accidents in the factories, all in addition to the care of the sick and of those mentally afflicted.

During the first half of the nineteenth century repeated extensions were made, until in 1858 the old infirmary attained its final shape, being surmounted by a dome holding a public clock. The lunatic asylum was transferred to Cheadle and the baths were closed. The final extension was expected to provide an additional 180 beds, but in the calculation the beds had been placed too close to each other. In fact, the gain was only 85, making a total of 277. While these alterations were taking place the infirmary passed its centenary. Since its doors were first opened 940,000 patients had been treated, one-ninth of them in-patients.

Hospital Fever

Hospital fever was a constant worry, and it was hoped that the new wards would lessen its ravages, but this was not the case. To make matters worse the board decided to admit cases of infectious fevers to one of the wings. The routine work of the hospital was maintained by placing the beds closer together until they totalled 310. The inevitable result was a sharp rise in the mortality figures.

A convalescent home was started at Cheadle in 1867 and a fever hospital at Monsall in 1871. But the wards continued to remain unhealthy. The board in desperation called for an independent investigation into the condition of the hospital. The report, dated 1876, disclosed a sorry state of affairs, particularly with regard to the drainage. The modern reader must agree that the only thing to be done was to pull down the building and start afresh elsewhere. There was a suitable plot of land close to the growing university, then known as Owens College. The board proposed this course of action to the trustees, but the trustees would not agree, preferring that the hospital should remain in the centre of the town rather than in its "damp and foggy suburbs." Thus perished the first attempt to move the infirmary from the Piccadilly site. Thirty-two years were to pass before this was finally accomplished.

The board now determined to rebuild and extend the hospital. There was a good deal of land available around the building, but the civic authorities wanted the site for a town hall. Generous terms were offered, but the board could not accept. The city council therefore decided to prevent any encroachment on land not already used for building. The authorities of Owens College endeavoured to persuade the board to move the hospital to Oxford Road, and when negotiations fell through they sided with the city council and effectively blocked successive schemes brought forward by the board. These bodies so influenced the trustees that in 1902 they agreed to the policy that there should be no rebuilding of the infirmary on the Piccadilly site, and ordered that the building and site should be sold on suitable terms and that a new infirmary should be built elsewhere without delay. Thus ended a long and bitter controversy. The members of the board resigned in a body and were replaced by trustees in favour of the new scheme. Within six years the infirmary as we now know it had been designed and built. It was ready for occupation in the autumn of 1908, and on a raw foggy December morning the patients were transferred to the new building. No time was lost in demolishing the old infirmary, its place being taken by a lovely garden where the townsfolk can sit and rest in the centre of the busy city. May it always remain thus.

The New Building

The new building, designed to hold 500 beds, with accommodation for an additional 100 if needed, was formally opened on July 6, 1909, by their Majesties King Edward VII and Queen Alexandra. The chairman of the board of management (Mr. William Cobbett) read an address of welcome and was knighted on the spot. Thereafter the board pursued through peace and war a remarkably progressive policy. Even the extensive damage wrought in two air raids in 1940 was not allowed to interfere, and in the darkest days plans were laid for the big developments which subsequently took place.

Many famous men have served on the medical staff of the infirmary. Charles White is renowned in the history of midwifery. Thomas Percival's medical ethics are still well known and respected. John Ferriar did much to develop public health services in the rapidly growing township. Peter Mark Roget wrote the *Thesaurus* beloved by addicts of the crossword puzzle. Edmund Lund introduced Lister's principles into the hospital. Walter Whitehead left his name in the annals of surgery. William Thorburn was a pioneer in neurosurgery. Graham Steel and James Ross made outstanding contributions in cardiology and neurology respectively. R. T. Williamson's book on the spinal cord is a classical work and is still in use. Lloyd Roberts, William Milligan, Archibald Donald, and Alfred Barclay developed their specialties within its walls. Arthur Burgess showed the practical possibilities of dealing surgically with acute abdominal emergencies. E. D. Telford laid the foundations on which vascular surgery has developed. John Morley, Geoffrey Jefferson, and Harry Platt have added greatly to its prestige, and the tradition is being worthily maintained. Lord Moran, Lord Webb-Johnson, and Sir William Fletcher Shaw, presidents simultaneously of the three Royal Colleges, all studied in the Manchester Medical School.

On July 5, 1948, the board of management lost control of its infirmary, which passed into the care of the board of governors of the United Manchester Hospitals under the chairmanship of Mr. Colin Skinner, who had been its treasurer for nearly 20 years. No happier or better appointment could have been made. Since the appointed day there have been many serious problems, including the amalgamation of the five constituent hospitals and the co-ordination of the services they provide with those provided by the hospitals belonging to the regional board. In addition, the rapid advance of scientific methods in medicine has necessitated the provision of much more adequate facilities.

In spite of increasing economic difficulties steady progress has been maintained. The neurosurgical block was opened in 1950 by Lord Webb-Johnson. More important still was the approval given by the Ministry to the erection of a big new building for radiology and clinical pathology, two vital and rapidly expanding departments that are badly hampered by cramped accommodation.

Thus the Manchester Royal Infirmary has kept and is keeping pace with the astonishing advances made in medicine during the past two decades. In many fields it has led the way, and it can truly be said that the hospital has a world-wide reputation as a centre of treatment and research and as a teaching school.

A.C.T.H. AND CORTISONE IN RHEUMATIC FEVER

A preliminary report of an international study of the relative merits of A.C.T.H., cortisone, and aspirin in the treatment of rheumatic fever was made recently in Chicago. The results of the study are at present inconclusive. The work is being conducted in 13 research centres in the United States, Great Britain, and Canada by the American Heart Association's Council on Rheumatic Fever in conjunction with the Medical Research Council. British representatives on the panel which made the report were Professor A. Bradford Hill and Dr. E. G. L. Bywaters.

The plan of study provides for uniform criteria for the diagnosis of rheumatic fever and for the degree of rheumatic activity required for the admission to the study, the random allocation of patients to the three treatment groups, a defined dosage schedule of the drugs for a fixed period of time, a specified period of observation following treatment, and a long-term follow-up schedule. It also lays down precisely the frequency and type of clinical and laboratory observations to be carried out on each patient. To date, in all three countries, 658 cases have been admitted to the study, and the analysis of rather less than half of these is the basis of the preliminary report. So far no consistent pattern of response to treatment is evident, and no firm conclusions can at present be drawn concerning the drug most effective in the control of the acute illness. The cases have not been under observation sufficiently long to provide data on the prevention of rheumatic heart disease. Admission of new cases to the study will be brought to an end later this year. It is expected that a total of 750 cases will be available in all three countries for final analysis.

BRITISH EMPIRE CANCER CAMPAIGN

Delivering his presidential address to the 29th annual general meeting of the British Empire Cancer Campaign at the House of Lords on July 3, the DUKE OF GLOUCESTER announced that the Queen had graciously consented to grant her patronage to the Campaign. The new report, the Duke went on, "gives clear proof of the remarkable activity of the Campaign at home and overseas . . . the headings alone show plainly the extraordinary amount of hard work done by the council and committee of the Campaign . . . and the many devoted men and women working patiently in their laboratories and at the bedsides of the sick." The Duke of Gloucester also commented on the increasing liaison between the Campaign and research establishments in Canada and the United States, and said that the Exchange Fellowship system was "operating steadily and smoothly."

Proposing a vote of thanks, the chairman, Lord HORDER, gave as an example of the Duke's practical interest a message which he had sent to a cinema in Jersey, where a film *première* was being held to raise funds. This message crowned the success of the function, which raised £2,500 for the Campaign.

Moving the acceptance of the annual report (see also p. 84), Sir DANIEL DAVIES drew attention to a newly published book entitled *A Study of Cancer in London*, which surveys the work of all London hospitals treating

and diagnosing cancer over a five-year period ending in 1944. It covered 15,201 cases. All but 400 had been followed up. Only 29% of the cases were at a localized stage on coming for treatment; however, of 451 stage 1 cases treated by surgery and radiotherapy, 55% were still alive after five years. This once again showed the vital need for early diagnosis. With this, Sir Daniel coupled a plea for intelligent education of the general public about cancer. Seconding the report, Mr. J. C. BURLEIGH, of the Finance Subcommittee, said that the British Empire Cancer Campaign's income for the past year from donations, collections, and legacies had been nearly £260,000—over £34,000 more than in 1950.

Reports of Societies

TREATMENT OF ANURIA

At a meeting of the Section of Urology of the Royal Society of Medicine, on June 26, Dr. E. M. DARMADY opened the proceedings by reminding the audience that considerable advances had been made in the understanding of the anurias since he addressed the Section four years ago. The advances were due to a better knowledge of the pathology and natural history of the disorder, and of methods of combating the toxæmia of renal failure.

Whatever the cause of acute tubular necrosis, the most important factor was circulatory failure. The kidney was profoundly affected by anoxia, and could not remain severely ischaemic for more than six to eight hours without developing changes which were often irreversible. Dissections of the nephron by Oliver and his own group showed that the characteristic changes (irrespective of the aetiology) were disruption of the tubular wall and patchy degeneration of the tubular epithelium, which affected the upper as much as the lower nephron. There were three stages in the natural history of the syndrome, as described by Bull and his colleagues: the onset, which usually lasted about six hours after the initial stimulus, the stage of oliguria or anuria, which might last for up to 22 days, and finally a late diuretic phase during which large amounts of water and electrolytes were lost.

The causes of the precipitating circulatory failure were manifold and were difficult to classify, but might be divided into general causes (shock, trauma, etc.); reflex causes (calculus anuria, vascular injury, sulphonamide anuria, etc.); renal diseases, such as nephritis; allergic causes; and various toxins. Obstetrical accidents such as ante- and post-partum haemorrhage were also of great importance.

In treatment, splanchnic block and intravenous diuretics such as sodium sulphate were likely to succeed only if adopted at the very onset. The essence of treatment lay in the control of circulatory failure and fluid intake, and restriction of the accumulation of waste products. The last factor could be dealt with either by removal of the waste products by some form of dialysis, or by minimizing their formation. Assessment of cellular and extracellular hydration was often difficult, and, although estimation of the haematocrit and plasma specific gravity were helpful, measurement of the osmotic pressure of the plasma by the method of Hamburger was more promising. In the anuric stage overloading with salt and water was disastrous, and new methods of estimating the volume of the extracellular compartment were urgently needed. The late diuretic phase often occurred suddenly, and necessitated rapid replacement of water and electrolytes. Since hypokalaemia was likely to occur at this time it was helpful to have readily available stock solutions containing potassium (1.9% potassium lactate or 1.1% potassium chloride).

Dr. G. M. BULL said that anuria was due to three main causes: circulatory insufficiency, organic renal damage, and obstruction. He and his colleagues, Dr. A. M. Joekes and Dr. K. G. Lowe, had studied 112 cases of anuria or severe oliguria at Hammersmith Hospital from 1947 onwards: 48 were due to circulatory renal insufficiency, 59 to organic