seem most likely, with a mesenteric infarct to account for the second-day ileus. Pheochromocytoma was considered because of the glycosuria and hypertension, but, after perusal of the papers by Crowther (1951) and Swan (1951), was rejected as unlike any previous case so far reported at that time. Unfortunately, the paper of Gilliland and Daniel was not noted until the day of the necropsy, when the similarity in presentation and signs was remarked upon.

Presumably haemorrhage into the cystic, friable, necrotic tumour produced the severely shocked state which, with a curious lividity of the skin, persisted throughout the whole illness. The use of anticoagulants must have aided this considerably, as must the bout of hypertension. A troublesome feature of the case was the diabetes, more and more insulin being required to keep the urine clear of sugar as her blood pressure rose after the original fall.

I would like to express my thanks to Dr. J. C. Prestwich for permission to publish this case, which was under his care.

R. S. JELLIFFE, M.B., B.S.,
House-physician, St. Mary's Hospital, Portsmouth.

REFERENCES

- Crowther, K. V. (1951). *British Medical Journal*, 1, 445.
Gilliland, I. C., and Daniel, O. (1951). *Ibid.*, 2, 275.
Swan, H. J. C. (1951). *Ibid.*, 1, 440.

Pheochromocytoma Simulating Thyrotoxicosis

In spite of a growing literature of case reports and methods of investigation, the diagnosis of pheochromocytoma presents many difficulties, and it is probable that only a minority of cases are recognized during life. This state of affairs may be attributed to the apparent rarity of the tumour and the wide range of symptom-complexes it may produce. The tumour itself is often too small to demonstrate by physical methods, and other investigations may yield ambiguous results. The following case illustrates the similarity which may exist between thyrotoxicosis and the syndrome of hyperadrenalism.

CASE REPORT

A married woman aged 58 was referred to hospital on June 19, 1950, by Dr. G. C. Campbell, who suspected thyrotoxicosis. For 12 months she had suffered from paroxysmal vertigo associated with tinnitus and deafness on the right. More recently she had lost weight and complained of weakness and apprehensiveness. Her appetite remained excellent. There had been amenorrhoea since the normal birth of her second child 28 years before. Her previous health had been good. There was no family history of goitre and none of deafness.

Of slight build, the patient was evidently emaciated. Her weight was 5 st. 10 lb. (36.3 kg.). She wore a rather strained expression and was pale, but not obviously anaemic. Her hair was fine; the eyebrows and axillary and pubic hair were scanty. There was an impression of restlessness about her movements, which were rapid and bird-like. Other features were a resting pulse rate of 120, a blood pressure of 180/100, an apex beat in the sixth left intercostal space in the mid-clavicular line, and a loud systolic murmur heard over all areas and maximal at the apex. No exophthalmos was evident. There was a fine tremor of the outstretched fingers. The thyroid gland could not be felt. Examination of the patient's abdomen revealed no abnormalities. She was mildly breathless at rest, but no physical signs were found over the lung fields. The nervous system appeared normal apart from some nerve deafness on the right. The tympanic membranes were healthy and intact. The urine contained neither albumin nor sugar.

A threefold diagnosis of thyrotoxicosis, old rheumatic heart disease, and possibly otosclerosis was made. Mr. Arnold Gourevitch kindly saw the patient and agreed

that thyrotoxicosis seemed likely. At his suggestion an x-ray film of the thoracic outlet was taken. This showed no evidence of a retrosternal goitre.

On July 13 the patient was admitted to hospital, and on the 17th she was given methyl thiouracil, 600 mg. daily, and iodine. By July 23 she was able to say she felt much less agitated, but the pulse rate seldom fell below 100. Iodine was discontinued on July 27. By August 4 her condition was clearly not responding in the manner anticipated. Tachycardia and tremor persisted. On August 11 the pulse rose to a sustained rate of 150 and breathlessness increased. Venous engorgement appeared in the neck. The pulse rate was 170 on August 14. The pupils were constricted and the skin was moist. The blood pressure had fallen to 116/96. Methyl thiouracil was discontinued. On August 17 the patient was semi-conscious, pulse 120, and B.P. 100/70. On the 21st consciousness returned for a few hours, pulse 110. Death occurred on the 22nd.

Dr. A. G. Marshall conducted a necropsy, and I am indebted to him for the report, the positive findings of which were: "Thyroid 34 g. Smooth encapsulated surface. Cut surface showed excess colloid with no adenomata. One or two very small haemorrhages present. Lungs: gross emphysema with severe engorgement. Pericardial effusion (about 3 ml. clear yellow). Heart: no dilatation, no hypertrophy. Mitral valve moderately distorted, with thickened chordae tendineae, and clusters of small fibrinous vegetations on it. Other valves normal. Myocardium normal. Coronary arteries: double orifice of left. No atherosclerosis. Aorta: slight sclerosis. Stomach: engorged, containing altered blood. Liver: severely engorged. Adrenals: right, 6 g. normal; left, 20 g. Sections from the left adrenal show it to be composed of cords and columns of polyhedral cells with much cytoplasm and small nuclei. It is highly vascular, but there are no features to suggest malignancy. The appearance is highly reminiscent of the adrenal medulla. Unfortunately, as the tumour was unsuspected, no fixative agent containing chrome salts was available, and specific staining has failed."

DISCUSSION

In one of two cases of pheochromocytoma recorded by McCullagh and Engel (1942) a similar error of diagnosis was made and death took place shortly after thyroidectomy. In this case the diagnosis of thyrotoxicosis seemed to be further supported by the presence of a thyroid adenoma. Those authors observed that the B.M.R. may be significantly raised in hyperadrenalism.

Espersen and Dahl-Iversen (1946) describe another case in which a diagnosis of thyrotoxicosis led to treatment with methyl thiouracil. This was followed by improvement, but was discontinued because of toxic manifestations. Subsequent thyroidectomy led to a recession of symptoms for five months, when further attacks of hyperadrenalism suggested the correct diagnosis of pheochromocytoma, which was successfully treated by adrenalectomy.

In view of the similarity of the two syndromes it is surprising that this diagnostic pitfall has not been more widely published. Clinically the two conditions share many features. Of the symptoms [*sic*] of pheochromocytoma Hartman and Brownell (1949) include "palpitations, hypertension, headache, nausea, pallor, perspiration, agitation, anxiety, tremulousness, dizziness, dyspnoea, pain (precardial or epigastric, or of the extremities), vomiting, glycosuria, and sometimes prostration." In the case here reported there was considerable weight loss but no loss of appetite.

J. V. S. A. DAVIES, B.M., M.R.C.P.

REFERENCES

- Espersen, T., and Dahl-Iversen, E. (1946). *Acta chir. scand.*, 94, 271.
Hartman, F. A., and Brownell, K. A. (1949). *The Adrenal Gland*. Kimpton, London.
McCullagh, E. P., and Engel, W. J. (1942). *Ann. Surg.*, 116, 61.