Sharpey-Schafer (1946) found that myxoedema artificially induced by thiouracil in patients with chronic heart failure did not lead to a change in the work of the heart (cardiac output $\times$ mean blood pressure).

It seems possible, therefore, that the beneficial effect of induced hypothyroidism is, in part, the result of changes in the tubular reabsorption of salt and water. It is of interest, in this respect, that the patient who suffered from both mitral stenosis and spontaneous myxoedema was able to excrete salt and water normally although her renal clearances were at the same level as those of patients with low-output heart failure.

Summary

The renal circulation and the capacity to excrete salt have been studied in small groups of patients suffering from myxoedema or chronic rheumatic heart disease, and in one patient suffering from both conditions. In some patients from each of the two groups the cardiac output has been measured.

The glomerular filtration rate and renal blood flow were reduced to approximately the same extent.

In spite of the similar renal circulatory changes the patients with myxoedema were able to excrete salt and water, whereas that function was seriously impaired in the patients with rheumatic heart disease.

One patient with mitral stenosis and spontaneous myxoedema had renal clearances of the same order as the other patients studied, but was able to excrete salt and water.

The cardiac output was low in both groups, but, whereas the arteriovenous oxygen difference and the cardiac output per unit of oxygen consumed were abnormal in the patients with rheumatic heart disease, these values were normal in the patients with myxoedema.

The significance of these findings is discussed.

We are indebted to Professor C. H. Stuart-Harris for his help and for the opportunity of studying patients under his care, and to Dr. M. W. A. Gatman for many biochemical estimations.

REFERENCES

- Blumgart, H. L., Levine, S. A., and Berlin, D. D. (1933). *Arch. Intern. Med.*, 51, 866.
 Corcoran, A. C., and Page, I. H. (1947). *J. clin. Endocr.*, 7, 801.
 Davies, C. E. (1951). *Lancet*, 2, 1052.
 — and Kilpatrick, J. A. (1951). *Clin. Sci.*, 10, 53.
 Dick, A., and Davies, C. E. (1949). *J. clin. Path.*, 2, 67.
 McMichael, J., and Sharpey-Schafer, E. P. (1944). *Brit. Heart J.*, 6, 33.
 Merrill, A. J. (1946). *J. clin. Invest.*, 25, 389.
 Mokotoff, R., Ross, G., and Leiter, L. (1948). *Ibid.*, 27, 1.
 Sharpey-Schafer, E. P. (1946). *British Medical Journal*, 2, 888.
 Zarrow, M. X., and Money, W. L. (1949). *Endocrinology*, 44, 345.

An important service rendered by Post Office wireless stations on the coasts of Britain is the transmitting of medical advice to ships at sea. In the absence of a ship surgeon, illnesses among the ship's company are generally dealt with by the captain, so far as he is able, with the aid of the ship's medical chest. In serious cases where the captain feels that a doctor's opinion is desirable he may forward particulars to a coast station by radio. If suitable equipment is installed in the ship it is possible for the captain and the doctor to speak directly to each other by a combination of the radio link and the normal telephone system. These messages are sent either in plain language or in a special code devised to overcome language difficulty when dealing with foreign ships. During 1951, 469 medical messages were handled by Post Office coast stations.

THE TREATMENT OF LUMBAR INTERVERTEBRAL DISK PROTRUSION

AN ASSESSMENT OF CONSERVATIVE MEASURES

BY

R. S. HENDERSON, F.R.C.S.

Orthopaedic Surgeon, Royal Alexandra Hospital, Edmonton, Alberta; formerly Senior Registrar, Royal National Orthopaedic Hospital, and Chief Assistant, Orthopaedic Department, St. Bartholomew's Hospital

So common has become the diagnosis of intervertebral disk protrusion as a cause of backache and sciatica that it is important to emphasize the small part that operation plays in the treatment of this disorder. O'Connell (1951), in a review of 500 cases treated by operation, made it clear that most of his patients were referred to him only because long periods of conservative treatment from general practitioners, physicians, neurologists, or orthopaedic surgeons had failed. Of all patients seen in his clinic, he considered that less than 20% needed operation.

In a disorder with a tendency to natural cure a patient may receive many different lines of treatment while the natural history of the condition unfolds. The success or failure of such treatment is too often judged on a basis of trial and error. The purpose of this review is to assess, so far as is possible, the results of the various forms of conservative treatment carried out in the orthopaedic department of St. Bartholomew's Hospital over five years.

Material and Results

In all, 500 unselected patients have been observed for a period of not less than 18 months from their initial treatment. They attended the orthopaedic department between 1946 and 1951 with symptoms and signs giving rise to the diagnosis of lumbar disk protrusion. This number is only a very small fraction of all the patients with lumbar and sciatic pain who attended the hospital in that time. Those with backache but without evidence of lumbar root involvement or neurological disturbance have been excluded from this series. Of the total number, 314 were men and 186 women. Two patients have died since discharge from hospital—apparently from causes unconnected with their spinal condition; five recovered spontaneously while awaiting treatment.

The upset caused by disk protrusion is essentially subjective and the results of conservative treatment are therefore difficult to assess. In the accompanying Table the term "relieved" implies complete loss of pain and disability without recurrence during the period of review; "improved" indicates that the patient has been able to resume work or normal life. The number of patients completely relieved by treatment is very small when compared with O'Connell's (1951) figures for those treated by operation. But, as he himself said, patients requiring operation have suffered severe or prolonged disability and mild post-operative symptoms are unlikely to give cause for complaint. Conservative treatment, on the other hand, is given for all degrees of severity and rarely produces such dramatic changes as operation. Recurrence, moreover, is to be expected unless advice on such matters as the lifting of heavy weights without bending the spine and the avoidance of stooping is followed carefully.

Of the 500 patients, 406 were initially improved or relieved, and only 58 were considered eventually to require operation. Sixteen were not improved, but were deemed unsuitable for operation because of associated functional

disorders. A few were treated by epidural injection, but the number was too small to assess the value of this form of treatment.

Table of Results

Initial Treatment	Total	Re- lieved	Im- proved	No Initial Improve- ment	Recur- rence	Ultimate Improve- ment with Conser- vative Measures	Opera- tion
Rest in bed ..	123	15	72	36	33	26	43
Plaster jacket	71	14	38	19	19	33	5
Lumbo-sacral support ..	104	9	88	7	3	10	0
Manipulation	68	10	40	18	15	23	10
Exercises ..	129	34	81	14	11	25	0
Spontaneous re- covery ..	5	5	—	—	—	—	—
Total ..	500	87	319	94	81	117	58

Methods of Treatment

Rest in Bed.—Complete rest in bed for at least three weeks, preferably in hospital, was advised when the symptoms were severe or of sudden onset and root tension signs well marked. Although more than half were relieved or improved, a quarter suffered recurrence of symptoms. All those failing to show initial improvement with rest were ultimately treated by operation, as were one-third of those whose symptoms recurred. Continuous leg traction was not used in this series.

Plaster Jacket.—This form of immobilization was used when rest in bed was indicated but economically impossible, or sometimes to prolong rest when improvement was slow. Lasting improvement was obtained by plaster alone in about half the patients treated. This corresponds with the findings of Durbin (1948). Jackets were initially applied in the position of the deformity and subsequently changed if necessary.

Lumbo-sacral Support.—Usually patients were fitted with supports when their symptoms were mild and when their lives and work involved activity which made recurrence probable. Old people, child-bearing housewives, and policemen were the commonest candidates for this form of treatment. Twice as many men as women were fitted with spinal supports. Experience showed that, once fitted, very few were able to leave off their instruments and most patients were content to depend on them indefinitely.

Spinal Manipulation.—Patients treated by this method fell into two groups. Of 15 patients with spasm and signs of root tension only three were relieved or improved, and 10 eventually came to operation; despite the alleged dangers of paraplegia (Goldthwait, 1911), in no cases were symptoms or signs made worse. During manipulation, movements were always carried out gently and flexion was avoided. Though manipulation is not now generally advocated for the acute type of case, it is interesting to read old case records of "sciatic scoliosis" in the days before the real cause of this condition was recognized. Though movements, including full flexion, were then always carried out strongly, there were apparently no disasters and some dramatic and lasting cures. Of the second group, those with stiff backs, little root involvement, and a long history of moderately severe symptoms, more than half were relieved or improved. In addition to those recorded as being treated primarily by manipulation in this series, 20 patients were not improved by physiotherapy or exercises until after manipulation had been carried out. They had already undergone several other forms of treatment.

Extension Exercises.—It has long been realized that no form of treatment is complete unless exercises to strengthen the extensor muscles of the spine are taught at the same time. In an acute attack exercises are often not at first possible, since any exertion aggravates the pain. In this series, 129 patients with mild symptoms were primarily treated by extension exercises with or without some other

form of physiotherapy. It will be seen that over 90% of the patients were improved or relieved without any other treatment. O'Connell (1951) considered that exercises should be designed not only to strengthen the extensor muscles but also to restore full mobility.

Discussion

The number of patients with disk protrusion attending hospital to-day far exceeds those treated in the years covered by this review, no doubt because both doctors and patients are now "disk-conscious." Accurate diagnosis having been established by investigation, the main problem is to select the right treatment at the right time and reduce the period of disability.

The present investigation shows that for patients with severe root involvement the correct advice should be complete rest followed by graduated exercises. For mild symptoms, early institution of back exercises and physiotherapy is probably the right treatment. The majority of cases, however, lie between these extremes, and it is in these, with moderate disability, that the decision regarding immobilization is more difficult.

The effect of prolonged immobility may lead to "fibrous ankylosis" of the spine at the affected level (Crisp, 1945), while the protrusion becomes organized; or immobility may sooner or later promote adhesions between disk tissue and lumbar roots (as are often seen at operation) and in the periarticular tissues. If the protective spasm that so often obliterates the normal lumbar curve becomes fixed by such adhesions after long immobilization, the spinal extensor muscles will no longer be able to act at a mechanical advantage. As a result these muscles will waste and lose tone in the same way as the quadriceps when knee extension is lost. As the spine thus loses its natural brace a vicious circle will be brought into being, with increasing liability to strain and recurrent symptoms. This is the state of affairs for which manipulation, aiming to restore extension and rotational mobility, may be expected to be of most value. The presence of these adhesions becomes evident when passive extension of the spine in the prone position causes pain during routine examination. If their presence is not recognized many patients may find themselves with the burden of a supporting corset which will only partially relieve their symptoms and which they will never be able to discard.

Of the patients in this series, 75% were between 30 and 50 years of age; 15% were under 30, and only 10% over 50. Of those fitted with spinal supports nearly half were under 40. Though it is clear from the records that young patients were treated in this way only to allow them to carry on their work, it is suggested that generally these instruments should be reserved for old people with degenerative changes in their lumbar spines. There would seem to be a grave danger to-day that large numbers of active men and women are being condemned more or less indefinitely to these corsets, when more active measures might give greater relief.

Summary

A review is made of a series of 500 unselected cases of lumbar intervertebral disk protrusion observed over a period of not less than eighteen months from the starting of treatment.

An attempt has been made to assess the value of, and indications for, the various measures used in conservative treatment over a five-year period.

I should like to thank Mr. S. L. Higgs, Mr. H. Jackson Burrows, and Mr. W. D. Coltart for their help and permission to make use of case records of their patients, and Mr. J. E. A. O'Connell for his kind advice.

REFERENCES

- Crisp, E. J. (1945). *Lancet*, 2, 422.
 Durbin, F. C. (1948). *J. Bone Jt Surg.*, 30B, 487.
 Goldthwait, J. E. (1911). *Boston med. surg. J.*, 164, 365.
 O'Connell, J. E. A. (1951). *J. Bone Jt Surg.*, 33B, 8.