REFRESHER COURSE FOR GENERAL PRACTITIONERS

NEPHRITIS—II. PROGNOSIS AND TREATMENT

BY

ROBERT PLATT, M.D., M.Sc., F.R.C.P.

(Professor of Medicine, University of Manchester)

Prognosis in Type 1 Nephritis

In prognosis we quickly become aware that what may be significant to the statistician is quite inadequate for the clinician, who is called upon to deal with individuals rather than "populations at risk." What the parents of a child with acute nephritis want to know is not the overall recovery rate but whether their particular child is going to be in the 80% who recover or the 20% who do not. Of course it is important for both doctor and relative to know that the general chances of recovery are high, but by taking note of the course of events in the individual we can often get a good deal further.

Cases with an acute onset, however ill in the first stage, even with convulsions and anuria, often make a complete recovery. Severity of onset is certainly no indication of bad prognosis. Very often, after about a week or a fortnight of oliguria and oedema, there is quite a sudden change for the better. The urine volume, which may have been only 300 to 500 ml., suddenly increases, the oedema rapidly subsides, the puffy bloated appearance of the face disappears, the haematuria and albuminuria become rapidly less. If this happens as a definite and sudden change the prognosis is undoubtedly good and the doctor is fairly safe in saying that all will be well. If a few days later the blood pressure has also returned to normal (for instance, the diastolic pressure is 75 mm. or below) then the good prognosis amounts practically to a certainty. It matters not that some small amount of protein (say 0.5 g. per litre or less) remains in the urine for weeks or even for many months in some cases.

Conversely, if the oedema is slow to clear and after several weeks the patient has still a notable albuminuria (say 3 to 10 g. per litre) and his diastolic pressure is 90 to 100, a progressive and unfavourable course is likely. Recurrence of oedema at this stage usually indicates that the course will be rapidly progressive, with death within two years.

At a somewhat later stage, when the patient has apparently recovered but still has some albuminuria, the diastolic pressure and the amount of the proteinuria are still the best guides. If the diastolic pressure is still raised several months after the acute attack the chronic latent course will almost certainly ensue.

Acute nephritis very rarely recurs despite what the textbooks say. When recovery is complete and albuminuria has entirely ceased, no doubts about the future should be allowed to cloud the mind of the patient or his relatives.

In the latent stage the eventual progression to renal failure is inevitable, but the prognostic art is still important in deciding how long it is likely to take. Not that the information will necessarily be communicated either to patient or to relative, but it will influence the doctor's advice in helping the patient to choose a way of life. If the course is likely to be fifteen to twenty years it is mischievous to interfere very much with the patient's liberty to do what he wants, but if the end is likely in a year or two he would obviously be ill advised to embark on a new career or business venture in which illness and incapacity might bring great financial hardship.

Tests of renal function are of some assistance in this assessment. A maximum urinary specific gravity, after fluid restriction, of only 1010 to 1015 is bad, as is also a urea clearance of less than 40 to 50%. The symptom of nocturia usually, but not always, indicates termination within a year

or two. A high diastolic pressure (120 or over) is incompatible with long survival, whereas a relatively low one (say 90 to 100) is favourable. A significant rise in blood urea indicates the approach of the terminal stage.

Prognosis in Type 2 Nephritis

The overall prognosis of Type 2 nephritis, as already stated, is unfavourable. In most cases oedema can be overcome, but gross albuminuria remains and hypertension with renal failure eventually ensue. By far the most accurate guide to prognosis is again the blood pressure. I have never seen recovery in any of my cases in which the blood pressure rose, whether early or late in the disease. If, on the other hand, it stays consistently normal despite oedema and heavy albuminuria complete recovery is always possible, and occurs in perhaps 20 to 30% of these normotensive cases. Some recover completely after several years of varying and recurrent oedema. A raised blood urea augurs a bad prognosis with as much certainty as a raised blood pressure.

Treatment of Type 1 Nephritis

Although our ability to treat nephritis is limited and our knowledge is incomplete, it is impossible even to make a logical attack on the symptoms without having some concept of the disordered physiology and biochemical changes of nephritis.

In acute nephritis the oedema seems to be due to salt and water retention. Whether this is in turn due to altered renal haemodynamics is not quite clear. The water retention gives rise to haemodilution, and the total body water is increased. These are the conditions which are probably responsible for causing convulsions, heart failure, and pulmonary oedema, all of which are complications to be avoided. When anuria or severe oliguria occurs it is not because the body lacks fluid, but because the kidney is not responding to the normal stimuli regulating fluid excretion. To give copious fluids in acute nephritis is therefore like giving sea-water to a drowning man. He has already quite enough.

All cases should be kept in bed throughout the acute stage. Fluid intake should be limited to 500–600 ml. until oedema is subsiding and diuresis established—that is, usually for a week or two. Salt should not be given. Calories should be provided in the form of carbohydrate and fat. Bread, butter, jam, biscuits, fruit, and sugar make up the diet at this stage. Milk is unsuitable, since it contains too much fluid, salt, and protein. When a favourable turn in the symptoms has occurred a light diet containing not much salt or protein is normally prescribed for a week or more before returning to normal food.

If an active infection such as tonsillitis is still present penicillin should, of course, be given, but it does not seem to influence the course of the nephritis, neither do the antihistamine drugs. Cases of acute nephritis, with wet lungs (pulmonary oedema) are very prone to secondary infection, which again requires penicillin. In rare cases in which total anuria occurs it may be treated by the conservative method now used in acute tubular necrosis. The object is to give the patient calories without protein or electrolytes. The method is described in Dr. D. A. K. Black's article (*British Medical Journal*, 1950, 1, 893) under "acute renal failure."

When a patient with acute nephritis has recovered he no longer requires treatment or burdensome restrictions on his

freedom, a fact insufficiently grasped by many doctors. Even in the latent stage it is doubtful whether dietary restriction influences the course of the disease, and it is usually better to let the patient do as he pleases.

The treatment of uraemia—that is, the terminal phase—has already been described in this series (*British Medical Journal*. 1950, 1, 893).

Treatment of Type 2 Nephritis

Except in those rare cases mentioned in Part I of this article which appear to be due to calomel or troloxidone, there is no such thing as curative treatment of Type 2 nephritis. All we can do is to make an attack on the oedema, which, after all, is usually the only symptom that troubles the patient, at any rate until the last stages of his disease.

The sequence of events in the formation of oedema in Type 2 nephritis appears to be as follows: heavy albuminuria causes loss of plasma albumin (hypoalbuminaemia). The normal transudation of fluid from capillaries to interstitial fluid is usually limited by the osmotic pressure of the plasma proteins. In hypoalbuminaemia this osmotic pressure is low and excess of fluid escapes, giving rise to oedema. The loss of fluid from the plasma causes a tendency to haemoconcentration, with a fall in blood volume, and to this the normal response of the kidney is to retain salt and water to build up the blood volume. Thus a vicious circle sets in, the kidney retaining more saline fluid to "replace" that which is lost from the plasma in oedema.

This vicious circle we can attempt to influence at various points. Obviously the right thing would be to stop the albuminuria, but this we have no means of doing. We can, however, give a high-protein diet, with a view to helping the body to replace its protein loss. This can be supplemented by hydrolysed protein foods. In addition we can restrict salt intake and give intravenous protein or other substances of high molecular weight.

For practical purposes a high-protein/low-salt diet is prescribed in the first place. If this is insufficient to keep the patient oedema-free we may be more strict and insist on salt-

that is, as much as is normally contained in two to three litres of plasma. Intravenous medication, therefore, needs to be on a grand scale if used at all.

Fortunately, simple mechanical means of removing oedema are often very successful, and, once free from the major load, the patient may be able to keep his oedema in check by dietary measures alone. In grossly oedematous patients paracentesis of the chest and abdomen may be required. In addition the use of Southey's tubes, now that we have antibiotics to overcome infection, is often most successful. The patient is prepared by being kept propped up in bed for several days, with the legs as low as possible to drain the oedema into the dependent parts. While still in this position, two Southey's tubes are inserted into each leg, and the fluid is drained off into suitable receptacles through fine rubber tubes (Fig. 1). Southey's tubes can be kept in position for three to five days. In this way many litres of fluid may be removed, enormously increasing the comfort and morale of the patient (Fig. 2). It is interesting that fluid disappears not only from the legs but from the whole body, including the face, if the tilted posture of the patient is properly maintained.

Mercurial diuretics have a small but sometimes useful part to play in Type 2 nephritis. They apparently do no harm, but are not very effective, causing usually only a slight diuresis and minimal salt loss, not comparable with their dramatic effect in cardiac oedema.

So much for the accepted forms of treatment. In recent years other means of therapy have been tried. It has been known for a long time that oedematous nephritis is apt to undergo unpredictable spontaneous changes which may sometimes unexpectedly lead to complete remission or apparent recovery. Sometimes this has occurred after infections. In children with nephrosis the effect of infection by malaria has been tried, since it is relatively harmless and controllable, with remission or recovery in some cases. Nitrogen mustard has also been used and probably acts in a similarly non-specific manner.

A sodium depletion more efficient than that achieved by diet can be obtained by the use of ion-exchange resins, but these require careful control and are not yet suitable for use outside hospital. Finally, the effect of A.C.T.H. and cortisone has, of course, been tried. It can be said with some assurance that these agents do not seem to influence the disease fundamentally. During their administration sodium retention generally occurs and oedema increases. On withdrawal, however, a diuresis often takes place which may be followed by a prolonged remission in some instances, even with disappearance of proteinuria.

The treatment of renal oedema may be compared with that of cardiac oedema. By one method or another it is usually possible to keep the patient dry, and by so doing we add greatly to his comfort. It is rarely that we can honestly claim to have influenced the underlying disease.

Next Refresher Course Article.—"Disseminated Sclerosis," by Dr. J. W. Aldren Turner.

FIG. 1.—The use of Southey's tubes in renal oedema.

free bread, and supplement the protein with hydrolysates. If this still does not suffice, intravenous dextran (a high-molecular-weight carbohydrate solution) or intravenous human serum albumin may be tried if available. Intravenous concentrated human plasma is not advised for various reasons. It must be remembered, however, that, while such measures sound very logical, the patients are often losing from 10 to 20 g. of protein a day in the urine—

FIG. 2.—Weight chart showing the effect of draining oedema by Southey's tubes together with the relative lack of effect (in this case) of dextran and a mercurial diuretic.