FAILURE OF MASSIVE DOSES OF VITAMIN B₁₂ IN ACUTE LEUKAEMIA

BY

IAN WELSH, M.B., B.Ch., B.Sc.

University of Witwatersrand; Nuffield Dominion Travelling
Fellow, Nuffield Department of Medicine, Oxford

This trial of vitamin B₁₂ in acute leukaemia was carried out at the suggestion of the Medical Research Council. Clinical workers had reported the apparent occurrence of remissions in patients with malignant disease who had been treated empirically with vitamin B₁₂. It was conceivable that vitamin B₁₂ might promote the differentiation of leukaemic cells into normal leucocytes, on the analogy of the restoration of normal blood formation in pernicious anaemia. Messrs. Glaxo Laboratories made available a supply of vitamin B₁₂ in ampoules containing 1 mg., which is approximately 1,000 times the maintenance dose in pernicious anaemia. The dosage decided on was 1 mg. intramuscularly a day, regardless of the type of the acute leukaemia or the age of the patient.

The co-operation of physicians and clinical pathologists in the Oxford region enabled us to treat and follow up 25 unselected cases of acute leukaemia in less than a year. Diagnosis was made on clinical and blood findings and confirmed by marrow aspiration. Marrow puncture was repeated in the few cases in which the peripheral blood showed much change. The cases were not divided into treated patients and untreated controls, because the usual course of acute leukaemia, including the frequency of partial and complete remissions, is now well known (Bierman *et al.*, 1950; Southam *et al.*, 1951), and any important modification as a result of treatment should soon become apparent. The prescription of blood transfusions and antibiotics was left to the discretion of the physician in charge of the patient, but other remedies such as x rays, radiomimetic drugs, A.C.T.H., and cortisone were not used.

The results are shown in the Table. The patients treated consisted of 11 females and 14 males, their ages varying between 3 and 69 years; 10 were 14 years of age or less, and the average age was 25 years. The cell types were as follows: 10 acute myeloblastic or myeloid, six undifferentiated cell type, five lymphoblastic, and four monocytic. The average time from onset of symptoms to diagnosis was

approximately seven weeks. The survival after diagnosis ranged from 156 days to 2 days, with an average of 52 days. The total duration of the disease thus averaged about 14½ weeks, and there were no instances of noticeably long survival.

All the patients are now dead. None showed a complete remission; the nearest was Case 2, who had a good clinical remission lasting three months, during which time his blast cells dropped from 20% to 5%, but the haemoglobin did not rise above 60% and the platelet count remained at 50,000 per c.mm.

No patient became afebrile on treatment, nor was there any substantial retrogression in the size of the spleen or glands. Only objective signs of clinical improvement, such as gain in weight or rise in haemoglobin, have been recorded. Six cases (Nos. 1, 2, 3, 7, 8, and 9) had a partial remission, and an additional 10 felt better but did not show any objective change in their condition. No toxic effect attributable to the B₁₂ was observed.

Summary

Twenty-five patients with acute leukaemia have been treated with massive doses of vitamin B₁₂. None lived for more than five months after beginning treatment. No complete remissions occurred, but six patients had partial remissions and a further 10 reported feeling better.

There is no evidence from this series to suggest that the prognosis or clinical course of acute leukaemia is altered by adding vitamin B₁₂ to other treatment.

Thanks are due to Professor L. J. Witts, in whose department much of the work was performed and whose help is gratefully acknowledged; to Messrs. Glaxo Laboratories for a generous supply of vitamin B₁₂; and to the physicians and clinical pathologists of the Oxford region for their co-operation.

REFERENCES

- Bierman, H. R., Cohen, P., McClelland, J. N., and Shimkin, M. B. (1950). *J. Pediatr.*, 37, 455.
Southam, C. M., Craver, L. F., Dargeon, H. W., and Burchenal, J. H. (1951). *Cancer*, 4, 39.

USE OF A.C.T.H. IN AGRANULOCYTOSIS

BY

H. FULD, M.D., M.R.C.P.Ed.

Physician, Sefton General Hospital, Liverpool

The papers by Discombe (1952) and by Dudley Hart *et al.* (1952) prompt me to publish the following notes. In the former article the relatively frequent occurrence of agranulocytosis on the Continent is compared with the rarity of the condition in this country, and is explained by the popularity which amidopyrine has gained and still maintains outside Great Britain. My own experience (Fuld, 1931) of having observed three fatal cases of agranulocytosis within six months in 1930 in a medium-sized German hospital where "pyramidon" (amidopyrine) was the standard drug, only second to aspirin in its frequency of administration, supports Dr. Discombe's conclusions. This is, however, only in retrospect, as I was quite unaware of the relationship of amidopyrine to agranulocytosis at that time.

In the past 19 years I have, of course, come across a number of patients with more or less severe leucopenic reactions following sulphonamides and thiouracil preparations, but no true case of agranulocytosis was met with until May, 1952.

Case Report

A woman aged 65 was referred by her doctor on May 30, 1952. She had had remarkably good health until a year previously, when she complained of lack of energy and

Results of Treatment

No.	Sex	Age in Years	Diagnosis	Duration of Symptoms Before Diagnosis	Survival After Diagnosis in Days	Total B ₁₂ in mg.	Partial Remission
1	F	4	Acute undifferentiated	8 weeks	156	68	+
2	M	40	Acute myeloblastic	2 "	130	130	+
3	F	4	" undifferentiated	10 "	120	48	+
4	F	17	Acute monocytic	12 "	103	103	0
5	M	69	" "	8 "	96	96	0
6	M	63	" myeloblastic	14 "	90	16	0
7	F	29	" monocytic	3 "	90	65	+
8	M	44	" lymphoblastic	2 "	70	70	+
9	M	3	" "	8 "	65	60	+
10	F	3	" stem cell	12 "	62	56	0
11	M	30	" monocytic	12 "	58	48	0
12	M	14	" myeloblastic	4 "	45	45	0
13	M	22	" undifferentiated	2 "	44	44	0
14	M	34	Acute lymphoblastic	4 "	40	40	0
15	M	56	" "	3 "	35	35	0
16	M	23	" myeloid	? mths	21	21	0
17	F	19	" "	7 "	18	18	0
18	M	4	" "	3 weeks	11	11	0
19	F	43	" myeloblastic	9 "	8	8	0
20	F	54	" myeloid	5 "	8	8	0
21	F	47	" lymphoblastic	16 "	6	6	0
22	M	30	" myeloblastic	8 "	6	6	0
23	F	41	" erythraemic myelosis	3 "	5	5	0
24	M	14	Undifferentiated	4 "	5	5	0
25	F	34	" "	12 "	2	2	0

general weakness which had persisted ever since. Her appetite had deteriorated during the last three months. In March she had a short course of sulphonamides for a septic finger, and again ten days later for a boil on her back which never healed properly. She did not notice any change in her general condition before and after this treatment. During the last few weeks she rested a good deal, but was not entirely confined to bed.

On examination she appeared a rather pale thin woman with evidence of recent loss of weight. Apart from a radiologically confirmed slight enlargement of her heart and a systolic apical murmur there were no significant changes in her cardiovascular system. Her palate was granular. There were no ulcers in her mouth or pharynx. Her spleen reached 1 in. (2.5 cm.) below the costal margin, and was hard. Her blood count on May 30 was: red cells, 2,550,000; haemoglobin, 48%; colour index, 0.94; white cells, 600 (polymorphs 3%, lymphocytes 78%, monocytes 19%). X-ray films of the skeletal system showed Paget's disease of the right femur.

A diagnosis of agranulocytosis was made, and she was put on treatment which included "cytamen," iron, folic acid, ascorbic acid, the vitamin-B complex reinforced by pyridoxine, as well as blood transfusions (2 pints (1.1 litres) on June 4 and on June 24). She also received large doses of penicillin, and, in the later course of her illness, streptomycin. On June 3 her peripheral count had not materially altered: red cells, 2,580,000; haemoglobin, 49%; colour index, 0.96; white cells, 650 (all lymphocytes). Her bone-marrow count consisted of: blast cells, 0.5%; premyelocytes, 1%; myelocytes, 29%; metamyelocytes, 4%; neutrophil polymorphs, 0.5%; eosinophils, 0.5%; normoblasts, early 3.5%, intermediate 11.5%, late 4.5%; monocytes, 2%; lymphocytes, 43%. The myeloid/erythroid ratio was 1.8:1.

On June 11 her white cells numbered 950 without any lymphocytes being seen. On June 19 her red-cell count was 3,220,000; haemoglobin, 59%; colour index, 0.9. There were 800 white cells, all lymphocytes. On June 23 her temperature rose rather high and remained so for three days in spite of antibiotics. An irritating laryngitis and tracheitis had developed and there was a small clean ulcer on her hard palate. On June 24 her white cells numbered 600. No granulocytes were seen.

A.C.T.H. was started on June 25, and for six days she received 20 mg. six-hourly. On July 1 the dose was cut to 20 mg. twice daily, and the treatment was stopped after a total of 600 mg. had been given in nine days. After 140 mg. of A.C.T.H. had been injected the bone-marrow content was: myelocytes, 16%; metamyelocytes, 13%; neutrophils, 2%; lymphoblasts, 2%; lymphocytes (large and small), 60%; normoblasts, early 1%, intermediate 4%, late 2%. At that time she had 500 white cells per c.mm., with 6% neutrophils and 94% lymphocytes in her peripheral blood. On the sixth day of treatment with A.C.T.H. her temperature again rose and she developed laryngitis and diarrhoea, which did not respond to antibiotics. On the eighth day her peripheral blood count was: red cells, 2,600,000; haemoglobin, 54%; colour index, 0.9; white cells, 450. No granulocytes were seen.

A.C.T.H. was stopped on July 3, as neither the patient's general condition nor her bone-marrow function or peripheral blood showed evidence of recovery. She was again put on the remedies which were given before the trial with A.C.T.H., but her condition steadily deteriorated and the last blood count taken on July 11 (20 hours before her death) revealed only 200 white cells with complete absence of granulocytes. She still maintained her red-cell and haemoglobin level. Her temperature reached 104° F. (40° C.), and the diarrhoea persisted. Pericardial friction rubs were heard and she developed oedema of the legs. On the evening of that day she sank into coma and died in the early morning hours of July 12.

Conclusions

This patient had been ill for many months before she came under observation, and I believe that her blood dis-

order existed before sulphonamides were prescribed. Neither antibiotics nor A.C.T.H. had any appreciable effect on the outcome.

I believe that this case belongs to the group of idiopathic agranulocytosis cases which appear to remain unaffected by A.C.T.H. Caldwell (1950) and McMillin (1951) both deal with drug-induced cases of agranulocytosis successfully treated by A.C.T.H. and cortisone respectively. Dudley Hart *et al.* wrote with reference to A.C.T.H. or cortisone that "they may act only in some cases of acute drug-induced agranulocytosis." My own experience entirely supports their opinion.

Idiopathic agranulocytosis, first described 50 years ago, still remains a problem so far as aetiology is concerned and a challenge in respect to treatment.

Summary

A case of agranulocytosis of unknown origin is described. The patient failed to respond to treatment, which included A.C.T.H. over a period of nine days.

The prognosis for agranulocytosis of unknown aetiology remains grave in spite of the introduction of cortisone and A.C.T.H.

REFERENCES

- Caldwell, A. L., *et al.* (1950). *Canad. med. Ass. J.*, **62**, 506.
Discombe, G. (1952). *British Medical Journal*, **1**, 1270.
Fuld, H. (1931). *Dtsch. Arch. klin. Med.*, **170**, 100.
Hart, F. Dudley, *et al.* (1952). *British Medical Journal*, **1**, 1273.
McMillin, J. S. (1951). *Amer. J. med. Sci.*, **222**, 392.

Medical Memoranda

Fractured Shaft of Femur Complicated by Multiple Systemic Emboli

CASE REPORT

A policeman aged 41 was admitted to hospital on March 4, 1951, having fallen sideways off his bicycle on to his left hip, and sustaining a comminuted fracture of the upper third of the left femur. At the time there was no evidence of a nerve or vascular lesion. X-ray films showed a comminuted pertrochanteric fracture of the femur, with a long posterior fragment. A Steinmann pin was inserted into his tibial tubercle and a balanced skeletal traction set up. It was thought, because of the comminution, that closer apposition of the fragment could be obtained by applying a clamp to the Thomas splint. This did, in fact, improve the alignment.

At 7.40 a.m. on April 4, exactly a month after admission, he had a sudden right hemiplegia. This cleared up rapidly, and when seen again at 11 a.m. he had only a residual right facial weakness. His blood pressure was 100/80. Dr. Kembell Price thought the condition was due to an embolism. At 5 p.m. the patient complained that his right leg was cold and painful for an hour. On examination the right leg felt cold and there was complete paralysis of all his muscles below the knee. His right popliteal fossa and calf were extremely tender. The posterior tibial and dorsalis pedis arteris could not be felt. A diagnosis of popliteal embolus was made.

Operation.—At 8 p.m. skeletal traction was converted into fixed traction and a spinal anaesthetic was given. Exploration of the popliteal vessels was carried out through a long midline incision. It was noted that after the induction of anaesthesia vasodilatation did not occur in the right leg, although it did in the left. The popliteal vessels were exposed. The popliteal artery felt hard and was white. It was opened and a 3-in. (7.6-cm.) clot removed. This, however, was not followed by a sudden spurt of blood from the proximal end, and it was not until the last piece of the clot was removed from the portion of the artery which lay in Hunter's canal that the artery was found to be clear. The