professors came all too soon. He left Cambridge with deep regret, full of gratitude for the kindness and friendship he had received from Corpus Christi College and from that College of his young enthusiasm, St. John's College, whence came the invitation which he received and accepted with great pride—to give the Linacre lecture in 1941. He returned to Cambridge for the last years of his life, and he was occupied till the end in writing *Some Chapters in Cambridge Medical History*, published in 1946 by the University Press.

Conclusion

Sir Walter Langdon-Brown died in 1946 at the age of 76 honoured by all and loved by his friends, and holding the devoted affection of the nearest and dearest to him. He used to speak with pleasure and satisfaction of the survival of the institutions, of his beloved university in particular, throughout future generations. I would like to speak of the survival of his life's work, the central thesis of which is best defined in his own words. "As we are animals we conform to biological laws. Our future evolution physically is improbable and mentally unlikely. Therefore only a psychological evolution remains. This necessitates co-operation, not merely instinctive as with bees and ants, but intelligent and voluntary. Now we, like all animals, carry with us vestigial traces of our past ancestry, not least in our mental processes. To develop psychologically we must understand ourselves, and it should help us to do so if we can find ways to investigate those hidden depths in our minds from which we draw our impulses."

Psychology has for some years been chiefly concerned in looking back, and it has been a preoccupation of psychologists to search out the origin of thoughts and feelings. This has been a contribution to knowledge as was Charles Darwin's work on the origin of the species. Langdon-Brown asks us to look forward and to find ways—new ways—to investigate the hidden depths of the human mind. Godwin Baynes, speaking of a moral act or response being invariably individual in origin, said, "A moral feeling can be imparted to a community, but its source is in the solitary individual soul." The same holds good for original ideas; they too have their origin in the individual mind. The work and teaching of Sir Walter Langdon-Brown was spread on so large a canvas that it must be viewed from a distance to be seen as a whole. His contribution to medicine was no discovery of factual knowledge, nor was it the presentation of some ideas; it was a wide conception which has given a fresh inspiration and a new direction to the understanding of the human mind.

According to the Swedish International Press Bureau the Swedish Medical Association has agreed on a *Codex Ethicus Medicorum* embodying 15 rules of ethics for medical practitioners in Sweden. Simultaneously a new set of rules for medical specialists has been agreed upon. These two briefs of rules aim at maintaining the high standard of Swedish physicians. They contain instructions on the relationship between the doctor and his patient, and clearly define the duty of the physician towards the preservation of human life, professional secrecy and ethical points about monetary reward, publicity, etc. It is stressed specially that a physician should refrain from advertising himself or his work in an importunate or flagrant way and not draw inappropriate attention to his person when giving information to the Press.

GENERALIZED OSTEOARTHRITIS AND HEBERDEN'S NODES

BY

J. H. KELLGREN, F.R.C.P., F.R.C.S.

AND

R. MOORE, M.R.C.P.*

(From the University Rheumatism Research Centre at the
Royal Infirmary, Manchester)

Osteoarthritis is usually considered to result from degenerative changes in the articular cartilage and other joint structures which occur to some extent in most individuals with advancing age, but which may occur prematurely in joints subjected to abnormal wear and tear from some special occupational stress or as a result of faulty joint mechanics.

Premature degeneration of a severe nature and more rapid in evolution occurs in joints in which the articular cartilage has been damaged by trauma, infection, or avascular necrosis of the underlying bone. Rapid degeneration may also occur in joints deprived of the protective reflexes afforded by normal pain and proprioceptive sensibility.

Although the degenerative arthritis of old age may affect many joints, the premature degeneration resulting from previous damage by trauma, etc., tends to affect only one or at the most a few joints, because the degenerative process is secondary to these causative factors, which as a rule are operative in only one or a few joints and, since the degenerative process is slowly progressive, the resultant disability also tends to be slowly progressive, though acute painful incidents following trivial injury or unusual stress may be superadded.

The clinical and radiological features of degenerative arthritis are well known, and patients with this condition are seen with great frequency, particularly in orthopaedic clinics; but in clinics dealing largely with rheumatic disease one often sees polyarthritic patients in whom the joint changes in the later stages resemble those of degenerative arthritis, though they differ somewhat in that they often have a rather acute spontaneous onset. Furthermore, these patients tend to present a definite pattern of joint involvement characterized by affection of the distal interphalangeal joints of the fingers, and the first carpo-metacarpal joints in the hand, the great toes, and first tarso-metatarsal joints in the feet, the interfacetal joints of the spine, the knees, hips, and other limb joints. These patients also present certain other characteristic clinical and radiological features which differ from those found in classical degenerative joint disease on the one hand and polyarthritis of the rheumatoid type on the other, and we have come to classify these cases under the heading of primary generalized osteoarthritis, which we consider to be a distinct clinical entity.

More than 150 years ago William Heberden recognized the fact that the nodes so often seen over the distal interphalangeal joints of the fingers represented a condition distinct from either gout or rheumatism, and in 1805 Haygarth described 34 cases of multiple arthritis associated with Heberden's nodes which he classified separately under the heading of "nodosity of joints." A

*Nuffield Fellow in chronic rheumatism.

polyarticular osteoarthritis was also mentioned by Adams (1857) and Charcot (1881). In more recent years Cecil and Archer (1926), while classifying 182 cases of degenerative arthritis attending their rheumatism clinic, noted that only 20 were of the classical monarticular type and only 17 of the senile type, while 145 were polyarticular and usually associated with Heberden's nodes. Because this latter group was largely composed of middle-aged women they considered this form of arthritis to be "menopausal." Francon (1950) also describes a generalized form of arthritis associated with Heberden's nodes.

On the other hand, Stecher (1948) considered these nodes to be a distinct entity, although he did note a tendency to involvement of other joints (Stecher, 1946b); but in this connexion it must be remembered that Stecher was engaged in a genetic study of the incidence of asymptomatic nodes in the general population, and he was less concerned with the more severe forms of this condition which send the sufferer to the rheumatism clinic. Stecher, however, made a most valuable contribution when he separated the post-traumatic type of node from the idiopathic inherited variety.

Interest in the question of whether Heberden's nodes are associated with generalized arthritis has been stimulated in this department because a routine examination of all the joints of the body is carried out in every patient attending the clinic, even though they may complain only of a single joint, and this routine has allowed us to recognize more clearly the patterns of joint involvement characteristic of the various forms of articular disease, and in particular that associated with Heberden's nodes.

The following description of what we consider to be primary generalized osteoarthritis is based on the experience of 391 cases of osteoarthritis taken from a total of 1,813 cases of arthritis of all types seen in this department during the three years preceding 1951. In addition we have made a detailed analysis of the 196 cases of osteoarthritis seen during the year 1950. Of these, 120 had either Heberden's nodes or arthritis of the first carpo-metacarpal joints or both; 20 were of the classical monarticular degenerative type, and 56 were classed as miscellaneous because, although their arthritis was polyarticular, the characteristic involvement of the hands was not found.

Sex and Age

In the first instance we will describe the features of those cases with characteristic involvement of the hand joints. The condition is found predominantly in middle-aged women; thus of the 120 analysed cases 110 were in women, whose ages at the onset of the disease varied from 35 to 73, with a mean of 52 years. There was, however, no accurate correlation with the menopause, and in 22 cases the arthritis began while the menses were still normal. The age of onset in the 10 men ranged from 20 to 63, with a mean of 50.7 years.

Pattern of Joint Involvement

This is one of the most characteristic features of primary generalized osteoarthritis. Thus of the 120 analysed cases 103 had Heberden's nodes, 79 had arthritis of the first carpo-metacarpal joints, and 62 had both. In a few cases the arthritis also affected the joint between the scaphoid and the multangulum majus, but the rest of the carpus and the metacarpo-phalangeal joints which are so characteristically affected in rheuma-

toid arthritis were usually spared, only very slight changes in these joints being found in 19 cases. On the other hand, the proximal interphalangeal joints were often severely involved, these joints being affected to some extent in 49 cases. The joints of the feet were affected in 50 cases, and in 40 of these the metatarso-phalangeal joint of the great toes was the one maximally affected.

In 22 cases there was also some involvement of the lateral metatarso-phalangeal joints, but the changes in these were always less than in the great toe, which is the reverse of what is found in rheumatoid arthritis; there was also some tarsal arthritis in 20 cases, and in several the joint between the first cuneiform and the first metatarsal was markedly affected. The knees were affected in 64 cases, nearly always bilaterally, and the hips in 36, while the spine was affected in 57, being the most frequent part to be involved after the hands and knees. The remaining joints were but rarely affected, and usually only to a lesser degree in those florid cases in which nearly all the joints of the body were involved.

Nature of Joint Lesion

Each affected joint tends to pass through two stages: an initial relatively acute phase during which the joint may be warm, red, and exquisitely tender; and, some months later, a chronic phase characterized by bony outgrowths around the joint margins. In the acute phase the joint is usually slightly swollen and may contain a small effusion, this excess fluid being particularly noticeable in the knees. The massive soft-tissue swelling and synovial thickening characteristic of many cases of rheumatoid arthritis is, however, not seen. During this initial acute phase spontaneous pain may be very troublesome. It often comes in severe spasms, particularly at night-time, and waking the patient in the early hours. When the finger-joints are severely affected and the skin over them becomes reddened and tender the pain acquires a characteristic burning, tingling, and bursting quality, and the patients place their hands outside the bed to obtain relief. This type of pain is particularly evident in those cases in which small cystic swellings form over the finger-joints, finally to burst through the skin, discharging a little glairy fluid (Fig. 1). More usually the pain is of the deep aching type, and is not well localized to the joints, often spreading widely up and down the limbs. This deep pain is also markedly aggravated by cooling the part, and these patients avoid cold water and dread exposing their hands in cold weather, although vasospasm is not a feature of this condition.

During the acute phase a few minutes of arterial occlusion produced by a cuff on the arm will cause a rapid and severe increase of pain: in this respect the pain resembles that of acute suppurative lesions such as whitlows (Addis, Jepson, and Kellgren,

FIG. 1.—Early node with discharging cyst (Case 1).

1950), and indeed two of our more acute cases were initially thought to be suffering from pulp-space infections of the finger. When a large joint such as the knee is affected in this acute manner the pain may be very severe, and the condition is not unlike that of a low-grade suppurative arthritis, though the constitutional disturbance usually associated with infection is absent.

After a period of several months the acute phase subsides and the clinical signs of inflammation disappear, giving place to the chronic phase, which is characterized by massive bony outgrowths around the joint margins. At this stage there is little pain, and, in spite of the unsightly deformity and gross radiological changes, the joints often retain a fair range of movement and function

progresses first one joint becomes involved and then another, each joint passing through a more or less painful phase and going on to bony enlargement, which is relatively painless.

By no means all the cases present with an onset which is acute enough to resemble a pyogenic process; indeed, only 21 of the 120 cases had incidents of this type in one joint or another, the more usual story being one of a sub-acute or even gradual onset. Nevertheless, a more or less

FIG. 2.—Multiple nodes and enlargement and adduction deformity of first carpo-metacarpal joint (Case 2).

is largely restored; thus permanent crippling very rarely ensues. The appearance of the hand in a more advanced case is shown in Fig. 2. Note the bony ridges over the terminal joints of the finger, the enlargement and deformity of the proximal interphalangeal joints, and also the bony enlargement, partial subluxation, and adduction deformity of the first carpo-metacarpal joint, giving the hand a curious squared appearance.

Fig. 3 shows the hand of a very advanced case. Although in this case nearly all the joints show bony enlargement the characteristic appearance is still retained. The condition in the hand is quite characteristic and easily recognized, but in the less accessible joints bony enlargement is less readily appreciated, and in the spinal articular processes it may be easily overlooked. Thus in the spine there is often a history of severe back pain and disability lasting several months followed by partial recovery which leaves the patient with only slight limitation of spinal movement, particularly in side flexion and rotation, and it is only by radiography that the marked bony outgrowth on the articular facets and spinous processes can be demonstrated. As the disease

FIG. 3.—Advanced case with bony enlargement of all joints.

painful initial phase in each affected joint is the rule in the more severe cases. The characteristic course of the disease is well illustrated by the following case histories.

Case 1

A housewife aged 56 had noticed increasing deformity in the distal and proximal interphalangeal joints of the hands accompanied by much pain for 15 years. During this time acute joint incidents had been noticed, in which first one and then another of the distal interphalangeal joints of the hands would become acutely swollen, painful, and red, and a cystic swelling would appear at the lateral aspect of the joint. This became so painful that she evacuated it with a needle, after which the pain would subside for about a fortnight, to recur as the swelling filled up again. Gradually the swelling became painless, stopped discharging, and left a hard bony knob at the base of the distal phalanx. This series of events then occurred in another distal interphalangeal joint over a period of six months before the discharge finally ceased. For four years she had been troubled by pain and stiffness in the left ankle and had had a goitre from the age of 14 years. The menopause occurred at 46. One sister had Heberden's nodes.

On examination she was seen to be a healthy woman weighing 8 st. 12 lb. (56.2 kg.). A small diffuse goitre was present. There was no clinical evidence of visceral disease. The hands showed large tender Heberden's nodes, and enlargement, tenderness, and limited movement of most of the proximal interphalangeal joints and both first carpo-metacarpal joints. There was also some swelling of the left ankle, and its movements were slightly limited. The distal interphalangeal joint of the right middle finger was very swollen and an acutely tender cystic swelling was present on the lateral edge of the joint, which had discharged a yellowish jelly-like fluid. The erythrocyte sedimentation rate was 35 mm. in one hour, the haemoglobin 84%, and the white blood cells 5,700 per c.mm.

Case 2

A housewife aged 65 had first developed pain and swelling of the distal interphalangeal joints and the first carpo-metacarpal joints 12 years ago. Acute episodes had occurred in the distal interphalangeal joints, lasting months and leaving Heberden's nodes in their wake. At this time the shoulders also became stiff. Eight years ago the knees had become swollen, painful, and stiff, and after a further four years the lower part of the back had become painful. She had been gaining weight for about four years. She had pernicious anaemia at 57 years and had passed a tapeworm at 63. The menopause occurred at 54. There was no family history of rheumatism.

On examination she was seen to be a robust woman weighing 10 st. 12 lb. (69 kg.). There was no visceral disease. Painful Heberden's nodes were present on all the fingers; both first carpo-metacarpal joints were painful, swollen, and limited, especially the left one; and both knees were slightly swollen and painful, and contained effusions. Movement in the right was from 5 to 110 degrees flexion, while that in the left was normal. She attended monthly for observation, and in the third month suddenly developed acute pain in the right knee, which became very much more swollen owing to increase in the effusion; the overlying skin was reddened, and movement was reduced to 5 to 70 degrees flexion. The joint had, in fact, the appearance of an acute arthritis, and she took to her bed for a fortnight, by the end of which time the pain began to subside. A plaster leg cylinder was then applied and she was encouraged to use the leg. After six weeks the plaster was removed, as the knee had returned to its previous condition; its movement was again what it had been before the acute incident. When she was first seen the erythrocyte sedimentation rate was 15 mm. in one hour; it rose to 26 mm. during the acute joint incident. The haemoglobin was 86% and the white cell count 5,300 per c.mm.

Case 3

A housewife aged 60 had noticed pain and swelling of the first carpo-metacarpal joints of both hands with weakness of the grip six years ago. Several months later the distal interphalangeal joints of the middle fingers of each hand became acutely swollen and painful, and a cystic swelling appeared on the dorsum of the middle left distal interphalangeal joint which discharged a yellowish fluid intermittently for a considerable time. The acute swellings subsequently involved the distal interphalangeal joints of the other fingers of both hands, eventually becoming painless over a period of three years, leaving bony swellings typical of Heberden's nodes. Four years ago the right hip gradually became troublesome: she noticed stabbing pains in the thigh on walking, the hip gradually developed a flexion deformity, and she suffered from severe pain in the groins, particularly when lying in bed. At this time walking became difficult.

Eighteen months ago she sought surgical advice, and, after a right subtrochanteric osteotomy had been performed, was nursed in a double hip spica for the next five months. On removal of the plaster it was noticed that the left hip and lumbar spine had become stiff and painful. Since removal

of the plaster the knees have been painful and have gradually lost their range of movement, and she is now only able to walk a few yards with the aid of sticks. Her general health has remained good throughout, and there has been no weight loss. The menopause occurred at 53 years. There was no family history of rheumatism.

On examination she was seen to be a pale slim woman in good general health but distressed by pain on movement in both hips. Her blood pressure was 210/110, but there were no clinical signs of visceral disease. Pronounced Heberden's nodes were present on the fingers and there were marked bony enlargement and deformity of the first carpo-metacarpal joints and some bony enlargement of the proximal interphalangeal joints, but the wrist movements were full and painless. Dupuytren's contracture of the fourth and fifth fingers was present in the right hand. The knees were slightly swollen and tender, and their movement was moderately limited. There was bony enlargement of the first metatarsophalangeal joints of the feet, with some limitation of movement. Examination of the spine revealed a flattening of the lumbar curve and almost complete absence of movement in the dorsal and lumbar regions. The right hip was completely fixed in 10-degrees flexion, while the left hip was capable of movement through 15 degrees of flexion. The erythrocyte sedimentation rate was 13 mm. in one hour, the haemoglobin 98%, and the white cell count 4,400 per c.mm.

Other Clinical Features

In spite of the acute and apparently inflammatory nature of the initial arthritic process we have not observed any constitutional disturbance in these patients. Thus weight loss, which is such a feature of rheumatoid arthritis, was not observed, and the blood picture was not abnormal. On the other hand, the sedimentation rate, estimated by the Westergren method, was often somewhat raised; thus out of the 112 cases for which figures are available 66 had an E.S.R. of under 20 mm. in the hour. In 38 cases the E.S.R. was between 20 and 40 mm., but in only 8 cases was it over 40 mm. The mean value for the whole group was 19 mm. This contrasts with the findings in our group of 20 cases of monarticular degenerative arthritis, in which the E.S.R. was less than 20 mm. in every case, the mean value for the group being 11 mm. The differential agglutination reaction, which is so often positive in rheumatoid arthritis (Rose *et al.*, 1948; Ball, 1950), was also uniformly negative in these cases.

Soft-tissue lesions, such as thickening of the bursae, and widespread involvement of muscles and fascia were not observed. The florid multiple tendon lesions so characteristic of rheumatoid disease (Kellgren and Ball, 1950) were also not observed in these cases, although in 13 of them a few small firm nodules could be felt in the flexor tendons of the hands. There was also no tendency to vasospasm or excess sweating; neither were there any cutaneous or eye lesions or characteristic visceral disturbances. Thus the disease process seems to be largely confined to the joints, and particularly the diarthrodial joints.

It has been suggested by Fletcher (1947) that both obesity and hypertension occur with unusual frequency in patients with osteoarthritis. On the other hand, Stecher (1946a) did not find any such correlation with Heberden's nodes. Among our 120 cases there were no patients who were grossly obese, and in only four was the systolic blood pressure over 200 mm., and in only two was the diastolic pressure over 120 mm. Thus neither marked obesity nor serious hypertension appears to be a feature of generalized osteoarthritis.

Radiological Changes

The x-ray appearances of the stage of bony enlargement conform to those of osteoarthritis. There is narrowing of the joint space, marked sclerosis of the subchondral bone, and florid bony outgrowths at the articular margins. The radiographs reproduced in Figs. 4-9 illustrate the changes seen in the various joints commonly affected. The combination of Heberden's nodes and arthritis of the first carpo-metacarpal joint is a most characteristic finding, and the enlarged views of these joints show the marked tendency to bony enlargement that is so characteristic of the later stages. The changes in the spine (Fig. 7) are peculiar in that the intervertebral disks remain relatively unaffected, while the articular facets and the neural arches, including the spinous processes, become enormously enlarged and sclerosed, so that "kissing spines" are often seen. Thus the spinal changes conform to the osteoarthritis of Collins (1949), who distinguishes between these changes and the osteophytosis of the vertebral bodies which so frequently accompanies degeneration of the intervertebral disk.

In the hips (Fig. 8) the changes are less distinctive. There is a general loss of joint space and the femoral head tends to become elongated; in advanced cases it becomes enclosed in a deep sclerosed acetabulum, resulting from marked overgrowth of the acetabular margin and not from protrusio acetabuli. In several

cases crumbling of the upper quadrant of the femoral head was noted in one hip, but in these instances there was a past history of significant trauma to the hip. In the knee (Fig. 9) the radiological picture is again fairly distinctive. There is marked narrowing of the joint space combined with rounded "molten wax" bony outgrowths at the articular margins and a marked absence of the sharp-pointed osteophytes so commonly seen in the

FIG. 4.—Whole hand, showing marked changes in terminal interphalangeal and first carpo-metacarpal joints: there are also slight changes in some of the proximal interphalangeal joints.

FIG. 5.—Enlarged views of advanced finger node. The most marked changes are seen in the lateral view.

FIG. 6.—Full-size view of first carpo-metacarpal joint showing much new bone formation.

FIG. 7.—Lumbar spine, showing marked enlargement of facets and spinous processes.

degenerative arthritis which follows injury.

Although the overall radiological picture is certainly that of osteoarthritis, in that narrowing of joint space, subchondral sclerosis, and marginal bony overgrowth are present, there are certain differences between the radiological appearances of primary generalized osteoarthritis and those of classical degenerative joint disease.

FIG. 8.—Hips, showing early and advanced changes.

FIG. 9.—Knee, showing marked "molten wax" osteophytes.

The Miscellaneous Group

On analysing the miscellaneous group of 56 patients who had polyarticular osteoarthritis but without Heberden's nodes or arthritis of the first carpo-metacarpal joints it became apparent that many of them presented features which we have come to regard as characteristic of primary generalized osteoarthritis, and it seems probable that further experience may justify the classification of many of these cases under the heading of generalized osteoarthritis; we are not yet, however, prepared to make this diagnosis in the absence of Heberden's nodes or arthritis of the first carpo-metacarpal joints. Furthermore, we fully recognize that there are many patients in whom degenerative arthritis of the terminal finger-joints may follow injury or unusual occupational strain (Stecher's traumatic nodes), and that arthritis of similar pathogenesis may affect the first carpo-metacarpal joint; but these cases are easily recognized, and have been excluded from the present investigation.

Treatment

In view of the large number of middle-aged women in our series and the fact that other workers have considered this form of arthritis to be menopausal, we carried out a blind clinical trial of oestrogen therapy. Twenty severely affected middle-aged women were selected, and were divided into two groups of 10, which were as evenly balanced as possible in regard to age, severity, and duration of symptoms, E.S.R., and number of joints affected. Mr. Lloyd, the hospital pharmacist, kindly provided us with two lots of indistinguishable tablets, A and B, one inert and the other containing 0.3 mg. of dienestrol. These were administered in a dosage of one tablet three times a day for six months, 10 patients receiving tablets A and 10 tablets B. The patients were seen monthly and were fully assessed before treatment and at the end of six months. Neither group reported any marked subjective improvement, and examination revealed a slight deterioration in both groups. Thus the grip measured by the method of Janus (1950) fell slightly in both groups. The E.S.R. rose slightly in both groups, and in both there were a small number of previously affected joints which became painless and a larger number of previously normal joints which became painful during the six-months course. Most of the affected joints, however, remained unchanged. The mean values in the two groups before and after treatment are shown in the table.

	Before Treatment			After 6 Months				
	Grip	E.S.R.	No. of Joints Affected	Grip	E.S.R.	No. of Joints		
						Recover-	Un-	Newly
Control	215	22	22	180	26	1	21	4
Dienestrol	210	23	19	200	26	1	18	3

During the following six months the tablets were reversed, so that the patient previously receiving A now received B, and vice versa. The second six months has not quite been completed in all patients, but it is clear

that no change has taken place, except that while receiving B one or two patients noted uterine bleeding, and this, in fact, proved to be the active tablet. Although there were only 10 cases in each group the remarkable similarity in behaviour between the two groups suggests that dienoestrol in the dosage given fails to influence the arthritic process in any way.

In fact, we are not aware of any treatment which will influence the arthritic process, but many of these patients fear that they are suffering from rheumatoid arthritis and are grateful for an explanation of the nature of their complaint and its probable future course, and some relief may be obtained by the proper management of their arthritis. Thus acute painful episodes in the larger joints respond well to short periods of immobilization in plaster, and pain may also be temporarily relieved by local heat, which is most conveniently applied from an electric heating-pad that the patient can use in her own home. Aspirin analgesics may be helpful, but are not as effective in this condition as in rheumatoid arthritis. Surgical reconstruction of the larger joints is probably best avoided in these cases owing to the marked tendency for new joints to become affected. Fortunately most patients manage to continue a useful life, and it is only the exceptional case which becomes severely crippled.

Discussion

It would appear that the clinical syndrome which we have come to call primary generalized osteoarthritis has certain distinctive clinical and radiological features which enable it to be distinguished from secondary degenerative arthritis and also from rheumatoid arthritis and other allied polyarthritic conditions. Although generalized osteoarthritis has been recognized since the time of Heberden and Haygarth it appears to have been largely lost sight of in recent years, perhaps because it has been described under different names, such as nodosity of joints, menopausal arthritis, and so on. With the clearer definition of degenerative arthritis as an old-age change occurring prematurely in joints damaged by trauma, avascular necrosis, infection, or previous rheumatic disease, or in those worn out by occupational stress and faulty joint mechanics, a definite concept of osteoarthritis has developed.

The polyarticular form of osteoarthritis we have just described does not readily fit into this concept, and we suspect that this has led to its neglect and caused many cases to be classified as rheumatoid arthritis, or as atypical or "mixed" forms of arthritis. That rheumatoid arthritis may coexist with generalized osteoarthritis is very likely, as both diseases are common in middle-aged women, and in our series such a double diagnosis was made four times in 110 female cases. This, however, would be the expected number if the incidence were governed by chance, since in a recent field survey of the general population in a Lancashire town the incidence of rheumatoid arthritis was found to be 4% in the adult female population.

From the work of Stecher it would appear that idiopathic Heberden's nodes are exceedingly common in elderly people, and it seems likely that only in a minority of those affected are the symptoms severe enough to make the patient attend a clinic specializing in rheumatic disease, but in those who do attend such clinics multiple joint involvement is a common finding. We feel, however, that the question of the incidence of multiple joint involvement in people exhibiting asympto-

matic Heberden's nodes requires reinvestigation. In our series only 20 out of 120 patients gave a history of similar disease in their known relatives, but a very different incidence of Heberden's nodes might have been revealed if the relatives had been examined, and we have certainly seen families in which all the sisters were severely affected. We are therefore inclined to the view that generalized osteoarthritis as we see it in the clinic represents a severe form of Stecher's idiopathic nodes. If that is so, this syndrome, like the nodes, may result from an inherited defect, and should be regarded as a constitutional disease.

Although in the later stages the joint lesions resemble those of classical degenerative arthritis, we have been unable to find a description of the joint changes during the initial acute phase, and this point clearly requires urgent investigation, as the nature of the morbid process at this stage remains unknown. Our object so far has been to define what seems to be a distinct clinical entity, and thereby to stimulate further investigation into its pathogenesis and the aetiological factors which influence its development.

Summary

From a study of 391 cases of osteoarthritis attending a rheumatic clinic we have been able to define a distinct clinical entity for which we suggest the name of primary generalized osteoarthritis. This condition occurs most often in middle-aged women, and is characterized by a distinct pattern of joint involvement, by a course in which each affected joint passes through an initial painful and more or less acute arthritic phase, and by other distinctive clinical and radiological features.

Generalized osteoarthritis is a constitutional disorder affecting the diarthrodial joints and probably represents a severe form of idiopathic Heberden's nodes. A therapeutic trial of oestrogen therapy gave negative results, but patients suffering from this disease may be helped considerably by the proper management of their complaints.

REFERENCES

- Adams, R. (1857). *A Treatise on Rheumatic Gout*. John Churchill, London.
 Addis, H. St. C. C., Jepson, R. P., and Kellgren, J. H. (1950). *Clin. Sci.*, **9**, 271.
 Ball, J. (1950). *Lancet*, **2**, 520.
 Cecil, R. L., and Archer, B. H. (1926). *J. Amer. med. Ass.*, **87**, 741.
 Charcot, J. M. (1881). *Clinical Lectures on Senile and Chronic Diseases*. The New Sydenham Society, London.
 Collins, D. H. (1949). *The Pathology of Articular and Spinal Diseases*. Arnold, London.
 Fletcher, E. (1947). *Medical Disorders of the Locomotor System*. Livingstone, Edinburgh.
 Francon, F. (1950). *Rheumatism*, **6**, 17.
 Haygarth, J. (1805). *A Clinical History of Diseases*. Gadell and Davies, London.
 Heberden, W. (1802). *Commentaries on the History and Cure of Diseases*. T. Payne, London.
 Janus, O. (1950). *British Medical Journal*, **2**, 1244.
 Kellgren, J. H., and Ball, J. (1950). *Ann. rheum. Dis.*, **9**, 48.
 Rose, H. M., Ragan, C., Pearce, E., and Lipman, M. O. (1948). *Proc. Soc. exp. Biol., N.Y.*, **68**, 1.
 Stecher, R. M. (1946a). *J. Lab. clin. Med.*, **31**, 687.
 — (1946b). *Arch. phys. Med.*, **27**, 409.
 — (1948). *Ann. rheum. Dis.*, **7**, 1.

The largest intake of first-year medical students for many years is reported by the Association of American Medical Colleges, who announced that 7,381 freshmen were admitted for the 1951-2 academic year (*New York Times*, January 12). It was also announced that applications for admittance had declined for the second year in succession. The fact that 30% of this year's applicants were applying for the second time is taken to indicate that about one of every two students who apply for admittance is ultimately accepted.