the blood picture, sedimentation rate, or urine, and the only symptoms recorded were slight gastro-intestinal disturbances in two persons.

I am indebted to Dr. H. M. Archibald for his interest and co-operation; to Dr. Robertson at the I.C.I. Unit in Nairobi for the infected mosquito bites; to Dr. J. C. Hawksley, Dr. M. Maizels, and Dr. E. Graves, of University College Hospital, for the final count upon my blood and the sternal puncture; to Dr. J. C. Broom and Dr. S. R. M. Bushby for the determinations recorded in Table II; to the volunteers who submitted to trials with the new drug; and to Brigadier J. S. K. Boyd for his encouragement and advice at all times during the experiments.

REFERENCES

- Archibald, H. M. (1951). *British Medical Journal*, 2, 821.
Falco, E. A., Goodwin, L. G., Hitchings, G. H., Rollo, I. M., and Russell, P. B. (1951). *Brit. J. Pharmacol.*, 6, 185.

HYPNOSIS IN CHILDBIRTH

BY

A. M. MICHAEL, M.B., F.R.C.Sed., M.R.C.O.G.
M.M.S.A.

*Late Senior Registrar, Department of Obstetrics and
Gynaecology, St. Helier Hospital, Surrey*

Considerable interest has been aroused in recent years, both in this country and in the United States, in the phenomenon of hypnosis, with special reference to its application in the relief of pain during childbirth. Unfortunately the attitude of the general public and some medical practitioners towards this phenomenon has been, and too often still is, one of superstition, fear, and even open hostility. No doubt this is the result of witnessing lay demonstrations by medically ignorant and often completely irresponsible people making extravagant claims for hypnosis.

From time to time reports of cases of childbirth while under hypnosis have appeared in the lay press, arousing interest and discussion. An attempt has therefore been made to investigate these claims on a scientific basis, so far as this is possible, and it is with this idea in mind that this preliminary objective evaluation of the effects of hypnosis as an anaesthetic agent in childbirth has been undertaken.

Relief of Pain During Childbirth

In our present civilization, women have been conditioned to expect pain during childbirth, and are convinced that it is a necessary adjunct to labour. Anticipation increases pain largely through muscular and nervous tension. Following on the publication of the work of Grantly Dick Read in educating the patient in the value of relaxation, and in attempting to remove the fear and apprehension of childbirth, efforts are being made to induce the parturient woman to regard childbirth as a natural function which should cause no more discomfort than any other physiological process of the body. Read's technique depends on the use of education, relaxation, and suggestion, and by employing trance states one only carries his method a stage further.

Kroger and De Lee (1943) believe that the only method of producing relief of pain with complete safety for mother and infant, without altering the normal mechanism of labour, is hypnosis. They further maintain "that hypnosis has unrealized possibilities for making childbirth not merely the equivalent of a surgical operation but rather a satisfying psychological experience, which may fulfil a deeply felt and unformulated need of the mother." Greenhill (1947) states that hypnosis should be employed more often than it is at

present. Even if complete hypnosis is not to be resorted to, physicians should remember that repeated suggestion, with or without the aid of medication, can accomplish a great deal in labour, particularly for the relief of fear, as well as the pain of childbirth.

Prior to 1775 the phenomenon of hypnosis was often described as a miracle performed by high priests of ancient civilizations, particularly in the East. The scientific history of hypnosis began with Anton Mesmer. In this country James Braid, in 1841, found that trance-like states could be induced by holding any bright object in front of, and slightly above, the subject, so as to strain the eyes and quickly fatigue them. He first coined the terms hypnosis and hypnotism. During the past 30 years there has been an increasing wealth of publications dealing with the effective use of hypnosis in the fields of psychiatry and experimental psychology. There have been papers advocating the use of hypnosis as an anaesthetic agent in surgery and dentistry. Sampimon and Woodruff (1946) described 29 surgical and dental operations successfully performed under hypnosis while in a prisoner-of-war camp in Singapore during the Japanese occupation.

Hypnosis has been used in obstetrics for many years, and Schultze-Rhönhof (1922) reported an 89.5% success rate in a series of 79 cases. He came to the conclusion that hypnosis was a useful and harmless anaesthetic agent. Isolated enthusiastic reports of labours conducted under hypnosis have appeared in recent years in the medical press, but, with the exception of papers by Kroger and De Lee (1943) and Abramson and Heron (1950), no large series of cases have been reported in the English language. It should be pointed out at this stage that in Abramson and Heron's series of 100 cases the experimental group of patients (receiving hypnosis) were each given a mean amount of 103.5 mg. of pethidine during labour, whereas the control group of patients (not receiving hypnosis) received a mean of 123.6 mg. of pethidine. In the series of 30 cases reported here only those cases which failed completely during labour received anaesthetic agents.

Present Investigation

Material.—All subjects in this series were volunteers and were obstetrically normal at the 34th–37th weeks of pregnancy. No attempt was made to select patients, and only one case was rejected during the antenatal period. This patient could not be hypnotized after three attempts, and, owing to pressure of work, it was not deemed practicable to continue with this case. Vogt, of Berlin, with typical Teutonic thoroughness, persevered with one case through 600 attempts and 700 with another before inducing hypnosis (Le Cron and Bordeaux, 1949). All volunteers were drawn from patients attending the antenatal clinics of this hospital, and no elaborate preparations were made to secure optimum conditions for hypnosis.

Antenatal Preparation.—Patients were seen at varying intervals during the antenatal period in order to hypnotize them and to accustom them to the routine. A semi-reclining position on the examination couch in an un-darkened room was adopted, and in order to secure fixation of attention the gaze was focused on a fixed point

TABLE I.—Analysis of Cases in Which Hypnosis Failed During Labour

Case	Age	Parity	No. of A.N. Training Sessions	Depth of Hypnosis* Obtained During A.N. Sessions	Stage of Labour Hypnosis Failed
A	19	1	10	4	1st
B	16	1	6	3	"
C	33	2	7	3	Beginning of 2nd
D	35	4	3	4	"
E	36	2	11	4	Crowning of head
F	23	2	2	3	"
G	27	2	11	5	"

* Depth of hypnosis referred to in this column is in accordance with Le Cron and Bordeaux's classification.

TABLE II.—Analysis of Cases in Which Hypnosis Succeeded

Case	Age	Parity	No. of A.N. Training Sessions	Depth of Hypnosis* Obtained During A.N. Sessions	Remarks (Behaviour in Labour)
1	27	2	10	3	Excellent
2	31	2	10	5	V. good
3	30	2	5	3	"
4	33	2	7	4	Good
5	29	4	7	4	V. good
6	21	1	6	4	Excellent
7	28	2	8	3	"
8	29	1	9	3	" (episiotomy sutured)
9	40	2	7	4	Excellent
10	19	1	11	4	"
11	33	3	5	3	Good
12	25	2	6	4	Excellent (laceration sutured)
13	29	1	4	3	Excellent (second-degree laceration sutured)
14	27	2	5	3	V. good
15	23	2	9	4	"
16	23	2	9	3	Excellent
17	30	2	9	5	"
18	40	2	5	4	"
19	24	2	6	4	"
20	28	3	5	4	"
21	25	2	9	4	V. good
22	32	2	8	4	Good
23	25	2	10	4	First stage excellent†

* Depth of hypnosis referred to in this column is in accordance with Le Cron and Bordeaux's classification.

† This case was deliberately awakened in the second stage.

on the ceiling. These training sessions varied from two in number in one case to 11 in two others. No attempt was made to standardize the number of training periods given to each patient, and it was found that the number of training sessions per patient had no direct bearing on the resultant behaviour of the subject during labour (Tables I and II). No patient was seen before the 34th week of pregnancy, and one case was seen for the first time as late as the 39th week, 10 days before delivery.

Depth of Hypnosis

Various trance stages are known and fairly easily recognized, although no clear line of demarcation between these stages exists. In spite of many attempts to separate them, no classification has as yet been generally accepted. For general purposes five divisions of the various stages of susceptibility have been found to be quite satisfactory, with three of these designating actual hypnosis (Le Cron and Bordeaux, 1949): (1) insusceptible; (2) hypnoidal—some symptoms but no trance; (3) light trance; (4) medium trance; and (5) deep or somnambulistic trance.

In the training sessions attempts were made to get as deep a degree of hypnosis as possible, and the response differed from case to case because of the great variability in the susceptibility of the subjects. Similarly the depth of hypnosis varied with the experience and ability of the operator. Simple tests were performed in an attempt to gauge the depth of the trance. Here it should be stressed that patients apparently reaching a deep trance during the training sessions did not necessarily prove to be ideal subjects during the actual childbirth (Tables I and II).

Technique of Inducing Hypnotic Trance

This has been comprehensively dealt with in innumerable publications, and it has been found that any acceptable method of trance induction is suitable. During the antenatal period steps should be taken to establish rapport between the patient and those who will be ultimately responsible for her nursing during delivery. This can be quite easily established by careful and sympathetic discussion with the patient while she is in the trance state. It will prove of great value to the obstetrician in charge of the delivery, as his presence at the bedside is thus not required to so great an extent, since the midwife can "reinforce" the subject with a few suggestions of sleep, should she show signs of awakening.

During the training sessions the hypnotized subject is assured that as labour is a physiological process it should be no more uncomfortable than any other function of the body. In addition, the events occurring during a normal labour are explained to the patient. (It is assumed that the patient has already received some instruction in the physiology of the antenatal period. At this hospital, patients receive three lectures on the physiology of pregnancy, labour, and the puerperium.)

At each session post-hypnotic suggestion is given that labour will be painless, and that the only sensation the patient will experience will be a feeling of tightness in the lower abdomen and lower back during the uterine contractions. While under hypnosis the subject is told that she will look forward to the actual confinement with a feeling of confidence rather than one of fear and apprehension. During these antenatal training sessions the subjects are taught to relax, and post-hypnotic suggestion is given that when in labour they would remain relaxed with each uterine contraction. Relaxation is a condition which can be easily induced in the hypnotic state, and is much more complete than that which the subject can otherwise accomplish. After waking from the trance the subject has no recollection of the suggestion which has been given until it arises in her mind at the stated time or occasion.

The application of mass hypnosis during the training sessions has proved invaluable as a time-saving factor. This was introduced to the patients after they had been hypnotized alone on the first two occasions. Thereafter they were hypnotized in batches of six with satisfying results. In most cases it will be found that after the first session hypnosis can be induced merely on command, and thereafter the trance state can be deepened by appropriate suggestion.

Management During Labour

Properly trained, the subject should be able to hypnotize herself when labour begins, or during the training sessions post-hypnotic suggestion can be given to her to the effect that she would enter the trance state at the appropriate time. This, however, is not always satisfactory, as many will do this at the beginning of false labour. A few cases were instructed to enter the trance state only after admission to the labour ward—that is, after they had been seen and attended to by the midwife in the admission room. The most satisfactory method was for the obstetrician to give the command to the subject after she had been examined by him personally, and after he was convinced that labour was well established and the os at least two fingers dilated. At this stage the patient is again told that her sleep will be deep and continuous and that she will hear only the commands of the obstetrician-hypnotist and the attendant midwife. The subject is told to carry out all the instructions given to her by the midwife, and should any examinations (abdominal, rectal, or vaginal) be necessary they would cause no discomfort at all.

The normal mechanism of labour is not interfered with, and patients are found to be very co-operative, especially in the second stage of labour, at which time the expulsion of the baby is aided by the voluntary powers of the subject. If labour is somewhat prolonged, it has been found useful to waken her for short intervals, during which time meals can be taken, and bowels and bladder be attended to. These functions can, however, be carried out even while the patient is in the trance state.

It is advisable for the hypnotist to be present continuously during the second stage of labour, as it was during this expulsive stage that the majority of failures in this series occurred. During this stage, patients are instructed to bear down with each contraction, which they recognize by the tightness in lower abdomen and back. Immediately after each contraction, in this stage, the patients are "reinforced" with a few suggestions of sleep and told to relax completely. When the perineum is being distended by the presenting

part the perineal area is "anaesthetized" by repeated suggestion to the hypnotized patient. In most subjects episiotomies can be performed quite painlessly, and subsequent repair effected without a local or general anaesthetic. After "anaesthesia" has been invoked, it must always be removed and that part of the body returned to a normal condition of sensitivity before waking the patient.

The third stage of labour is also conducted with the patient in the trance state, and she is awakened only after she has been made comfortable following on the expulsion of the placenta.

Kroger and De Lee state that in the management of the prenatal period "post-hypnotic suggestion is given to the effect that the labour will be painless, and that the subjects will have no recollection of the entire procedure." In this series of cases the patients have been kept informed of the progress of labour, and there can thus be full conscious co-operation and participation by the mother in the birth of her child. This is said to be psychologically desirable.

In an attempt to assess the objective rather than the subjective effects of hypnosis in childbirth it was thought advisable to have for comparative purposes a control group of patients. These were obtained, after the experimental cases had been delivered, by referring to the hospital admission book and extracting therefrom details of normal deliveries of women, of similar parity, admitted to hospital immediately after the appropriate experimental cases. Tables III and IV show the mean age and parity of the patients in the two groups.

TABLE III.—Mean Age of Patients (in years)

	Control Group	Experimental Group
Multigravidae (24 cases) ..	27.70	29.96
Primigravidae (6 ") ..	25.50	22.67
Total (30 ") ..	27.27	28.50

TABLE IV.—Mean Parity of Patients

	Control Group	Experimental Group
Multigravidae (24 cases) ..	2.26	2.33
Total (30 ") ..	2.00	2.07

Results

In this series there were six primigravidae and 24 multigravidae. In view of the fact that the duration of labour in a normal case definitely varies according to whether the subject is a primigravida or a multigravida the two groups have been treated independently. Application of "Student's" *t* test shows that only in the second stage of labour in respect of the primigravidae is there a statistically significant difference (at the 5% level) between the control group mean of 88 minutes and the experimental group mean of 37 minutes (Table V). In spite of the fact that statistically, with the above exception, there is no significant difference in the above figures, the difference of the means between the control and experimental groups of both classes of patient indicates a tendency in favour of the treatment of the experimental group (Table V). The reasons for the virtual failure to pass the significance tests are, first, the small series of cases, and, secondly, the very great variations apparent in any single group.

TABLE V.—Duration of Labour

Stage of Labour	Primigravidae		Multigravidae	
	Control	Experimental	Control	Experimental
First ..	Hr. Min. 17 22	Hr. Min. 13 30	Hr. Min. 10 14	Hr. Min. 9 59
Second ..	1 28	0 37	0 29	0 35
Total duration	19 4	14 22	10 58	10 45

First Stage of Labour.—Of the 30 cases in the experimental group, 28 behaved extremely well during this stage of labour while under hypnosis. It must be again stressed that not one of these patients received any analgesia throughout this time. Co-operation was good and relaxation excellent. The two failures were young primigravidae, aged 16 and 19 respectively, who were brought into hospital after many hours of strong labour at home, and who were delivered within three and two hours respectively after admission to the labour ward. All attempts at inducing hypnosis in these two cases failed, as they were being subjected to repeated strong contractions occurring at short intervals.

Second Stage of Labour.—Analysis of the behaviour of the patients in this stage reveals that of the remaining 28 cases 25 were extremely satisfactory, co-operating well and carrying out all instructions. One woman was deliberately awakened at the beginning of the second stage of labour, following on an excellent first stage, as both the obstetrician-hypnotist and the midwife, with whom rapport had been established, were forced to leave her. The other two failures in this stage were both multiparae, one of whom attended seven antenatal training sessions, and the other only three. No difficulty was experienced in hypnotizing these two women during the antenatal period, and no explanation for their failure in this stage can be advanced (Table I).

Finally, three further cases failed during the actual delivery of the infants. Thus of the original 30 cases, 22 (73%) had completely painless labours. A corrected percentage should exclude the patient deliberately awakened, and this figure should then be 76.

Discussion

An attempt has been made to show that painless childbirth can be achieved without resort to analgesia or anaesthesia. In an entirely unselected series of 30 cases, 76% had completely painless labours, without having received any drug whatsoever, and it is felt that with proper selection of cases this figure could be improved upon. Though statistically there was no significant difference in the duration of the first stage of labour between the experimental and the control groups, it was noted that the patients in the former group were most restful and co-operative, and required less attention than the patients not receiving hypnosis. This impression was gained by the nursing as well as the medical staff, and was all the more welcome and valuable because at the beginning of the series they were sceptical, and often openly hostile to the experiment.

Subsequent discussions with the patients taking part in this experiment revealed that they were all very pleased with the experience they had undergone. Even the "second-stage failures" reported that they had had marked relief from pain and discomfort until such time as hypnosis failed; even then they were not unduly distressed by the nature of the uterine contractions, and found that they could relax completely in the intervals between contractions. Only one woman expressed a desire for analgesia in the event of a subsequent pregnancy.

The percentage of individuals that can be hypnotized is disputed by various authors, but in this series only one case was rejected, as she could not be hypnotized after only three attempts. A possible explanation of the high percentage of hypnotizable subjects obtained in this series is that the pregnant woman is in a more than usually suggestible state and so is an exceptionally good subject for hypnosis; also, in addition, volunteers are particularly likely to be suitable patients. One of the greatest difficulties in the practice of hypnotism, however, is the unreliability and unpredictability of its effects.

One possible drawback to the use of hypnosis is the time expended by the hypnotist during the prenatal training period. This, as has already been stressed, can be overcome by the introduction of mass hypnosis. Also, by establishing rapport between the patient and the attendant midwife, the

hypnotist is not required to be present at the bedside during the first stage of labour. The training of the labour ward staff in the management of the hypnotic patient in labour is most important, and can be easily accomplished. The main points to be observed in the handling of the patients are quietness and gentleness, and prior to the performance of any manipulative procedure the patient should be informed of what is about to be done.

Doubt exists in the minds of some about the safety of hypnosis. In a leading article in the *British Medical Journal* (1949) the following statement appears: "There is no evidence that in capable and conscientious hands hypnosis, repeated as many times as is desired, carries with it any physical or psychological danger to the patient." The question of a disturbance in the normal mother-child relationship has been raised, but it is felt that this relationship, rather than being impaired, is actually enhanced, as the mother retains no unpleasant memories of the childbirth, and therefore has no opportunity, either consciously or subconsciously, to blame the infant for any suffering she may have experienced.

Summary

An attempt has been made to show that prenatal training in hypnosis can benefit the parturient woman by assuring relaxation and abolishing the pain of labour.

There is evidence to show that the duration of labour is shortened.

The nursing staff can be easily trained to deal with the hypnotic patient, and it is submitted that this method is both practicable and safe for the mother and infant.

I wish to express my thanks to Miss D. Daley for permission to undertake this investigation in her department. In addition, thanks are due to Miss E. Wilson and to the medical and nursing staffs of the department of obstetrics and gynaecology for constructive criticism and help, and to Mr. Harris Schermann for preparation of the statistical data.

REFERENCES

- Abramson, M., and Heron, W. T. (1950). *Amer. J. Obstet. Gynec.*, 59, 1069.
British Medical Journal, 1949, 2, 427.
 Greenhill, J. P. (1947). *De Lee's Principles and Practice of Obstetrics*, 9th ed., p. 266. Saunders, Philadelphia.
 Kroger, W. S., and De Lee, S. T. (1943). *Amer. J. Obstet. Gynec.*, 46, 655.
 Le Cron, L. M., and Bordeaux, J. (1949). *Hypnotism Today*, 2nd printing. Grune & Stratton, New York.
 Sampimon, R. L. H., and Woodruff, M. F. A. (1946). *Med. J. Aust.*, 1, 393.
 Schultze-Rhonhof, F. (1922). *Zbl. Gynäk.*, 46, 247.

DIABETES MELLITUS IN SURGICAL PATIENTS

BY

C. F. ROLLAND, M.B., Ch.B., M.R.C.P.Ed.

(From the Department of Therapeutics, University of Edinburgh, and the Clinical Laboratory, Royal Infirmary, Edinburgh)

Every week two or three diabetic patients are admitted to the surgical wards of the Royal Infirmary of Edinburgh. Thus the problems which arise in the care of diabetics undergoing surgical treatment are common. This paper discusses the subject from the medical point of view and refers to a series of 300 cases treated during 1948, 1949, and 1950, excluding obstetrical patients and those admitted for extraction of teeth. The management of the diabetes in all these patients has been the responsibility of one medical charge.

Age Distribution

Table I shows the ages of 300 consecutive surgical diabetics arranged in five-year groups. Most of the patients were over 55 years old when they required surgical treatment, and

TABLE I.—Age Distribution

Age Group in Years	No. of Patients	Age Group in Years	No. of Patients
1-5	0	56-60	50
6-10	0	61-65	60
11-15	3	66-70	48
16-20	5	71-75	36
21-25	2	76-80	22
26-30	6	81-85	2
31-35	5	86-90	0
36-40	12	91-95	1
41-45	14	96-100	0
46-50	16		
51-55	18		300

a considerable proportion were over 70. Few were young. These facts have an important bearing on the difficulties to be anticipated in dealing with the surgical diabetic. Degenerative disease, particularly of the cardiovascular system, becomes apparent in many diabetics at an earlier age than in other patients. Such degeneration plays a part in determining the mortality rate and type of treatment required. Many of the more serious problems which arise in the surgical treatment of diabetics are those encountered in the surgery of the aged.

The Surgical Disorders

Table II shows the incidence of some of the conditions which called for surgical treatment in this series of cases. Gangrene of the foot was the most common single disorder,

TABLE II.—Incidence of Conditions Necessitating Surgical Treatment

Cardiovascular disease:		
Gangrene of foot	50	} 55
Hypertension	1	
Intermittent claudication	2	
Angina pectoris	1	
Femoral vein thrombosis	1	
Eye diseases:		
Cataract	40	} 55
Glaucoma	6	
Others	9	
Infections:		
Abscesses	14	} 59
Carbuncles	11	
Cellulitis	6	
Acute otitis media	8	
All others	20	
Trauma:		
Fractured femur	18	} 34
Others	16	
Neoplastic disease: All types	30	
Miscellaneous	67	
	300	

and cataract was the next. There were 59 patients with infective lesions, a considerable proportion of which were of staphylococcal origin. Under the heading "all others" were cholecystitis, appendicitis, pyelonephritis, and similar disorders. Fractures of the femur and neoplastic diseases were common, as they tend to be in elderly patients. The miscellaneous group consisted of a great variety of diseases without any particular connexion with diabetes.

Untreated Diabetics

As many as 41 of the patients in this series were not known to have diabetes mellitus at the time when they were admitted to the surgical wards. Of these 41 undiagnosed diabetics only 5 were under 50 years of age, while 24 were over 60. They suffered from a variety of surgical disorders. Twelve had cardiovascular disease, mostly gangrene of a foot; nine had gynaecological conditions, and six had infective lesions. In two patients the recognition of the diabetes was delayed until diabetic coma developed after operation. In one of these the operation was an emergency procedure for the relief of a strangulated femoral hernia. The other was a planned operation—dilatation of the cervix and curettage of the uterus in a patient who had post-menopausal