hypnotist is not required to be present at the bedside during the first stage of labour. The training of the labour ward staff in the management of the hypnotic patient in labour is most important, and can be easily accomplished. The main points to be observed in the handling of the patients are quietness and gentleness, and prior to the performance of any manipulative procedure the patient should be informed of what is about to be done.

Doubt exists in the minds of some about the safety of hypnosis. In a leading article in the *British Medical Journal* (1949) the following statement appears: "There is no evidence that in capable and conscientious hands hypnosis, repeated as many times as is desired, carries with it any physical or psychological danger to the patient." The question of a disturbance in the normal mother-child relationship has been raised, but it is felt that this relationship, rather than being impaired, is actually enhanced, as the mother retains no unpleasant memories of the childbirth, and therefore has no opportunity, either consciously or subconsciously, to blame the infant for any suffering she may have experienced.

Summary

An attempt has been made to show that prenatal training in hypnosis can benefit the parturient woman by assuring relaxation and abolishing the pain of labour.

There is evidence to show that the duration of labour is shortened.

The nursing staff can be easily trained to deal with the hypnotic patient, and it is submitted that this method is both practicable and safe for the mother and infant.

I wish to express my thanks to Miss D. Daley for permission to undertake this investigation in her department. In addition, thanks are due to Miss E. Wilson and to the medical and nursing staffs of the department of obstetrics and gynaecology for constructive criticism and help, and to Mr. Harris Schermann for preparation of the statistical data.

REFERENCES

- Abramson, M., and Heron, W. T. (1950). *Amer. J. Obstet. Gynec.*, 59, 1069.
British Medical Journal, 1949, 2, 427.
 Greenhill, J. P. (1947). *De Lee's Principles and Practice of Obstetrics*, 9th ed., p. 266. Saunders, Philadelphia.
 Kroger, W. S., and De Lee, S. T. (1943). *Amer. J. Obstet. Gynec.*, 46, 655.
 Le Cron, L. M., and Bordeaux, J. (1949). *Hypnotism Today*, 2nd printing. Grune & Stratton, New York.
 Sampimon, R. L. H., and Woodruff, M. F. A. (1946). *Med. J. Aust.*, 1, 393.
 Schultze-Rhonhof, F. (1922). *Zbl. Gynäk.*, 46, 247.

DIABETES MELLITUS IN SURGICAL PATIENTS

BY

C. F. ROLLAND, M.B., Ch.B., M.R.C.P.Ed.

(From the Department of Therapeutics, University of Edinburgh, and the Clinical Laboratory, Royal Infirmary, Edinburgh)

Every week two or three diabetic patients are admitted to the surgical wards of the Royal Infirmary of Edinburgh. Thus the problems which arise in the care of diabetics undergoing surgical treatment are common. This paper discusses the subject from the medical point of view and refers to a series of 300 cases treated during 1948, 1949, and 1950, excluding obstetrical patients and those admitted for extraction of teeth. The management of the diabetes in all these patients has been the responsibility of one medical charge.

Age Distribution

Table I shows the ages of 300 consecutive surgical diabetics arranged in five-year groups. Most of the patients were over 55 years old when they required surgical treatment, and

TABLE I.—Age Distribution

Age Group in Years	No. of Patients	Age Group in Years	No. of Patients
1-5	0	56-60	50
6-10	0	61-65	60
11-15	3	66-70	48
16-20	5	71-75	36
21-25	2	76-80	22
26-30	6	81-85	2
31-35	5	86-90	0
36-40	12	91-95	1
41-45	14	96-100	0
46-50	16		
51-55	18		300

a considerable proportion were over 70. Few were young. These facts have an important bearing on the difficulties to be anticipated in dealing with the surgical diabetic. Degenerative disease, particularly of the cardiovascular system, becomes apparent in many diabetics at an earlier age than in other patients. Such degeneration plays a part in determining the mortality rate and type of treatment required. Many of the more serious problems which arise in the surgical treatment of diabetics are those encountered in the surgery of the aged.

The Surgical Disorders

Table II shows the incidence of some of the conditions which called for surgical treatment in this series of cases. Gangrene of the foot was the most common single disorder,

TABLE II.—Incidence of Conditions Necessitating Surgical Treatment

Cardiovascular disease:		
Gangrene of foot	50	} 55
Hypertension	1	
Intermittent claudication	2	
Angina pectoris	1	
Femoral vein thrombosis	1	
Eye diseases:		
Cataract	40	} 55
Glaucoma	6	
Others	9	
Infections:		
Abscesses	14	} 59
Carbuncles	11	
Cellulitis	6	
Acute otitis media	8	
All others	20	
Trauma:		
Fractured femur	18	} 34
Others	16	
Neoplastic disease: All types	30	
Miscellaneous	67	
	300	

and cataract was the next. There were 59 patients with infective lesions, a considerable proportion of which were of staphylococcal origin. Under the heading "all others" were cholecystitis, appendicitis, pyelonephritis, and similar disorders. Fractures of the femur and neoplastic diseases were common, as they tend to be in elderly patients. The miscellaneous group consisted of a great variety of diseases without any particular connexion with diabetes.

Untreated Diabetics

As many as 41 of the patients in this series were not known to have diabetes mellitus at the time when they were admitted to the surgical wards. Of these 41 undiagnosed diabetics only 5 were under 50 years of age, while 24 were over 60. They suffered from a variety of surgical disorders. Twelve had cardiovascular disease, mostly gangrene of a foot; nine had gynaecological conditions, and six had infective lesions. In two patients the recognition of the diabetes was delayed until diabetic coma developed after operation. In one of these the operation was an emergency procedure for the relief of a strangulated femoral hernia. The other was a planned operation—dilatation of the cervix and curettage of the uterus in a patient who had post-menopausal

bleeding. Both of these patients recovered. A third patient, admitted because she suffered from pyaemia with multiple abscesses, developed diabetic coma before the diabetes was recognized. There was no operation in this case. She died after the ketosis had been corrected and the blood sugar had returned to normal levels.

The discovery of these 41 hitherto undiagnosed diabetics emphasizes the necessity of testing the urine for sugar soon after the admission of any patient to a surgical ward, whatever the surgical disorder may be. The fact that even a minor operation, with a short period of anaesthesia, may precipitate diabetic coma in an unprepared diabetic makes this routine as essential as the recording of the name and address of the patient.

In these 41 patients the diagnosis of diabetes presented no difficulties. In most cases there was a history of typical symptoms, such as thirst, polyuria, pruritus vulvae, or loss of weight, together with the finding of glycosuria and sometimes ketonuria. In patients with glycosuria who did not have a typical history the finding of a fasting blood sugar higher than 120 mg. per 100 ml. was diagnostic. In none of these cases was a glucose-tolerance test necessary to confirm the diagnosis.

The Fatal Cases

Of these 300 patients 28 died during their stay in the hospital, making the overall mortality rate 9.3%. It is important to note that this figure includes all the deaths and not only those immediately attributable to the operation. In fact, in five of the fatal cases there was no operation (Table III), usually because the patient was considered to be *in articulo mortis*. Only two of the patients died within the

first day of the post-operative period. In three cases the time intervening between operation and death was as long as 62, 86, and 96 days.

Most of the patients who died were old. Fifteen of them were over 70 and only two were under 58. Most old diabetics suffer from degenerative vascular disease, and evidence of this was found in all patients upon whom post-mortem examinations were made. The presence of such disease certainly increases the risks of the post-operative period. Thus one patient died of cerebral thrombosis and at least four others of myocardial infarction. Six deaths occurred in the group of patients whose diabetes was undiagnosed at the time of their admission to hospital.

Nine of the fatal cases had been treated for gangrene of a foot, seven for malignant disease, and six for fracture of a femur. Thus three conditions which normally carry a high mortality rate in non-diabetic patients accounted for 22 of the 28 fatalities in this series. In only one case did diabetic coma play any part in the fatal issue, and even here its importance was uncertain, as at the time of death the ketosis had been corrected, the blood sugar was 140 mg. per 100 ml., and the patient had a severe staphylococcal pyaemia.

A death rate of 9.3% may seem to be high, but it compares quite favourably with the figure of 6.95% which was the overall mortality rate in the 1,007 patients admitted to one of the general surgical charges in the hospital during 1949. The average age of this latter group was considerably less than that of the diabetic patients. The factors of most importance in the deaths of the diabetic patients were: (1) their age, the majority being over 70; (2) the presence of degenerative vascular disease, which was no doubt accelerated if not caused by the diabetes; and (3) the incidence of serious disorders such as gangrene and malignant disease. It is particularly striking that diabetic coma, which used to be responsible for so many deaths in the pre-insulin era, is now of minor importance. It is in fact a preventable complication.

Management

The management of the diabetes in these patients was in the hands of the medical charge which is responsible for the diabetic out-patient clinic. The sister in charge of the diet kitchen was responsible for the patients' diet, and their urine specimens were tested in the side-room of the metabolic ward to which the diet kitchen is attached. This arrangement has many advantages. In particular, control is centralized and the easy availability of out-patient records provides valuable information about the patients and their previous treatment. At all times a close friendly co-operation was maintained with the surgeon in charge of each case. It is essential that the physician and the surgeon should understand each other's intentions and difficulties, and this co-operation was the most important single factor in the successful treatment of these patients.

The aim of the medical staff was to arrange the control of the diabetes in such a way as to permit the surgeon to treat his patient as he would wish to do if no diabetes was present. In the great majority of cases this aim was achieved so long as reasonable notice of the surgical programme was received. In only one case was a planned operation postponed because of an unexpected increase in the severity of the diabetes on the morning appointed for the operation.

Pre-operative Preparation.—In most cases the operations were not emergency procedures. The time before the operation was occupied in the assessment and control of the diabetes. In most cases the normal diet was continued until the day before the operation. When patients were taking protamine-zinc insulin in doses of over 20 units the insulin treatment was rearranged so as to control the diabetes with two or three doses of soluble insulin daily for two days before the operation. When patients were taking less than 20 units of protamine-zinc insulin daily this dose was continued until the day of the operation, when soluble insulin was substituted. This arrangement was designed to avoid the risk of hypoglycaemia, which is incurred when a patient

TABLE III.—*Details of Fatal Cases*

Sex and Age	Diagnosis	Operation	Interval Between Operation and Death	Cause of Death
M 78	Gangrene of foot	Mid-thigh amputation	39 days	Bronchopneumonia
F 67	" " "	" " "	86 "	Myocardial infarct
F 64	Secondary carcinoma liver	Laparotomy	5 "	Cholaemia
M 78	Fracture neck of femur	Nil	—	Shock
F 75	" " "	" " "	—	Pulmonary embolus
F 28	Appendicitis	Appendicectomy	5 days	Thyrototoxic crisis
F 76	Gangrene of foot	Mid-thigh amputation	14 "	Myocardial infarct
M 62	Carcinoma rectum. Perforation colon	Laparotomy	2 hours	Peritonitis
F 69	Gangrene of foot	Mid-thigh amputation	2 days	Bronchopneumonia
F 64	Carcinoma ampulla of Vater	Transduodenal resection	20 "	Hepatic failure.
F 73	Fracture neck of femur	Reduction	4 "	Duodenal fistula Bronchopneumonia
M 58	Thrombosis deep veins legs	Nil	—	Pulmonary embolus
F 70	Fracture neck of femur	Smith-Petersen nail	18 days	Bronchopneumonia
F 83	Fracture shaft of femur	Steinmann's pin	23 "	" "
M 79	Intestinal obstruction	Nil	—	" "
M 38	Carcinoma pancreas	Cholecyst-gastrostomy	18 days	Cholaemia
M 73	Gangrene of foot	Mid-thigh amputation	3 "	Pulmonary oedema. Bronchopneumonia
F 73	" " "	" " "	96 "	Myocardial infarct
F 76	" " "	Below-knee amputation	36 "	Bronchopneumonia
F 76	" " "	Mid-thigh amputation	12 hours	Cerebral thrombosis
F 61	Staphylococcal pyaemia	Nil	—	Pyaemia. Ketosis
F 66	Carcinoma pancreas	Cholecyst-gastrostomy	27 days	Hepatic failure
F 70	Fracture neck of femur	Smith-Petersen nail	62 "	Bronchopneumonia
F 70	Abscess anterior abdominal wall	Incision	19 "	Pyaemia
F 61	Carcinoma stomach	Laparotomy	20 "	Carcinomatosis
F 72	Carcinoma rectum	Colostomy	14 "	Myocardial infarct
M 58	Pyloric stenosis	Partial gastrectomy	7 "	Peritonitis
M 65	Gangrene of foot	Mid-thigh amputation	7 "	Pulmonary embolus

comes to operation with a large depot of long-acting insulin. In infected cases, antibiotic treatment, usually with penicillin, was administered during the pre-operative days to minimize the infection and prevent its spread.

The Day of Operation

Opinions in the standard textbooks on this subject differ over the best method of controlling the diabetes on the day of the operation. Some workers starve the patient and give no insulin before the operation (Barach, 1949); others give protamine-zinc insulin in a reduced dose to those patients who have been accustomed to take it (Wilder, 1941; Root, 1946); yet others give a small carbohydrate feed together with a dose of soluble insulin at a suitable interval before the operation (Lawrence, 1950). In this series the latter plan was adopted; the soluble insulin was given about four hours, and the carbohydrate about three and a half hours, before the operation. The object was to send the patient to the operating theatre with a reasonably normal carbohydrate metabolism and an adequate store of glycogen in his liver. For the carbohydrate feed many clinics use a solution of glucose flavoured with orange juice. Some diabetics unaccustomed to sweet foods find this nauseating, and this minor disadvantage was obviated by the use of Benger's food with lactose in milk. This proved very satisfactory and there were no complaints of vomiting during anaesthesia. In patients who were to have gastric operations an equivalent amount of carbohydrate was given by intravenous injection of glucose.

Soluble insulin was preferred for the pre-operative dose because of the shorter duration of activity and the greater flexibility of treatment which it permits. Hypoglycaemia due to soluble insulin is more easily recognized and more rapidly corrected than that due to protamine-zinc insulin. Patients who had been controlled by diet without insulin were given a small dose, about 12 units of soluble insulin, with the pre-operative feed, to help them to weather the stress of the operation. The estimation of the correct pre-operative dose for those who require insulin for everyday treatment may be a difficult problem. It is essential to study the individual patient from all points of view rather than to attempt to work out and follow a routine. In this series the dose was usually from one-half to two-thirds of the usual morning dose of soluble insulin, reckoned on the basis of control by two daily doses of this type of insulin. On the one hand, it is important to give enough to prevent the occurrence of ketosis in the immediate post-operative period, and, on the other hand, particularly in the older patients, it is dangerous to risk the complication of hypoglycaemia, which is difficult to recognize in the anaesthetized and sedated patient.

Post-operative Period

During this period the diet was modified to suit the surgical condition. The usual practice was to give a light fluid diet on the evening of the day of the operation and then according to the following diet sheet:

First Day (Carbohydrate, 150-165 g.; calories, 1,300-1,400)

Breakfast.—Four tablespoonfuls strained porridge. Quarter-pint milk (150 ml.) from allowance. Tea with milk, if desired.

Mid-morning.—Quarter-pint (150 ml.) fruit juice plus $\frac{1}{2}$ oz. (10 g.) glucose.

Lunch.—3 oz. (90 g.) puréed fruit or fruit juice as jelly. Cereal pudding made with quarter-pint milk (150 ml.) from allowance, plus $\frac{1}{2}$ oz. (10 g.) cereal. Milk from allowance to drink.

Tea.—One egg, lightly boiled, poached, or scrambled. 1-1½ oz. (30-45 g.) bread with butter from allowance. Tea with milk from allowance.

Supper.—Benger's food, made with quarter-pint (150 ml.) milk from allowance plus $\frac{1}{2}$ oz. (15 g.) Benger's. 1-1½ oz. (30-45 g.) bread with butter from allowance.

10 p.m.—Remainder of milk to drink.

Allowance for Day.—1½ pints (900 ml.) milk; $\frac{1}{2}$ oz. (15 g.) butter. Two plain biscuits—for example "rich tea"—can be substituted for 1 oz. (30 g.) of bread.

Second Day

1 oz. (30 g.) of bread is added to breakfast.

1-1½ oz. (30-45 g.) easily digested meat, plus 1½ oz. (45 g.) mashed potato, added to lunch (carbohydrate, 175 g.; calories, 1,500).

From the third day the amount of meat, vegetable, and bread is increased daily until the diet reaches normal proportions.

In certain cases it was necessary to supply carbohydrate by intravenous infusion of glucose solutions for the first day or two after the operation, but this was restricted to patients in whom the surgeon wished to avoid oral feeding. Until a normal diet was taken soluble insulin was used and long-acting insulin was avoided. When the patient was able to take his usual food protamine-zinc insulin was recommenced.

The Anaesthetic

When ether was the standard general anaesthetic, and even more so in the days of chloroform, diabetics fared badly. The dangers of ketosis were accentuated by liver damage and the not infrequent post-operative vomiting. In consequence it was usual to recommend that spinal or regional anaesthesia should be employed if at all possible and that nitrous oxide and oxygen should be given preference, where feasible, if inhalation anaesthesia was essential. Most of the patients in this series were anaesthetized with intravenous thiopentone, and the anaesthesia was maintained with cyclopropane and oxygen. Curariform drugs were used when relaxation was especially important. A few patients had spinal anaesthetics. The results were remarkably satisfactory. Post-operative vomiting was rare and most of the patients made rapid and uncomplicated recoveries from the anaesthetic. General anaesthesia of this type is so satisfactory that there is no longer any advantage to be gained by the use of regional anaesthesia in diabetic patients.

Emergency Operations

In general the same principles were observed in emergencies as in planned operations. Sometimes there was difficulty because little information was available about the patient's previous treatment, and special problems were encountered when the diabetes was poorly controlled at the time the need for operation arose. Under these circumstances, and especially if the patient was ketotic, the operation was sometimes delayed for a few hours to allow energetic treatment with soluble insulin to effect an improvement in the control of the diabetes. In some such cases the insulin was given in part by the intravenous route for more rapid action. It was at these times that the value of close co-operation between the physician and the surgeon was greatest. When the surgeon could countenance no delay in the operation, and the patient was ketotic, the physician had to do the best he could. Large doses of soluble insulin, regulated by frequent blood-sugar estimations, and the use of intravenous infusions of normal saline proved successful in the few cases of such gravity encountered in this series.

Surgeons and anaesthetists often ask for a system that will enable them to care for diabetics in emergencies when no physician is at hand to help. From what has been said it is evident that it is impossible to devise any mathematical or rule-of-thumb method to suit all diabetics who require operation. There are so many factors to be considered, and the treatment has to be flexible so that it can be varied at short notice to meet the rapidly changing situation. When such occasions arise it seems wise that the anaesthetist should take a share of the responsibility for the diabetes, to free the surgeon for his own work. Soluble insulin should be used. It should be given in large doses (40 to 100 units) if the patient is ketotic. The doses should be regulated by blood-sugar estimations and urine tests. An intravenous infusion of saline should be used if the patient is dehydrated, and the stomach should be washed out if he has been vomiting. If the diabetes is reasonably well controlled, soluble insulin

should be given in small doses (8 to 20 units) every four hours on the basis of urine examinations. It may be necessary to give such patients 20 g. of glucose by intravenous injection with the pre-operative dose of insulin, and fluid feeds containing 20 g. of carbohydrate every four hours for the first day after the operation.

Special Difficulties

Diabetic ketosis is often complicated by symptoms which suggest the presence of an acute abdominal emergency. These symptoms usually subside as the ketosis is corrected. Two young patients in this series had unnecessary laparotomies performed on account of the occurrence of symptoms suggestive of acute appendicitis while they were ketotic. Such a situation may give rise to grave diagnostic difficulties. The patient is often rather uncooperative and there may be physical signs such as abdominal tenderness and rigidity. The presence of a leucocytosis is of no assistance, as there is nearly always a very high white blood count in diabetic coma. In such cases the history is of more value than the examination. If it is found that thirst, polyuria, and drowsiness preceded the abdominal symptoms it is likely that ketosis is responsible for the whole clinical picture, but when the symptoms appear in the reverse order it is more likely that the ketosis is the result of an acute abdominal emergency.

Conclusions

Modern developments have improved the prospect for the surgical diabetic to such an extent that the overall mortality rate in such patients is little higher than that in non-diabetics. Degenerative vascular disease is common in diabetics and gives rise to most of the serious difficulties encountered. The diabetes of a considerable number of patients admitted to surgical wards was undiagnosed before admission, and the recognition of the disorder by routine urine-testing is of paramount importance before such patients are subjected to operative treatment. General anaesthesia is now so satisfactory that there is no special need for regional anaesthesia in the diabetic patient. Full co-operation between the surgeon and a physician who is specially interested in diabetes is of the greatest importance in the successful management of surgical diabetics.

Summary

The problems of the control of diabetes in surgical patients are discussed.

Details are given of a series of 300 consecutive surgical diabetics.

The majority of the patients were over 50 years of age.

Gangrene of the foot was the most common single disorder.

Forty-one of the patients were not known to be diabetic when they were admitted to the surgical wards.

The overall mortality rate was 9.3%.

Diabetic coma was unimportant as a cause of death.

My thanks are due to Professor D. M. Dunlop for his encouragement and help in the preparation of this paper; to the surgeons of the Edinburgh Royal Infirmary for their co-operation and permission to use details of their cases; to Sister E. M. Mutch for her invaluable assistance; and to many others who have helped in various ways.

REFERENCES

- Barach, J. H. (1949). *Diabetes and its Treatment*, pp. 158, 162. New York.
Lawrence, R. D. (1950). *The Diabetic Life*, p. 150. London.
Root, H. F. (1946). In E. P. Joslin's *The Treatment of Diabetes Mellitus*, p. 679. London.
Wilder, R. M. (1941). *Clinical Diabetes Mellitus and Hyperinsulinism*, p. 213. Philadelphia and London.

INCIDENCE AND TYPE DISTRIBUTION OF CAPSULATED *H. INFLUENZAE* STRAINS

BY

BETTY DAWSON, M.B., Ch.B.

AND

K. ZINNEMANN, M.D.

(From the Bacteriology Department, School of Medicine,
Leeds)

In 1945 Straker reported on the incidence of capsulated *Haemophilus influenzae* strains in healthy children. She found that the incidence varied from 8 to 19% in the individuals examined. Little is known about the proportion in which each of the six known types of *H. influenzae*, a-f, contribute to the carrier rate in this country. Straker merely stated that some of the strains isolated from children belonged to type b, and that in 60 adults swabbed regularly for four years capsulated strains were found occasionally, but that type b was absent. Platt (1937) examined 86 nasopharyngeal strains and identified three of these as type e and one as type a. There is a certain element of doubt regarding his findings, as his typing results were obtained with purified fractions of the capsular substance, while his experiments when employing the usual precipitino-agglutination technique gave confusing results. Apart from these two reports no investigations have been carried out on the incidence of the six *H. influenzae* types in this country. Ounsted (1950, 1951) suggested that *H. influenzae* meningitis in children arises from infection with type b strains, the virulence of which has been increased by several previous passages through partially immune older sibs. In view of this interesting hypothesis exact knowledge of the incidence of the *H. influenzae* types has gained an added importance.

Present Investigation

The investigation was undertaken with a view to obtaining data on the occurrence and distribution of the six *H. influenzae* types in the nasopharynx of children in the Leeds area. The recent M.R.C. inquiry into the effectiveness of different prophylactic whooping-cough vaccines provided us with a convenient source of per-nasal swabs from children with upper respiratory infections. These children were either whooping-cough suspects to be examined bacteriologically or their contacts.

Technique.—The children were seen by Dr. G. R. Baxter, who is in charge of the immunization centre at Leeds, and he took nasopharyngeal mucus by means of per-nasal swabs mounted on a flexible curved and non-corroding wire, with a view to isolating *H. pertussis*. The swabs were delivered to the laboratory with the least possible delay and were inoculated at once to Bordet-Gengou plates containing 5 units of penicillin incorporated in 15 ml. of medium. From these plates subcultures were made after three to four days' incubation at 37° C. to 10% heated blood (chocolate) agar plates, and any *H. influenzae* strains detected in this way were examined for their colonial appearance. Strains with mucoid colonies thus isolated were examined for their agglutinability by means of the slide-agglutination technique with specific type sera a-f, and, if positive with one of these sera, the Neufeld test ("capsule swelling") was carried out with the homologous and the five heterologous sera.