REFRESHER COURSE FOR GENERAL PRACTITIONERS

^ FOOD-POISONING_X

BY

^ JOAN TAYLOR, M.B., B.S., B.Sc., D.P.H.

Director, *Salmonella* Reference Laboratory, Colindale

Food-poisoning is a difficult term to define, but through common usage it is accepted as meaning an acute intestinal disturbance coming on within a few hours of the consumption of unwholesome food or drink. It is distinguished from other food-borne infections such as enteric fever, dysentery, or primary abdominal tuberculosis by the occurrence of acute gastro-enteritis, though intermediate cases do occur in which the differential diagnosis, particularly from that of dysentery, may cause difficulty.

Food-poisoning may be due to injurious chemical substances, poisonous plants, or harmful bacteria or bacterial products in food. The most usual type of food-poisoning encountered in this country is bacterial in origin, being caused by organisms of the *Salmonella* group. The disease is due to the multiplication of these organisms in the intestinal tract of the host and is known as the "infective type." Food-poisoning due to the ingestion of bacterial products is also fairly common. Some bacteria while growing in food are able to produce toxic substances which on ingestion cause vomiting and diarrhoea; this disease is known as the "toxin type" of food-poisoning. In a report on food-poisoning in England and Wales during 1950, of 3,979 recorded incidents, 2,170 were of bacterial origin (2,021 salmonella infection), 2 chemical, and 1,807 of unknown causation.

Infective Type

The *Salmonella* group of organisms, which now comprises about 250 recognized species, has its natural reservoir in birds, mammals, and reptiles. The organisms multiply in the intestinal tract of man, causing swelling of the mucosa of the stomach and small intestine and the production of a slimy exudate. The mucosa may become ulcerated. In severe cases these lesions may extend to the large intestine, and there may be an accompanying septicaemia. It is believed that a large number of salmonellae must be ingested in order to give rise to clinical symptoms. The organisms are widely distributed in nature and may be transferred to food in a variety of ways: the food may come from an infected animal, or may be contaminated with infected faeces from mice, rats, pets, or other animals; flies may transfer infected material to food; or human carriers may infect food during handling. Carriers sometimes give a history of a recent attack of diarrhoea, but many have had no intestinal disturbance nor have they been contacts of known cases.

As in most instances many bacteria or large quantities of their products must be ingested to cause symptoms, the most dangerous foods are those which are good culture media and which remain at a warm temperature long enough for bacterial multiplication to take place. These conditions are provided by meat dishes, gravies, and soups, particularly if reheated some time after preparation, and also by foods such as custards prepared at low temperatures. Most tinned foods are autoclaved during processing and are sterile, but once opened they are as liable to contamination as other foods. Small tins of ham can be autoclaved, but large tins are pasteurized, as autoclaving results in loss of palatability and changes in appearance. As a result, pathogenic

organisms may be present in tinned ham, particularly in the large tins, and therefore it is wise to keep the tin in the refrigerator to limit multiplication and not to store it for too long. Hens' eggs are rarely infected, but a number of cases of food-poisoning have been traced to the consumption of ducks' eggs. The latter should be placed in boiling water for at least 10 minutes before being eaten and should not be used in the preparation of lightly cooked foods.

A number of other organisms are believed to cause food-poisoning, though their mode of action is unknown. Foods heavily contaminated with faecal organisms such as *Bact. coli* and *Proteus* have caused disease though the organisms themselves have not been shown to be pathogenic or to produce toxins.

One particular type of food-poisoning, caused by the eating of meat dishes which have been cooked the previous day and allowed to cool down overnight, is seen especially in those eating canteen meals, and is apparently caused by the anaerobic organism *Clostridium welchii*.

Toxin Type

Food-poisoning due to bacterial toxins is most commonly caused by the toxin produced by *Staphylococcus pyogenes*. The toxin is formed in the food during the growth of this organism. It is not destroyed by ordinary cooking processes, so that an infected food which is recooked may still cause disease. The food is usually contaminated by a food-handler who may have a septic lesion on the hands or may transfer the organism from his nose or throat via his hands to the food. Food which has been stored before consumption is the most common source of this disease. Ice-cream, cheese, as well as "made-up" reheated foods, are usually the cause. Other organisms such as non-haemolytic streptococci are known to cause the toxin type of food-poisoning. Little need be said about botulism. This toxin type of food-poisoning is extremely rare in Great Britain, only four incidents ever having been recorded. The disease is caused by the toxin of *Clostridium botulinum*, which acts on the central nervous system, causing diplopia, ocular paresis, and sometimes aphasia accompanied by vomiting and constipation.

Chemical Poisons

Food-poisoning may also be due to injurious chemical substances in the food, though such cases are comparatively rare. There are, however, many instances on record in which food has been in containers which have been used for chemical poisons. Some drinks may be contaminated with antimony or zinc as the result of placing acid liquids—for example, lemonades—or acid stewing-fruits in cheap enamel or zinc vessels. The symptoms usually develop very soon after drinking the fluid.

Death Cap

Some foods are poisonous in themselves. Poisoning due to the "death cap" (*Amanita phalloides*), a fungus eaten in mistake for mushrooms, is well known. The mortality rate is high, and children are particularly susceptible. The incubation period is between 4 and 24 hours, when the patient complains of very acute abdominal pain, vomiting, and diarrhoea. The vomit and faeces may contain blood and mucus. In severe cases anuria may follow, and later jaundice and cyanosis.

Allergy

Diarrhoea and vomiting may occur in individuals who have idiosyncrasies to certain foods. These attacks rarely last more than 24 hours. There is an acute onset soon after the food is eaten. Abdominal pain may be severe, but there is no rigidity. Though this syndrome may occur at any age, it is more common in childhood. Sometimes there is a history of other allergic manifestations in the patient or his family. The diarrhoeal stool from these patients usually contains eosinophil cells. The symptoms are cured by the injection of 3 to 5 min. (0.18 to 0.3 ml.) of adrenaline (1:1,000).

Investigation of Suspected Cases

In all suspected cases of food-poisoning it is important that the patient and the suspected food should be investigated as soon as possible, as knowledge of the causal agent may determine the treatment of the patient and the control measures to stop the spread of the disease. The patient's faeces, collected preferably during the acute stage of the disease and before starting treatment, should be sent to a laboratory. Laboratories provide sterile screw-capped receptacles containing spoons with which a portion of faeces may be transferred to the receptacle. If faeces are not available a sterile throat swab should be passed into the rectum, and care taken to see that the swab shows faecal staining; no lubricant or antiseptic should be used. Such swabs, which are suitable for bacteriological purposes only, must be transported to the laboratory before they begin to dry. Vomit may also be sent to a laboratory, although it rarely assists the investigation. Organisms of the *Salmonella*, *Shigella*, and staphylococcal groups are usually readily isolated if no treatment has been started. Suspected food, if available, should always be sent to the laboratory, as the proof that a particular food has caused disease depends on the correlation of the findings in the patients with those in the food. The assessment of the role of any particular organism isolated from food, other than the accepted pathogens such as salmonellae, shigellae, and some types of staphylococci, is extremely difficult.

Clinical Picture

The diagnosis of food-poisoning when a number of persons are affected usually presents few difficulties, but the diagnosis of an individual case may be far from easy. It should be borne in mind that an accusation of food-poisoning is a common form of blackmail brought against expensive restaurants. In such cases the doctor should never give a certificate to say that the patient has suffered from food-poisoning without the fullest supporting bacteriological and epidemiological evidence.

The symptoms in any single outbreak vary considerably in severity: some patients complain of no more than a vague discomfort which is disclosed only on questioning, while others are severely ill. These latter range from those whose main symptoms are diarrhoea and vomiting to those in whom these symptoms may be overshadowed by a clinical picture more closely resembling that of enteric fever. The symptoms vary with the agent which has caused the food-poisoning, but only the more usual types encountered in this country are dealt with. The most common type is that due to the bacterial infection of food and drink.

The incubation periods of the different types of food-poisoning may give some indication of the cause. Chemical poisons are usually rapid in their effect, symptoms occurring with a sudden onset in about half an hour or less. The toxin type, being due to the absorption of preformed bacterial toxin, also has a short incubation period, though it is longer than that of chemical poisons. The incubation period of the toxin type is between 2 and 12 hours, and there is also a sudden onset. Food-poisoning caused by the multiplication of organisms in the intestine has a longer

incubation period, usually between 12 and 48 hours. The onset of this type may be sudden, though a number of patients complain of feeling ill before the main symptoms occur.

*Incubation Period of Food-poisoning**

Chemical	Bacterial Toxin		Infectious
	Staphylococcus	<i>Cl. welchii</i> and Streptococci	
0-30 minutes	2 hours	12 hours	20 hours

In all types the patient usually complains of nausea, vomiting, and diarrhoea. Commonly there is also headache, abdominal cramps, sweats, shivering, and giddiness. At the height of the illness the patient may be severely prostrated. In the toxin type the disease is of sudden onset and may be very similar to sea-sickness: the more severely ill patient is in a state of collapse and unable to stand; he sweats, and does not care whether he lives or dies. These patients commonly have a subnormal temperature; it is rarely above 99° F. (37.2° C.). They recover rapidly, usually in 24 hours.

In the infective type the disease develops rather more slowly and is accompanied by a rise in temperature, commonly to 102° F. (38.9° C.). The abdominal pain is of a colicky nature, starting usually in the epigastrium but spreading over the whole of the abdomen. Though the abdomen is tender there is no rigidity. The disease usually lasts for about four days and the patient complains of lassitude for some days more. The severely ill patient may look drawn and show signs of dehydration. If death supervenes it is usually about the tenth day of the disease, but death is unusual except in the very young, the old, or those with some additional serious illness.

The vomit is watery; it may be bile-stained, but is only tinged with blood if vomiting is excessive. The stools are watery, usually with faecal colouring, and they are very offensive. Blood is not often present and the mucus tends to be slimy, whereas in dysentery pus and blood are usual. In salmonella infections some patients may show the rose spots and the clinical picture of enteric fever. Patients generally recover rapidly. A certain number of the infective type may be operated upon as cases of appendicitis. It is probable that in these the appendix is inflamed, but the symptoms usually subside as the patient recovers, without necrosis of the appendix.

Complications may occasionally be encountered with salmonella infections. Very rarely empyema, purulent arthritis, or osteitis occur. Meningitis may be found more particularly in young babies. Occasionally patients have a long febrile period sometimes of a septicæmic type. After an attack of food-poisoning some patients become prone to dyspepsia and intestinal upsets.

Differential Diagnosis

Correct diagnosis is important from the point of view both of treatment and of preventing fresh cases. In dysentery vomiting is seldom a major symptom and rarely lasts for long. In addition the stool is usually blood-stained and contains large numbers of pus cells. Intestinal obstruction, intussusception, or volvulus usually cause localizing signs in the abdomen which are absent in food-poisoning. In practice these conditions are rarely confused with food-poisoning.

A disease which occurs in this country is extremely difficult to distinguish from food-poisoning of the types already described. This condition, which occurs more commonly in winter and spring, cannot be traced to any particular food. It may start by affecting a number of persons at the same time, and when institutional outbreaks occur

*From *Modern Trends in Gastro-enterology*, edited by F. Avery Jones. Butterworth and Co., London.

secondary cases are the rule, usually with an incubation period of three to five days. A good review of this disease has been given by Hargreaves,* who refers to it as "epidemic diarrhoea and vomiting," though its more usual name is "the winter vomiting disease." In America similar outbreaks have been studied, and as the result of various investigations it seems probable that a filterable agent, probably a virus, is excreted in the faeces. The onset of the disease is sudden and not uncommonly occurs at night. The main symptoms are vomiting and diarrhoea, but vertigo may be a prominent symptom. Some patients have complained of acute upper abdominal pain, and some tenderness is elicited on deep palpation over the liver area. The differential diagnosis between these patients and true cases of food-poisoning presents many difficulties.

Treatment

Rest in bed is essential in all definite cases of food-poisoning, as recovery is always more rapid and relapses less common if this is observed. The treatment should be symptomatic. If poisoning by chemicals or fungi is suspected the patient's stomach should be washed out with the specific antidote if the offending chemical is known, otherwise with normal saline. In the early stages of other types of food-poisoning patients who are vomiting should be given a glass of warm water containing one teaspoonful of sodium bicarbonate in order to get rid of excessive mucus. If as the result of vomiting and diarrhoea the patient becomes dehydrated intravenous glucose saline should be administered. Persistent diarrhoea may be treated with bismuth carbonate or subnitrate. Opium in any form should be used only if there is no doubt about the diagnosis.

A specific antiserum for poisoning due to *Amanita phalloides* is available.† This serum should be injected intramuscularly and given as early as possible in the disease. The dose for a child is about 20 ml.

There are no specific remedies for bacterial food-poisoning. Antibiotics have been tried, but they have not given good results in the ordinary disease in which diarrhoea and vomiting are the main symptoms. Rarely some cases caused by salmonella infection may have a long-continued fever suggestive of septicaemia. In such cases chloramphenicol should be tried, as some patients seem to improve clinically; the drug does not, however, appear to shorten the period during which the organisms are excreted in the faeces, nor does it have any effect on the carrier condition. The usual dose of chloramphenicol for adults is 50 mg. per kg. of body weight in 24 hours. It is given at intervals of six hours, and should be continued until symptoms have been controlled, after which the dose should be halved for a further three to four days before stopping treatment. During the convalescent period the change to normal diet should be made gradually, particularly in patients who show a tendency to relapse.

Prevention of Spread

In salmonella food-poisoning the patient's excreta should be treated as in typhoid by the addition of 1:20 carbolic acid, corrosive sublimate, or other disinfectants which are known to act on Gram-negative organisms. The patient should be warned of the possibility that the organisms may be excreted for some time after recovery and told to wash his hands after defaecation and before taking food. He should not be allowed to handle food until laboratory tests have shown that the organism is no longer being excreted.

The prevention of bacterial food-poisoning depends on the education of the public in personal hygiene, in clean

methods of food storage, and in clean methods of food preparation. Emphasis should be placed on the desirability of keeping food cool, preferably in a refrigerator, until it is ready for cooking; if it is not eaten immediately after cooking it should be rapidly cooled down and kept cool until it is reheated.

Under Section 17 (1) of the Food and Drugs Act, 1938, it is compulsory for a registered medical practitioner who becomes aware of or who suspects that a patient whom he is attending is suffering from food-poisoning to notify immediately the medical officer of health of the district, giving particulars of the case. If this is done early, laboratory investigations may determine the cause of the disease, the food at fault, and the source of the contamination. Such information will help to stop the spread of the disease.

Next Refresher Course Article.—"Tenosynovitis" and Tendovaginitis," by Mr. D. Lloyd Griffiths.

MEMORANDUM ON FAMILY PLANNING, WITH PARTICULAR REFERENCE TO CONTRACEPTION*

The subject of family planning is clearly of fundamental importance to the welfare of the community. The Royal Commission on Population, after sifting evidence from many sources, expressed itself as in no doubt of the need for a positive family-planning policy. The National Health Service Act makes provision for the care of pregnant and parturient women but does not lay down any plan of action in respect of family planning, though it may be argued that this should come under the heading of general medical services.

Family planning can be considered under three main headings: (1) the need for facilities for contraception; (2) marriage guidance; and (3) the treatment of involuntary sterility. As regards marriage guidance and the treatment of infertility, there is general and indeed increasing support from all sections of the community, irrespective of religious beliefs. But on contraception opinion is still sharply divided, and a certain number of persons, including all Roman Catholics, regard the use of appliances and withdrawal as morally unjustifiable in any circumstances. They object to public moneys being spent on any clinics where contraceptive teaching is given, or on contraceptive appliances. The use of the "safe period" is, however, accepted by the Roman Catholic Church, and, where spacing is indicated for health or social reasons for such patients, this method may be advised.

Inquiry into the Position

Towards the end of 1949 the Family Planning Committee of the Medical Women's Federation instituted inquiries to ascertain the position of family planning in respect of its three aspects—contraception, marriage guidance, and the treatment of involuntary sterility—as it stood after more than one year's working of the National Health Service Act.

The medical officers of health of 197 local authorities were circularized and were asked the following questions:

*A memorandum prepared by the Family Planning Committee of the Medical Women's Federation: Miss Beatrice Turner, F.R.C.S., F.R.C.O.G. (chairman), Dr. Annis Gillie, Dr. Olive Gimson, Dr. Margaret Hadley Jackson, Dr. Isabelle Little, Dr. Mary Macaulay, Dr. Joan H. Buchanan (honorary secretary), with Dr. Doris M. Odum and Miss Josephine Barnes, D.M., F.R.C.S., as *ex-officio* members.

*British Medical Journal, 1947, 1, 720.

†From the Central Public Health Laboratory, Colindale Avenue, London, N.W.9 (Telephone: Colindale 6041).