REFRESHER COURSE FOR GENERAL PRACTITIONERS

TREATMENT OF BURNS—II

BY

J. P. BULL, M.D. AND D. M. JACKSON, M.D., F.R.C.S.

(From the M.R.C. Burns Unit, Birmingham Accident Hospital)

A further important step in the prevention of infection in burns is early grafting. Only when a wound has healed is the risk of infection finally over. For this reason, as well as to minimize scarring, skin grafts are regarded as the best dressing for a granulating area and should be used as early as possible.

Depth of Burning

The modern classification of the depth of burning is into two degrees—partial-thickness and full-thickness skin-loss. Every recent burn which is diagnosed as full-thickness skin-loss should be considered for excision down to normal tissue followed by immediate grafting. Sometimes this is contraindicated by shock, or by difficulty in diagnosing full-thickness skin-loss or in defining its limits, but excision remains the ideal treatment. If, however, this treatment is not possible, the area of complete skin destruction can always be diagnosed by the end of three weeks, and usually by 16 days after injury, and grafting should not be delayed beyond this time. Every full-thickness burn should therefore be grafted either at once or at the latest within three weeks of injury.

The diagnosis of the depth of a burn is notoriously difficult. Whether partial- or full-thickness skin-loss is present depends on whether the bases of the hair follicles and sweat ducts are destroyed, and this cannot be discovered by inspection of the surface of the skin. A white avascular coagulated surface is no indication of a full-thickness skin-loss burn; in fact, on the face, scalp, palms, and soles it often heals as partial skin-loss within a fortnight. Similarly, within 24 hours of the injury a visible circulation in the burned skin, as judged by redness which blanches on pressure, is no certain indication of partial-thickness loss; for circulation, initially present, may cease several hours after burning, and the lesion may then prove to be full-thickness skin-loss.

The pin-prick test for analgesia is a useful and quick guide to the depth of burning. If part of a burn is pricked repeatedly and firmly with a sterile hypodermic needle, and some pain sensation is present, that part of the burn will heal as partial-thickness skin-loss within three weeks. If analgesia is present (and the burn is not produced by a chemical such as carbolic acid), it is either full-thickness loss or very deep partial skin-loss, taking over three weeks to heal. In certain sites where the skin is thin, such as the back of the hand and fingers and the dorsum of the foot, this test is often enough to indicate primary excision and grafting; but on the face, scalp, palms, and soles the epithelial elements extend more deeply and analgesia is by no means sufficient evidence for primary excision. Although this test is easy to perform, it must be interpreted carefully in conjunction with the history and the site of burning if the excision of partial-thickness skin-loss burns is to be avoided. Generally speaking any area of full-thickness skin-loss more than 1 inch (2.5 cm.) across at its narrowest diameter will take three weeks or more to heal, and should be grafted at once. On the face and fingers even smaller areas may best be treated by grafting.

Grafting of Full-thickness Burns

The most useful type of graft for resurfacing burns is the split-skin graft cut with a Blair knife. The superficial quarter or half of the thickness of the skin is taken in this

way (see Diagram) and a donor area is left which heals in 10 days and can be used to supply further grafts in 16 to 20 days. This is the type of graft used routinely on excised and granulating areas and where the texture of the skin is unimportant: in areas where thicker skin of uniform texture is required, and where avoidance of graft contraction is important, a three-quarter-thickness dermatome graft, a full-thickness graft, or a skin flap should be used. One of the advantages of primary excision is that a full-thickness graft, if indicated, can be applied initially rather than at a second operation.

Diagram of the skin showing the position of epithelial elements. A=Epidermis. B=Dermis. C=Subcutaneous tissue. D=The plane dividing partial-thickness from full-thickness skin-loss. E=Hair follicle. F=Sweat gland and duct. 1=Thin or intermediate split-skin graft (usually cut with a Blair knife). 2=Three-quarter-thickness split-skin graft (usually cut with a dermatome). 3=Full-thickness skin graft.

Extensive Burns

The management of extensive full-thickness skin-loss burns raises further problems. Frequently the first of these is how to keep the patient alive until grafting is possible two and a half weeks after injury. Blood transfusion and artificial feeding are important aids to this end.

The haemoglobin percentage should never be allowed to drop below 80%, and between 120% and 130% is the figure at which to aim when estimating the amount of blood to be transfused. Recently a patient with a 64% burn who survived required 20 litres of blood between injury and complete healing: this is an unusual amount, but it illustrates the large transfusions sometimes needed. When repeated transfusions are likely it is important that blood should be of the patient's own group and Rh factor, and the greatest economy in veins is called for. Haemoglobin estimations should be carried out every two or three days in these cases, as a fall of 5–10% daily may occur.

After the stage of shock anorexia is common and difficulty is often experienced in getting the extensively burned patient to take food. If enough protein can be given to patients in this condition, the appetite is usually restored and they begin to take ordinary meals once more. Usually, however, an ordinary full diet is inadequate to prevent gross

wasting due to the utilization of endogenous protein. A high protein and high calorie diet is required, and this may conveniently be given as a continuous drip feed through an indwelling nasal tube. A useful mixture is: 18 oz. (511 ml.) milk, two eggs, $1\frac{1}{2}$ oz. (42 g.) milk protein, and 1 oz. (28 g.) sugar. This provides 780 calories to the pint (568 ml.), and the usual supplementary feed is 3 to 4 pints (about 2 litres) a day. This type of supplementary feeding should probably be given from the second day to all burned patients who have sustained more than about 20% of full-thickness skin-loss.

A further problem is how to secure skin cover as soon as extensive areas of complete skin destruction begin to granulate. Skin grafting in these cases is an urgent matter, and many patients have died because the danger of delay was not realized by the surgeon. A few of the principles of grafting extensive areas may be mentioned.

1. The maximum amount of skin cover should be obtained at the first operation; the importance of this is shown by the dramatic improvement which often follows during the 24 hours after a large burn is covered.

2. Great care must be taken in preparing patients for extensive grafting operations. The haemoglobin must be at least 80%, and a blood transfusion must be running before an area of 5% or more of full-thickness skin-loss is prepared and grafted. In suitable cases tourniquets, elevation of the limbs, and tilting the operating table can be used to limit blood loss.

3. If the whole burn cannot be grafted at once, the critical areas must be done first; they are the face and neck, hands and feet, and the flexor surfaces of joints.

4. The grafting programme should be planned so that the best use can be made of the donor areas. Where it is avoidable the patient should not be turned over while under the anaesthetic. Even with careful planning it is easy for the surgeon to become unpunctual in carrying out his own grafting time-table.

5. An early decision should be made on the suitability of a case for the use of skin grafts from another person (homografts). If homografts are indicated they should be used early in treatment to gain the greatest benefit from skin cover. When applied in sheets homografts take like autografts, but disintegrate about three weeks later after an immunity has developed against them, leaving the area granulating once more. By using $\frac{1}{4}$ -in. (1.3-cm.) wide strips of homograft, and alternating these with similar strips of autograft, the homografts appear to take permanently. It is probable, however, that the epithelial part of the homograft is replaced by autograft epithelium before immunity has developed.

6. All unused skin graft should be stored in a refrigerator: it will then remain viable up to three weeks, and probably up to six weeks. Small areas of graft failure can be dressed with this stored skin and further operation so avoided.

Special Features of Burns of Certain Sites

Hands.—In most full-thickness skin-loss burns of the hand in civilian practice only the skin and subcutaneous tissue are destroyed; in these cases early full function can be obtained, even if the whole hand is granulating two and a half weeks after the accident. The essential features of treatment to obtain this result are elevation of the limb from the first day to control oedema, immobilization of the hand and wrist in the position of function until epithelization is complete, and the completion of skin cover within three to four weeks of the injury. Sheets of split-skin graft are satisfactory for the back of the fingers and hand, but on the palmar surface a three-quarter-thickness dermatome graft after primary excision gives a better result. Following flexor surface grafts of the fingers splinting in extension is advisable for two to three months, and this can be maintained conveniently without complete immobilization by using a leather glove with stay bones incorporated over the backs of the fingers, hand, and wrist. Deep lesions involving tendons, nerves, vessels, and bones are usually the result of burns during unconsciousness, and burns by electricity or by the hand being firmly held between hot rollers or in faradic spasm on an electric fire bar. Immediate excision of these deep burns is often unwise, since it is exceedingly difficult to define the limits of necrosis in the first few days. Nevertheless, excision of non-viable

tissue is still indicated as soon as demarcation is present, and this is essential if healing is to be complete within the critical three to four weeks. Surgery in these cases has the same objective as in other types of hand injury—to restore the action of the hand as a pincer and a vice before it can become stiff and useless.

Face.—It is generally recognized that full-thickness skin-loss burns of the eyelids should be grafted as soon as possible to prevent ectropion, which may possibly lead to corneal ulceration and blindness. It is important, however, that the whole face and neck should be treated as a critical area and grafted not later than three weeks. Areas of complete skin destruction on the head and neck often result from falling into the fire in an epileptic fit. These burns can usually be excised and replaced with a dermatome graft on the day of injury, and the cosmetic result may be so good that further treatment is unnecessary.

Feet.—It is not generally appreciated that split-skin grafts cut with a Blair knife give excellent results in deep burns of the feet. Even on the sole and over the tendo Achillis they stand up well to rough wear, and they have proved satisfactory as the routine graft for molten-metal burns in foundry workers, who stand for long hours in heavy boots in a hot, dirty environment. They are contraindicated only if the burn has destroyed the pad of subcutaneous tissue or paratenon, and bone, joint, tendon, or nerves are exposed. After grafts on the feet have taken, controlled return to walking is necessary to prevent damage by capillary haemorrhages beneath the graft. This danger is avoided by dressing the graft on the fifth day after operation, and, if it has taken satisfactorily, giving active exercises in bed for two days, hanging the legs over the edge of the bed for one day, sitting out for one day, and walking on the tenth day.

Bone.—A free graft will take satisfactorily on periosteum, but, if the periosteum is destroyed by burning, the exposed bone should be regarded as dead. Burns down to and including bone, which occur most commonly on the skull, should be excised down to viable tissue and skin grafted to secure early healing. A split-skin graft on marrow is satisfactory for wound closure and sometimes as permanent skin cover for the skull, but it is not satisfactory where it will be subject to pressure or rubbing.

Joints.—In applying free grafts near to a joint there are several important principles which should be followed. The joint should be extended to make the burn on the flexor surface as large as possible before grafting. Scars between the graft and sound skin, and between separate sheets of graft, should be planned to run round the limb and not across the flexor crease, where they are likely to form tension bands. Splinting is never required for grafts on extensor surfaces if they are stretched from the start by active exercises. Often, however, grafts on flexor surfaces benefit from splinting combined with active exercises for periods up to three months. By this time the tendency for the graft to contract has usually ceased unless there is gross shortage of skin, when further grafting will be required.

Respiratory Passages.—It is unusual for the respiratory passages to be burned even when the face is severely burned by flames, presumably because the breath is held. A conflagration in an enclosed space is the commonest cause of this lesion; more rarely it may follow the swallowing of hot liquids. If the lungs are involved a large amount of exudate is rapidly poured out and the patient tends to drown. The prognosis is very grave and care should be taken not to give excessive amounts of intravenous fluid. Early tracheotomy, with administration of oxygen and frequent suction to keep the air passages clear, offers the best hope. Tracheotomy is also indicated for oedema of the glottis following the drinking of hot liquids if obstruction is severe. It is contraindicated for the noisy breathing and gross oedema of the subcutaneous tissue which accompanies burns of the neck; in these cases, if the patient sits erect in bed the oedema quickly descends on to the trunk.

Chemical and Electric Burns

Chemical Burns.—It is often impossible to tell at once whether a chemical burn from an unknown caustic is due to acid or alkali, so that valuable time may be wasted in searching for an appropriate antidote. Treatment is made easier by a buffer solution which will neutralize both types of caustic. A mixture of monobasic and dibasic phosphate is recommended which has the following composition: monobasic potassium phosphate (KH_2PO_4), 70 g., and dibasic sodium phosphate ($\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$), 180 g., in 850 ml. of water; this makes 1,000 ml. of solution. The solution is non-irritant and can be used in the eyes. The caustic should be washed off immediately and vigorously either with water or with this solution and a compress of the buffer applied for 20 minutes. The chemical burn should then be treated as described for a thermal injury.

Electric Contact Burns.—These are usually painless and deep. Until the blistered skin is removed they tend to look unimportant, but because of their depth the slough is slow to separate and the scarring great. They can often be treated by primary excision and grafting, but it is sometimes difficult to decide how much tissue is devitalized. Electric contact burns on the hands may coagulate the tissues down to tendon, and in these circumstances a skin flap is indicated to preserve finger movement.

Conclusion

The modern management of burns owes much to new knowledge of physiology, bacteriology, and pathology. This has been combined with the general surgical principles of the treatment of open wounds and the special skills of plastic surgery. As a result many burned patients can now be saved whose injuries until recently would have been thought certainly fatal. For instance, instead of children with burns of one-third of the body surface probably dying, four-fifths of such patients should now survive. It is evident that further advances will be made. Much effort is now being applied to the prevention of burns and scalds, particularly of the all too frequent domestic injuries. The fuller understanding and control of the shock period, better prevention of infection, and technical improvements in grafting are all immediate possibilities. Assessment of the value of changes in detail of therapy is not easy and usually demands study of carefully controlled series. Comparison of mortality results has now been simplified so that series treated under different regimes can be validly compared from this point of view. The other two important criteria of success—speed of healing and completeness of function—should also be available when judging the value of new methods.

Next Refresher Course Article.—"Diseases of the Gall-bladder," by Mr. Harold C. Edwards.

The Minister of National Insurance has asked the National Insurance Advisory Committee to review, in the light of changes in the cost of living and other relevant circumstances, the amounts by which the National Insurance benefits of long-term hospital in-patients are reduced. Benefit is at present reduced after eight weeks in hospital by 5s. a week if the patient has a dependant, and by 10s. a week in other cases. After a year in hospital benefit is further adjusted, leaving the patient with 10s. a week if the treatment is for respiratory tuberculosis and with 5s. a week in other cases. But if the patient has a dependant the balance may be paid to the dependant. The committee will consider representations from interested organizations or persons. Representations, in writing, should be sent to the secretary, National Insurance Advisory Committee, 30, Euston Square, London, N.W.1, before June 20.

THE NEW MINISTER OF HEALTH

On May 7 it was announced from 10, Downing Street, that the Minister of Health, Captain H. F. C. Crookshank, had tendered his resignation, and that the Queen had been pleased to approve the appointment of Mr. I. N. Macleod as Minister of Health. Captain Crookshank succeeds Lord Salisbury as Lord Privy Seal and will continue his duties as Leader of the House of Commons.

The Rt. Hon. IAIN NORMAN MACLEOD, M.P., P.C., is 38 and the eldest son of a doctor, the late Dr. Norman Macleod, who for many years was in practice at Skipton, in Yorkshire. He was educated at Fettes and at Gonville and Caius College, Cambridge, where he took an honours degree in history.

In the 1945 General Election Mr. Macleod contested the Western Isles, a constituency in which he had previously lived but where no Conservative candidate had come forward since 1931; he was unsuccessful in a three-cornered fight. He returned to the Army and served for a while in Norway, before being demobilized in 1946.

[Luther-Smith, Enfield

After the war Mr. Macleod took to politics in good earnest. He joined the Conservative Parliamentary Secretariat, and worked under Mr. R. A. Butler. In 1948, when the Party's Home Affairs Research Department was established, he became its joint head with Brigadier Enoch Powell, and when the latter left he continued in sole charge until just before the 1950 General Election. This department provides background information for Conservative Members of Parliament, and also undertakes the detailed examination of policy. In 1946 Mr. Macleod was adopted as the prospective Conservative candidate for Enfield, Middlesex, and, after the division of the constituency under the Representation of the People Act, 1948, for Enfield West. He won this seat for the Conservatives in 1950 with a majority of over 9,000 in a three-cornered contest, a figure he improved upon in the 1951 election in a straight fight. After serving as secretary of his party's Parliamentary Committee on Insurance and Pensions in the 1950 Parliament, in the next Parliament Mr. Macleod was made chairman of the Conservative Parliamentary Committee on Health and Social Security in succession to Sir Hugh Lucas-Tooth. He was also chairman of a group of nine newly elected Conservative M.P.s who in 1950 published their views on home policy under the title *One Nation*. They dealt with such questions as the social effects of redistribution of wealth, population policy, housing, education, health, insurance and assistance, the care of old people, social policies for industry and the countryside, local government, and payment for the social services. Mr. Butler lent his weight to the book with a commendatory foreword. Mr. Macleod is joint author (with Brigadier Powell) of another political pamphlet, *The Social Services: Needs and Means*, an essay originating during the passage of the National Health Service and National Insurance Bills of 1951, when it appeared to the authors that neither major party had clearly thought out the relationship between the social services and the means test. This pamphlet was the subject of a recent leading article in the *Journal* (January 26, p. 207).

Since being appointed Minister of Health Mr. Macleod has been made a Privy Councillor. He does not have a seat in the Cabinet.