REFRESHER COURSE FOR GENERAL PRACTITIONERS

HEART BLOCK

BY

GEOFFREY BOURNE, M.D., F.R.C.P.

Physician and Officer-in-Charge, Cardiological Department, St. Bartholomew's Hospital

Heart block can be defined as a condition in which the impulse from auricles to ventricles is partially or completely interrupted. This interruption may involve the main stem in the bundle of His, in which case, although the impulse is from time to time slowed or permanently arrested, the method of spread in the ventricular muscle beyond this obstruction is normal in type. Bundle-branch block is a condition in which partial or complete interference occurs in the transmission of the impulse down either the right or the left branch of the bundle. Bundle-branch block may occur in cases of ordinary heart block.

Lesions of the Main Bundle

The bundle of His may be affected by some pathological process in such a manner that the transmission of the impulse from auricles to ventricles is slowed down but not obstructed. This condition is known as latent heart block, for no ventricular beats are dropped out, and there is no clinical evidence of the presence of the block. In the next stage beats are dropped from time to time when the transmitting power of the bundle begins to tire. After a dropped beat, however, the bundle is rested so that the transmitting time picks up again. This is called the stage of dropped beats (Fig. 1). These two varieties of heart block are slight in degree; they are usually found in acute conditions such as diphtheria or acute rheumatism. The pause of a dropped

FIG. 1.—Heart block with dropped beats. Leads I, II, III, and CR4. In Lead I the P-R interval of the first two beats is 0.28 and 0.36 second respectively. The third auricular impulse fails to traverse the bundle of His, and no ventricular response is evoked. The ST changes of a recent posterior infarct are also present.

FIG. 2.—Complete heart block. Leads I, II, and III. In Lead II it is clear that the regular P waves have a rhythm which is quite independent of the QRS or ventricular complexes.

beat is silent: in the pause due to a premature beat the extrasystole can always be heard through the stethoscope, if it is carefully listened for. In low-grade heart block the loss of function of the bundle is an indication of the general effect of the toxin or the disease upon the state of the heart muscle as a whole.

More severe grades of heart block are as a rule found in more chronic pathological states, the lesion affecting the bundle being often a local one, generally arteriosclerotic but occasionally syphilitic. The usual arterial supply to the bundle of His is a separate branch of the right coronary artery, so that heart block is an infrequent complication of coronary thrombosis, which generally involves this vessel lower down. Congenital varieties of heart block are not uncommon, constituting perhaps 10% of all cases of high-grade block. Heart block usually occurs in auricular flutter, where the auricular rate is usually 200 beats a minute or more. This high auricular rate over-stimulates the bundle of His and fatigues it. The heart block in this condition does not indicate disease of the conducting tissues.

The more severe grades of heart block usually involve a delay in the bundle sufficient to stop many of the impulses from auricle to ventricle. The usual varieties found are 2:1, 3:1, or complete block. In 2:1 block every other auricular impulse fails to reach the ventricle. In 3:1 block two out of three fail to do so, and in complete block no auricular impulses are transmitted to the ventricle. In the latter condition ventricular activity is evoked by a separate pacemaker situated in the bundle of His above its division into two branches. From this position, usually at a regular rate of from 30 to 40 beats a minute, impulses arise and stimulate the ventricles to contract (Fig. 2).

Effect of Heart Block on Cardiac Function

Partial heart block in which few of the beats fail to reach the ventricle from the auricle has no significant effect upon the cardiac function. The heart muscle, however, is considerably weakened by the general myocarditis of which this slight degree of block is a sign. Clinically, therefore, the presence of this slight block is valuable as an indication of myocardial weakness. These types of block are as a rule recovered from completely if the patient survives the acute disease. This recovery in the case of acute rheumatic carditis may be unexpectedly good. I have seen three patients, all of them adolescents or adults, who suffered from acute rheumatic carditis so severe as to cause pericarditis, high fever, and an unusually high grade of heart block. In spite of the severity of the disease and its widespread effect on the heart, all of these patients made a perfect recovery and were able to return to normal physical activities in course of time. They remained well for a number of years before they were lost sight of.

High-grade heart block presents a different type of physiological problem. The best example is complete block in which the ventricles are beating at a speed of 30 to 40 beats a minute. The bradycardia of necessity causes a long diastole. This has a number of important results: the ventricles are well filled; a big stroke output follows; the ventricles are stretched and a little enlarged; the systolic pressure is raised; the diastolic pressure is lowered; the coronary flow is prolonged. There are three clinical points, therefore, which must be remembered in heart block with bradycardia in order that the patient's true physical condition shall not be misunderstood. Some enlargement of the

ventricles is physiological in heart block, therefore part of the cardiac enlargement can be disregarded. The raised systolic blood pressure in such cases does not mean hypertension; a systolic reading of 180 to 190 is physiological. Finally, when this systolic pressure is associated with a diastolic reading of 70 to 80 this very big pulse pressure is also a physiological result of the bradycardia, and is not due to aortic regurgitation or to any similar pathological process.

A further result of heart block, which is not physiological in origin but due to a direct extension of the causative disease to the neighbourhood of the idioventricular pacemaker in the bundle of His, is seen in the form of Stokes-Adams attacks. When disease affects the activity of the ventricular pacemaker its function becomes unreliable and in some cases it ceases work completely, bringing the ventricles to a standstill. When the standstill lasts for four seconds or more the result is transient giddiness. If the period is prolonged further a brief spell of unconsciousness occurs. Should the time be longer still the unconscious patient suffers from generalized convulsions. During the Stokes-Adams attack there is no pulse at the wrist and no evidence of cardiac activity on listening with the stethoscope. Pallor followed by cyanosis occurs during the period of ventricular standstill, and also deep breathing. At the end of the attack, as the patient comes round, the body flushes red and very often breathing stops for a space, since the previous hyperpnoea has washed out most of the carbon dioxide from the blood stream. The chief diagnostic points suggesting a Stokes-Adams attack are: complete cardiac standstill; an abrupt onset; a duration averaging five to twenty seconds and not exceeding thirty seconds; and rapid and complete recovery to a normal state. Epilepsy, true or arteriosclerotic, vasovagal attacks, and the rare syncopal attacks of auricular flutter and aortic stenosis must be excluded.

Prognosis

The outlook in these high grades of heart block due to chronic myocardial disease is generally not too good, the average expectation of life being about five years. In trying to assess the prognosis, due attention must be paid to other factors such as a history of coronary disease, much enlargement of the heart, nocturnal paroxysmal dyspnoea, and so on.

Treatment

The treatment of the minor degrees of block which are seen in acute infections is that of the acute infection. Complete block and high-grade block need no treatment as such, but if heart failure is present the patient should be treated by rest in bed, mercurial diuretics, intravenous or intramuscular aminophylline, and by digitalis. The bradycardia of heart block is no contraindication to the use of digitalis if congestive failure is present. Digitalis acts by depressing the conductivity of the bundle of His. If this conductivity is already so depressed as to have become complete it is quite clear that the action of the digitalis has been forestalled, and that the drug cannot have any more effect upon the already slowed heart rate. In other words, digitalis cannot make more complete what is already complete.

Stokes-Adams attacks are usually so brief that treatment cannot be administered during them. Therapy must therefore be concentrated upon prevention. In most cases ephedrine hydrochloride, 1 gr. (65 mg.) by mouth three times a day after meals, is efficacious in preventing the attacks. In some cases the dosage may be less than this and in others it needs to be more. If ephedrine does not act perfectly its effect can sometimes be enhanced by atropine. This also can be given by mouth at the same time as the ephedrine, and the dosage again will vary from case to case: sometimes 1/200 gr. (0.32 mg.) three times a day will suffice, but in others the adequate dose will be found to be 1/100 gr. (0.65 mg.) or even 1/75 gr. (0.87 mg.). Should ephedrine be found inadequate adrenaline may be given two or three times daily by injection subcutaneously or intramuscularly. The dose should be 0.5 to 1 ml. Sublingual isoprenaline

may also be tried. Quinidine inhibits the activity of the idioventricular pacemaker, and its use in high-grade heart block is therefore very dangerous.

Bundle-branch Block

The myocardial lesion which involves one or other of the branches of His, and which leaves the main bundle unaffected, may occur in either acute or chronic conditions. Diphtheria, myocarditis, and acute rheumatism are again the most outstanding acute infective causes. The abnormality is only part of the general myocardial disease and its clinical importance is subsidiary.

Chronic bundle-branch defects may roughly be placed in one or other of two categories. In the first group the lesion may be one of several or of many other evidences of organic cardiovascular trouble. It may coincide with chronic coronary disease, aortic valve disease, or hypertension, and may be associated with symptoms of myocardial failure. Here it is merely one of many proofs of organic heart disease, and it is insufficient as such to add to the seriousness of the general situation. It can also be a manifestation of a cardiac infarct, appearing during the acute stage and disappearing as the patient recovers.

The second category is that in which a bundle-branch lesion, right or left, seems to be an isolated phenomenon, or at any rate the most outstanding abnormality present. In such patients the heart may not be enlarged or its enlargement may be slight and the function of the heart muscle may be good. One such patient when first examined 16 years ago had a normal electrocardiogram. At his next examination two years later a bundle-branch lesion was discovered. He has since remained in good health in spite of his cardiographic abnormality and is now aged 63. The electrocardiogram of a right or a left bundle-branch lesion is a sufficiently obvious deviation from the normal to constitute a picture in black and white which of itself might be frightening enough to give rise to unduly careful treatment or an unnecessarily bad prognosis (see Figs. 3 and 4). It is therefore important to realize that when a bundle-branch defect is the only major abnormality present, undue attention should not be focused upon it. The important points in

FIG. 3.—*Left bundle-branch block.* Leads I, II, III, and CR4. Note the slurring and the great prolongation of the QRS complexes in all leads. This is the constant characteristic of bundle-branch block.

FIG. 4.—*Right bundle-branch block.* Leads I, II, III, and CR4. The QRS is prolonged to 0.16 second. The S wave is deep and slurred in Lead I.

such cases are the heart size and the capacity of the patient to undergo exercise. If the heart is normal in size and if the patient suffers no shortness of breath or other significant symptoms, the bundle-branch lesion may be almost disregarded.

A bundle-branch lesion in other patients may occasionally be part of a latent but more extensive damage to the bundle of His. Such cases may be examined fairly often and yet be found to have a heart rate which is normal and without dropped beats; but in spite of this they are subject to sudden short fainting attacks. These are due to the fact that the idioventricular pacemaker is involved in the pathological process, so that from time to time a more extensive heart block intervenes with cessation of the idioventricular pacemaker and the onset of Stokes-Adams attacks. The important diagnostic points here are the presence of the bundle-branch lesion and the fact that the Stokes-Adams attacks are typical.

Next Refresher Course Article.—"Treatment of Burns," by Dr. J. P. Bull and Mr. D. M. Jackson.

A CANCER RESEARCH JUBILEE

Fifty years ago—in April, 1902—a public appeal was made for the establishment of a fund to be devoted to the investigation of the cause, nature, and treatment of cancer. Some £30,000 was quickly subscribed, and in July of that year the Royal Colleges of Physicians and Surgeons consented to take control of the investigation by setting up a general committee to formulate a scheme of research. In a supplement to the annual report of the Imperial Cancer Research Fund, presented at the recent 49th annual meeting, the work of the ensuing 50 years is described.

The first chairman of the fund was Sir William Church, who at the time was President of the Royal College of Physicians. He remained in the chair for 22 years. His successor, Dr. Sydney Martin, died in the year of his appointment, and was followed by Sir Humphry Rolleston, who served for 15 years, and he by the present chairman, Professor H. R. Dean. Sir Henry Morris, who was the leading spirit in the original discussions, was the first treasurer, to be succeeded by Sir William Watson Cheyne, Sir George Makins, and the present holder of the office, Sir Holburt Waring. The fund has had four directors. Dr. E. F. Bashford was the first, and it was largely due to his initiative that the first scientific reports attained such importance. He was followed in 1914 by Dr. J. A. Murray, who served until 1935; and he by Dr. W. E. Gye, now pursuing research in the field of cancer in Australia, and Dr. James Craigie, the present director. The earlier laboratories were in rooms allocated to the fund by the Royal Colleges in their examination halls, but in 1936 a property at Mill Hill was secured.

The story of the research carried out under the auspices of the Imperial Cancer Research Fund runs parallel with that in other laboratories. Much useful inquiry has been initiated by the Fund itself, but more perhaps has been done by taking up work reported from other sources, especially Continental and American, and assessing its value and carrying it further. When the Fund started operations cancer statistics were in a chaotic state, and one of the first pieces of work was to discover whether the alarming rate at which cancer was reported to be increasing at the beginning of this century was a true increase or whether it was due, thanks to greater attention to clinical recognition and improved diagnostic methods, to the discovery of hitherto "missed" or "inaccessible" cancers. It was found that most of the increase was referable to "inaccessible" sites, the stomach particularly. Another question tackled was whether cancer was confined to civilized races, and as a result of examination of a large number of specimens sent to the laboratories it was declared that cancer occurred among all races and in all climates throughout the Empire.

Again and again in this short history one is reminded of landmarks in cancer research: the work of Jensen on tumour transplantation, which was followed up in these laboratories; Warburg's discovery of aerobic glycolysis in cancer cells, which stimulated much new work on metabolism, although later some of his generalizations proved untenable; the announcement by Rous that a sarcoma of the domestic fowl could be communicated by means of cell-free preparations obtained by mincing and grinding the tumour; Shope's discovery of a tumour virus in the cotton-tail rabbit; Bittner's work on the transmission of mouse cancer by the mother's milk. Then there has been a vast amount of work done on cancers caused by exposure to irritating substances such as tar, and in the wide field of carcinogenesis in general. All this and much more has engaged the laboratories of the Fund, and notable contributions have been made by the scientific staff.

Perhaps, looking back on this history, a warning given by Dr. J. A. Murray, then director, as long ago as 1931 may still be appropriate. He begged the lay and the general medical public to appreciate the conservative and cautious attitude which those engaged in cancer research preserve towards the recurring extravagant announcements of new methods of treatment of cancer. The idea that it is all a gamble, and that success is equally likely to fall to the ignorant as to the instructed, should be discarded. To quote Murray: "Whatever the future of the cancer problem may be, it is certain that it will not yield to the sound of the trumpet."

Reports of Societies

SPREAD OF INFECTION IN THE FAMILY

At a meeting of the Section of Epidemiology and State Medicine of the Royal Society of Medicine on April 18 there was a discussion on "The Spread of Infection in Families." Introducing the speakers, the president, Professor ROBERT CRICKSHANK, pointed out the special interest of the family as the smallest epidemiological unit and the opportunities it presented for valuable observations by the general practitioner and the paediatrician. Such observations had been made in the past by Snow and Budd, and recently by Pickles, of Aysgarth.

Dr. W. C. COCKBURN, outlining the essential facts to be obtained in the investigation of infectious disease, said that, so far as the family was concerned, these were few and simple: the date of onset of illness and the age, sex, position in the family for the first case, and the number and type and date of onset of symptoms in the household contacts. For most investigations bacteriological co-operation was necessary; he instanced the American observations on the relative frequency of poliomyelitis virus in the excreta of the household and of less close contacts. Epidemiological information of this sort could be used for the following purposes: (1) The study of the "spreading power" of the disease: something was already known of the similarity of salmonella and shigella infections in this respect. (2) The mode of spread—for example, did this differ when the first case was in a child, an adolescent, or an adult? (3) The effect of age on susceptibility. (4) The relation of numbers of clinical cases to persons harbouring the organism without symptoms: recent analyses of food-poisoning infections in this country, carried out by the Public Health Laboratory Service, had shown that in family outbreaks there was a surprisingly high proportion of symptomless excretors. Dr. Cockburn believed that the difficulty of knowing the precise time of infection in a family made family studies less valuable for determining incubation periods.

Whooping-cough and Measles

Recent work on the evaluation of prophylactic agents had incidentally yielded interesting information about the spread of pertussis within the family. Inquiries connected with