REFRESHER COURSE FOR GENERAL PRACTITIONERS

TREATMENT OF BURNS—I

BY

J. P. BULL, M.D.

AND

D. M. JACKSON, F.R.C.S.

(From the M.R.C. Burns Unit, Birmingham Accident Hospital)

[This article is concerned chiefly with the hospital treatment of burned patients, including skin grafting. Although of prime interest to residents and registrars, we believe that many general practitioners will welcome this account of current methods of treatment. Some will be glad of it for more immediately practical reasons, as they may have charge of such patients in cottage hospitals or privately.]

Since long before man discovered the use of fire he has suffered injuries from burning. Four thousand years ago the ancient Egyptians recommended oil-soaked cloth as a dressing. In this country we continue to have open domestic fires, and two-thirds of serious burns and scalds occur in the home. The remaining one-third of civilian cases come from industry, where high temperatures, inflammable fluids, chemicals, and electricity are common hazards. Modern warfare produces injuries from flame throwers and incendiary and phosphorus bombs; added to these are the more recent problems of possible mass casualties caused by atomic weapons.

Whatever the source of heat, the effects on the patient are largely determined by whether clothing catches fire. Nightclothes and dresses of cotton and rayon are most commonly responsible in fatal domestic burns (just as oil- or petrol-soaked overalls can account for a similar fatal result in the factory or garage worker). As Leonard and Vera Colebrook have pointed out, adequate guards for open fires, whether coal, electric, or gas, are an obvious precaution.

The treatment of burns has long been a Cinderella of surgery. The apparent simplicity of the injury has led to its dangers being underrated. Too often treatment has been relegated to junior staff, and cases have seldom been taught upon in the teaching hospitals. This situation has improved recently, and with new knowledge of resuscitation, antibiotics, nutrition, and surgical techniques results are better. It is clear that the ideal treatment has still to be found, but a review of the aims of therapy and current methods will help the clinician to decide on the appropriate management of different types of burns in the varying circumstances of home, surgery, and hospital.

The Local Lesion and its General Effects

Different forms of trauma produce essentially the same burn lesion. Burns and scalds due to hot water, steam, burning clothing, molten metal, chemicals, and electricity all cause damage to the epidermis and varying degrees of damage to the dermis and perhaps to deeper structures. A temperature of 55° C. for 30 seconds produces the superficial erythematous lesion familiar in the minor domestic scald. A temperature of 60° C. for 30 seconds causes blistering and death of part of the Malpighian layer. Such "partial-thickness skin-loss" heals spontaneously in one to three weeks by regeneration of the remnants of this layer: the cells lining the hair follicles are particularly active in growing out to cover the denuded surfaces. A temperature of 65° C. or more for 30 seconds often causes complete destruction of epidermal elements. Sometimes a few cells of deeply penetrating sweat ducts survive, and these may give the late spontaneous healing of deep partial-

thickness skin-loss at four to six weeks. Otherwise the only cover can come from the edges, and by the time this slow and uncertain procedure can succeed extensive scarring will have occurred.

The injury elicits a prompt reaction in the local blood vessels: there is often a momentary vasoconstriction and the capillaries may be fixed in this condition in the coagulated dermis of high-temperature burns; but in the commoner less severe injuries this stage is immediately followed by extreme vasodilatation and increased permeability. Localization of the escaping fluid between the skin layers causes blistering, and its diffuse spread deep to the epidermis causes oedema. Whether and when blistering occurs depends on local conditions, and its presence is not diagnostic of any particular depth of damage, since both superficial and deep burns may show blistering. The fluid which leaks out into the blister has a high protein content intermediate between lymph and plasma. To correct for this loss more fluid is drawn into the blood stream from uninjured areas, but a large burn will exhaust these reserves and a reduction in circulating blood volume results.

An approximate index of this change is provided by the haematocrit, which measures the proportion of red cells to plasma; the haematocrit reading rises as plasma is lost from the blood and returns to normal when the loss is corrected by therapy. At the end of this period of high permeability many of the local capillaries remain dilated and are filled with static red cells. The red cells thus lost into the burn contribute to an anaemia which is probably accentuated by haemolysis of red cells that have passed through the burn and by subsequent delay in normal replacement. Though there is thus a loss of red cells, particularly in large burns, plasma loss predominates in the first 24 hours after the injury. Should replacement therapy be delayed or insufficient, circulatory collapse similar to that seen in traumatic shock results. If this continues the vasomotor control ultimately fails and no amount of added fluid will restore an effective circulation. We are still ignorant of the full mechanism of this irreversible shock.

Apart from the reduction of circulating volume, upon which all authorities are agreed, the other general effects of burns are as yet less clearly defined. Toxins of several types have been said to be released by the burned tissues: "sludging" of blood probably occurs, though its clinical significance is uncertain; the stress response is evoked and the implications of this are being studied. From the purely clinical point of view these fields of investigation are not yet sufficiently developed to give much help in the management of patients.

The loss of areas of skin has two important consequences. The normal barrier to surface infection is replaced by moist dead tissue, an excellent medium for bacterial growth. Similarly the normal surface barrier to fluid and protein loss is destroyed, causing increased nutritional demands. These two effects are increased with increase of area, and should infection be allowed to occur the fluid and protein loss is further increased and prolonged. These complications alone can cause wasting and anaemia without there being need to invoke any hypothetical toxin.

Aims of Treatment

The aims of treatment follow from our knowledge of the lesion and its course. The fluid loss continues for several

hours, and in large burns for a day or more. Maximum haemoconcentration is not immediate, and there is thus time to replace the loss and then to balance the continuing leak until the danger of circulatory collapse is past. One aim therefore is the prevention of shock by intravenous fluids. Prevention and correction of anaemia by blood transfusion are also required for large burns, and together with generous feeding help to maintain the natural resistance of the patient and favour healing.

The local healing of partial-thickness burns depends on the survival of epithelial cells. Nothing must be done to prejudice their growth; local application of antiseptics is therefore dangerous, as is also the development of sepsis in the burn. Local treatment must protect the burn from further injury and from infection. Where all epithelial structures have been destroyed, satisfactory new skin cover can be obtained only by grafting. So long as such raw areas remain they drain the body's resources and are avenues for the entrance of infection. The further aim of treatment in these cases is therefore to graft new skin as early as possible; this has the added benefit of reducing scar formation.

Control of Shock

If the plasma loss is great the reduced circulating volume will cause clinical shock to develop. The following degrees of shock may be mentioned:

Threatened Shock.—The loss of plasma from burns of only moderate extent can be compensated by the body's resources, so that, in spite of some haemoconcentration and diminished circulating volume, clinical shock does not appear. This state of threatened or latent shock usually passes off with rest, warmth, and oral fluids. If, however, compensation is disturbed by moving the patient, by exposure to cold, or by operation, clinical shock may be precipitated unexpectedly.

Reversible Shock.—The classical picture of pallor, restlessness, tachycardia, oligæmia, and lowered blood pressure will ensue if the compensation for fluid loss becomes inadequate. Transfusion of a colloid solution such as plasma will then be required to restore the circulating volume. If this is not administered in adequate amount and early enough, irreversible shock is likely to follow.

Irreversible Shock.—This type is similar clinically to the previous one, but differs from it in that it proceeds to a fatal outcome in spite of apparently adequate transfusion. It is therefore a post-mortem diagnosis which is necessarily relative to present knowledge, and it must not be made until all therapeutic measures have been tried.

Shock is a condition much more easily prevented than cured. Patients with severe burns often reach hospital within an hour of injury, and before shock is clinically apparent: the opportunity for preventing it is therefore frequently presented. In practice, burns of more than 15% of the surface area in an adult, and more than 10% in a child, will produce clinical shock unless the patient is transfused, so that an estimate of the extent of the burn is the first essential of diagnosis. A convenient method of memorizing the surface area proportions for different parts of the body is shown in the diagram, and an additional aid is that the palmar surface of the patient's outstretched hand is approximately 1% of his body area. By these means it is easy to estimate whether the area burned exceeds the critical figure above which shock may be expected. When a patient has been placed in the

The "rule of nines" simplifies the estimation of surface areas of the body. Each arm and the head and neck are approximately 9%, and each leg and the front and the back of the trunk are each 2 by 9%. The perineum and genitalia bring the total to 100%.

"shock" category the most important thing is to begin transfusion as rapidly as possible. Delay in treating such a case may easily convert it into one of irreversible shock, so that this is one of the few circumstances in medicine where speed is urgently necessary.

The composition of the fluid lost from the circulation is akin to plasma, so that the latter seems to be the obvious replacement fluid to choose, and it is in fact the one commonly used. Enough plasma should be given to replace the loss, and it should then be continued at a rate sufficient to maintain the plasma volume.

Transfusion

There are three practical aids for controlling the rate of transfusion: repeated haematocrit estimations, a surface-area formula, and the use of certain clinical signs. None of these is sufficient alone, but all should be used in conjunction.

The Haematocrit.—The haematocrit reading is a useful guide to plasma loss because it is sensitive and changes early; its main weakness is the difficulty of estimating the normal value for an individual patient before injury. Average haematocrit values, however, are known for the two sexes at various ages, and a numerical measure of the degree of plasma deficiency can thus be obtained. The haematocrit estimation can easily be done in the clinical room attached to the ward, even in artificial light and without much practice. Only 0.5 ml. of blood is required, and capillary blood taken from an area of warm skin is satisfactory. An alternative which may be useful in some circumstances is haemoglobin estimation. Blood-pressure readings for this purpose are useless, since a fall of blood pressure occurs late in shock due to burns.

The Surface-area Formula.—If enough plasma is given to restore the haematocrit reading to normal as soon as possible and then to keep it there, most adults will require an amount proportional to the area burned. This observation can be made the basis of a surface-area formula. A few cases, however, need more than this, sometimes much more, and these cases demonstrate the fallacy of relying entirely on a formula. For adults a total of 1-1½ litres of plasma is required for every 10% of the body surface burned, excluding superficial erythema; but the variation in plasma requirements between individual burned children is so great that we have been unable to find a reliable formula for forecasting their needs. An alternative used in some centres is a formula calculating the total fluid requirement for 48 hours from the patient's age, weight, and surface area burned, and controlling the rate of transfusion by a fixed time-table (see *Ann. Surg.*, 1947, 126, 1010; and *Lancet*, 1951, 1, 501). This formula rests upon the principle that the fluid loss from the circulation is proportional to the area burned, a relationship which is probably only approximately true.

Clinical Signs.—In addition to these two numerical methods, a careful watch must be kept on the patient: there is no place for dictating treatment at occasional visits to the bedside. The three most useful physical signs in the shock stage are thirst, lowered urinary output, and restlessness, all indicating the need for further resuscitation. The hourly urinary output is so valuable in the management of the shocked patient that an indwelling catheter should be used routinely in burns of over 20%. Left in for 24 to 48 hours the risk of urinary infection is small, and is quite outweighed by the advantages. A urinary output below 30 ml. an hour in an adult should be regarded as inadequate. A fall of blood pressure, especially in the second 24 hours, is a very serious prognostic factor and usually indicates the onset of terminal peripheral failure.

Past experience has established that the most satisfactory cases receive half of their plasma requirement in the first six to eight hours, and the surface-area formula enables one to make a rough estimate of what this should be. If a transfusion is required it is usually necessary to continue it for 24 hours, and sometimes up to 36 or 48 hours.

Dried plasma can be made up with normal saline or the usual distilled water. When saline is used the protein and sodium chloride in the transfusion fluid are about twice as concentrated as in the exudate; this is corrected by an equal volume of oral fluids in the form of fruit-drinks sweetened with glucose, 1½ oz. to 1 pint (75 g. to 1 litre). Should this route be impossible owing to vomiting, an equal volume of 5% glucose in water should be given intraven-

ously instead. The fluid lost into the burn takes with it its normal complement of electrolytes, so if no extra saline is given intravenously it is advisable to use 1.75% sodium lactate or 0.3% sodium chloride with 0.15% sodium bicarbonate in oral fluids.

Blood Transfusion.—Although plasma loss from the circulation is the principal cause of the diminished circulatory volume, red-cell destruction and loss also occur as a result of the direct action of heat, haemolysis, stasis, and sometimes haemorrhage from the alimentary tract and kidneys. Burns of sufficient extent to require treatment for shock need blood as well as plasma if anaemia by the end of the first week is to be prevented. The best time for giving this blood is debatable, but in view of the functional loss of blood from the circulation in shock due to pooling, there is some reason for giving it early in the shock treatment. This makes the interpretation of the haematocrit readings more difficult, however, and, provided the patient's condition and progress are satisfactory without extra blood, we frequently postpone transfusion till the end of the shock period.

Plasma Substitutes.—Such substitutes, though perhaps not possessing all the advantages of plasma, should be free of its disadvantages, which include the risk of infective hepatitis. One of these is "dextran," a polysaccharide obtained from *Leuconostoc mesenteroides*, which is now available and is being used increasingly. It is too early to say whether dextran can fully replace plasma, but in our present state of knowledge a reasonable procedure is to use up to one plasma volume of dextran, and then to continue the transfusion, if necessary, with plasma or blood.

Sedation.—Morphine should be given only to those burned patients who are restless, complain of pain, or are hysterical with fear; in these cases a single dose should be given intravenously to get an immediate effect and avoid delayed absorption due to peripheral stagnation. Pain is not a prominent feature of most extensive deep burns, and sedatives which themselves increase tissue anoxia are often not required.

Oxygen.—Cyanosis is common with extensive burns and may be the result of haemoconcentration, peripheral stagnation of the circulation, or oedema of the lungs. Usually the administration of oxygen is valueless, but sometimes the cyanosis is immediately replaced by a healthy pink colour, and in these cases it is presumably useful.

Heat.—The most comfortable room temperature for treatment is 70 to 75° F. (21.1 to 23.9° C.), and in this environment the patient will usually feel warm when the shock is controlled.

Control of Infection

Since only 10% of burns examined within a few hours of injury are contaminated with pathogenic organisms, recent burns should be regarded as clean wounds. Owing to their large surface area, however, they are very likely to become infected by direct contact or from the air, and special precautions are required to prevent this. Before an aseptic dressing technique was used and antibiotics became available sepsis was a common cause of death and accounted for much that was attributed to toxæmia.

There are two principal methods for treating burns at the present time—the closed method and the exposure method. The closed method is based on the knowledge that burns are originally clean wounds, and that with adequate measures they can be kept clinically uninfected. These measures consist of: (1) immediate first-aid cover for the burn, with a clean cloth only; (2) the early application of a prophylactic antibiotic cream covered by a dressing to act as an antibacterial barrier; and (3) the employment of a no-touch aseptic technique for dressing the burn, changing of dressings as infrequently as possible, and ideally an air-conditioned dressing-room to prevent air-borne contamination.

An effective dressing is obtained by spreading penicillin cream (1,000 units per gramme in "lanette wax S.X.") liber-

ally over the burn and surrounding skin with a sterile spatula, covering this with gauze to facilitate removal of the dressing, and then applying dry cotton-wool and a crêpe bandage. It is necessary to stress that dressings should contain a thick layer of dry cotton-wool which will act as an efficient barrier against bacteria; as soon as the dressings become soaked with exudate they can be penetrated by organisms from the outside of the dressings within a few hours. A thin plaster cast applied over this dressing is invaluable in children, for elevation of limbs, for immobilization, and to prevent removal of the dressing. A stronger penicillin cream (10,000 units per gramme) has proved effective against haemolytic streptococci, and "polymyxin" (0.1 mg. per gramme) in the same base has been shown to be of prophylactic and therapeutic value against *Pseudomonas pyocyanea* and coliform bacilli. Other antibiotics are under investigation, but their value is not yet proved.

The exposure method is based on an entirely different principle. Since it is difficult to prevent bacteria colonizing the surface of a burn, efforts are directed towards drying the necrotic dermis to a hard crust as soon as possible so as to make conditions unfavourable for bacterial survival. Accordingly, the burn is exposed and immobilized, and precautions are taken to prevent cracking of the scab. Penicillin in lactose (10,000 units per gramme) may be dusted on to the burn four-hourly during drying, and the treatment is usually combined with systemic penicillin until the burn is dry.

With both methods bacterial colonization on the burned surface occurs, but clinical infection is uncommon. This gives little cause for satisfaction, since it has been shown that even colonization of burns with *Str. pyogenes*, *Ps. pyocyanea*, and coliform bacilli delays healing and hinders successful grafting. It seems likely that bacterial colonization will ultimately be preventable by a combination of local chemotherapy and a bacteria-proof cover. At present the two methods of treatment are entirely complementary, the choice depending upon such factors as the age of the patient, the site and size of the burn, the climate, and the possibility of treating the injured person as an out-patient. In peacetime conditions the method which is easiest to carry out efficiently is the method of choice for the particular patient. No surgeon experienced in the treatment of burns would deny that the exposure method is best for burns of the face or buttocks which cannot be excised and grafted primarily. On the other hand, it is easier to give active exercises and postural change to elderly patients with extensive circumferential burns of the trunk by the closed method of treatment.

[Part II will be published in our next issue.]

One day in June—the date has yet to be arranged—another of the London County Council's blue and white plaques is to be erected at the Lyric Theatre, and will be unveiled at a short ceremony. The inscription on the plaque will read: "L.C.C. This was the home and museum of Dr. William Hunter, Anatomist, 1718–83." William Hunter crossed the border to come to London in 1740, and, after becoming a member of the College of Surgeons, reached the top of his profession by middle age, being appointed physician-extraordinary to Queen Charlotte in 1764. He was recognized as one of the leading anatomists and obstetricians of the day, and in 1770 he took up residence at Great Windmill Street, where over several years he built a house, lecture theatre, dissecting-room, and museum. The house has been incorporated into the Lyric Theatre which fronts on to Shaftesbury Avenue, but the old frontage in Great Windmill Street can still be discerned. It is said that his dissecting-room was actually where the theatre stage is now situated. To his anatomical and pathological collections Hunter added paintings, coins, medals, shells, corals, and rare books. There will be an exhibition of part of the Hunterian Collection in London from June to August at the Iveagh Bequest, Kenwood, on Hampstead Heath.