A HISTOLOGICAL STUDY OF ONCHOCERCIASIS TREATED WITH HETRAZAN

BY

FRANK HAWKING, D.M., M.R.C.P., D.T.M.

(From the National Institute for Medical Research, Mill Hill, London, and the Medical Department, Uganda)

[WITH SPECIAL PLATE]

During 1949 about 50 patients with onchocerciasis were treated with hetrazan at Jinja, in Uganda, East Africa. The clinical details of this work have been described elsewhere (Hawking, 1950). The present paper describes a histological study of nodules and skin biopsies taken before, during, and after treatment with this compound. Hetrazan ("banocide") is 1-diethylcarbamyl-4-methyl-piperazine. It was discovered by Hewitt *et al.* (1947). The dose used in the present work was 50 mg. of dihydrogen citrate salt (which contains 51% base) twice or three times daily, the dose being increased after the first two days until 250 mg. thrice daily was given. In normal persons the compound caused practically no toxic symptoms apart from slight nausea or anorexia; but in persons with onchocerciasis it caused violent allergic reactions (swelling, oedema, and tenderness of the skin, itching, a fine papular rash, enlargement and tenderness of adjacent lymph nodes, together with fever, headache, and other systemic symptoms) which lasted three to four days, and which were proportional to the density of the microfilariae in the skin rather than to the dosage of hetrazan. The microfilariae usually disappeared from the skin after several days, but they often reappeared after treatment ceased. The adult worms, as determined by fresh examination of excised nodules, did not seem to have been much affected by the hetrazan.

Methods

In order clinically to determine the number of microfilariae in the skin, frequent skin slips were made from a part adjacent to nodules—that is, usually near the anterior superior iliac spine or the great trochanter. These were taken by raising a fold of skin between finger and thumb and slicing off a small segment with a razor blade. The skin shaving (about 0.3 by 0.5 cm.) was placed in Ringer's solution under a cover slip and left for about half an hour to see if microfilariae would emerge. In one heavily infected patient 174 microfilariae emerged from a piece of skin measuring approximately 16 mm.² At the end of the microscopical examination the piece of skin was pressed (cut surface downward) upon a small piece of card, to which it readily adhered, and it was then fixed in formol-saline. From the large number of specimens thus collected suitable pieces have been selected for histological examination; serial sections were cut parallel with the skin surface and stained with haematoxylin and eosin.

In addition, nodules were excised from patients before and after hetrazan treatment. The results of examination in the fresh condition have been described in the previous paper (Hawking, 1950). The nodules were fixed in formol-saline and prepared for histological examination in the usual way. There was also a small amount of post-mortem material, fixed in formol-saline.

Description of Nodules

Specimens were available from five untreated patients and from nine patients after treatment with hetrazan, as follows:

Case 20, after 0.38 g. in 8 days	Case 37, after 5.8 g. in 15 days
" 23 " 4.26 " " 26 "	" 38 " 0.2 " " 24 "
" 30 " 2.30 " " 13 "	" 38 " 2.50 " " 26 "
" 32 " 2.9 " " 13 "	" 50 " 4.17 " " 15 "
" 33 " 4.75 " " 27 "	

Doses given in terms of base.

Each specimen consisted of one to six nodules bound together with fibrous tissue. Usually about three histological sections were prepared from each specimen and stained with haematoxylin and eosin. The appearances of worms and of the adjacent part of the nodules varied greatly. For convenience the worms may be classified into the four groups defined by Ashburn, Burch, and Brady (1949), which are briefly as follows: I, microfilariae normal; II, microfilariae necrotic; III, microfilarial production suppressed; IV, dead worms and their disintegrating remains. In some specimens, one histological section showed normal worms (Group I) while another showed only dead worms (Group IV); in such cases each section has been counted separately. Accordingly the normal group of nodules yielded seven sections, of which three were Group I, two were Group III, and two were Group IV; microfilariae were present in the wall of the nodule in four cases and absent in three. As regards the cellular reaction round the worms, it was acute (polymorphs, etc.) in two cases, chronic (monocytes and plasma cells) in one, and absent in two.

From the patients treated with hetrazan there were 11 sections, of which six were Group I, three were Group II, and two were Group IV; microfilariae were present in the wall of the nodule in two cases, rare in two cases, and absent in seven. The cellular reaction round the worms was acute in five cases, chronic in four, and absent in two. There is no significant difference between these two series, and no histological evidence that hetrazan had produced any adverse effect upon the adult worms, except that possibly an acute inflammatory reaction round the worms is more common after hetrazan treatment. It is possible that some effect might have been seen 60 days after treatment (as suggested by Ashburn *et al.*), but the significance of this (if it occurred) would be obscure. As indicated above, the cellular reaction round the worms could be classified into three groups: (a) no reaction, usually round healthy worms of Group I; (b) a chronic cellular reaction consisting chiefly of monocytes, plasma cells, and fibroblasts; round dead and dying worms there were often giant cells, some of which were large cells applied immediately to the worms and others were smaller ones scattered through the capsule; and (c) acute inflammation consisting chiefly of polymorphs and eosinophils. It may be concluded that there was no significant difference between the nodules from untreated patients and those from treated patients.

Description of Skin Slips

Untreated Patients.—The skin slips from the patients were taken from the neighbourhood of a nodule (usually near the anterior superior iliac spine), and in untreated persons the skin had macroscopically appeared normal. When the specimens from the untreated persons were examined microscopically various numbers of cells were found. In most of the specimens the number was probably within normal limits, allowing for minor reactions due to scratching, trauma, or skin irritations. The cells were often gathered round the small vessels, but some were also scattered diffusely through the connective tissue of the corium. The cells were mostly large pale angular ones with pale elongated nuclei (fibrocytes), oval nuclei (fibroblasts), or round nuclei (monocytes). A few plasma cells were often present. In some specimens there were a few lymphocytes and eosinophils, but these were uncommon. Most of the specimens contained many microfilariae, even though there had been an opportunity for these to migrate out while the specimen was immersed in Ringer's solution (see above). The micro-

F. HAWKING: ONCHOCERCIASIS TREATED WITH HETRAZAN

FIG. 1

FIG. 2

FIG. 3

FIG. 4

FIG. 5

PETER STORY AND W. E. GIBB: SARCOMA OF SPLEEN

FIG. 1.—Lung metastasis: many blood spaces, pleomorphic cells, and nuclei; occasional multinucleate and few mitotic cells. (H. and E. $\times 200$.)

FIG. 2.—Lung metastasis: predominance of long spindle-shaped cells. (H. and E. $\times 500$.)

FIG. 3.—Liver metastasis: compact layer of tumour cells round blood vessel. (H. and E. $\times 120$.)

FIG. 4.—Liver metastasis: necrotic zone with thin layer of tumour cells; islands of living cells round blood vessels. (H. and E. $\times 50$.)

filariae lay free in the subepithelial connective tissue of the corium and there was no cellular reaction to their presence (Special Plate, Fig. 1). No microfilaria was seen penetrating the epidermis. In a piece of skin from a post-mortem examination two microfilariae were seen lying between the collagen bundles of the reticular layer. In the material from one post-mortem examination no microfilariae were found in sections made from the lung, liver, kidney, adrenals, and pancreas; in specimens from the conjunctivae rare microfilariae were found, lying in the subepithelial connective tissue and exciting no cellular reaction.

Histology of Skin after Hetrazan

In specimens of skin taken 18–24 hours after the first dose of hetrazan there were numerous foci of inflammation (Figs. 2 and 3). This inflammation varied in the different patients, ranging from a very intense reaction down to a mild one; the severity of the histological inflammation was proportional to the severity of the clinical reaction, which was roughly proportional to the number of microfilariae initially present. In the severe cases the inflammatory foci consisted mainly of neutrophil and eosinophil polymorphonuclear leucocytes; monocytes and fibroblasts were common; lymphocytes and plasma cells were few. Fibrinous exudate was often present in the centre of the focus, and in very severe cases the nuclei of the cells near the centre were pyknotic and degenerate. The inflammation sometimes involved the stratum germinativum of the epidermis, so that the basal cells were separated by oedema, the nuclei were shrunken, and groups of inflammatory cells were present in the epidermis itself. Microfilariae were often but not always present in these inflammatory foci, phagocytic cells being clustered near them in many cases and the larvae showing signs of degeneration—for example, derangement, separation, or scattering of the nuclei. But in some cases the microfilariae appeared to be degenerate without the proximity of phagocytes; while in other places microfilariae might lie freely in the connective tissue without any marked inflammation around them. Presumably the inflammatory reaction is an allergic one due to the liberation of microfilarial products by hetrazan in a skin which is sensitive to them; but histologically this association is often absent. Clinically the inflammation of the skin was manifested by swelling of the skin, with papular rashes, itching, and systemic febrile reactions.

During the next five days the inflammation of the skin gradually diminished and its character changed (Fig. 4). In a typical late case after five days the inflammatory foci were circumscribed, the intervening connective tissue being normal; the cells were mostly monocytes, fibroblasts, and early

fibrocytes, together with lymphocytes, plasma cells, a few eosinophils, and rare neutrophils. No giant cells were seen. The extent of the inflammation varied in the different individuals. Some microfilariae lay in the foci, and the nuclei were deranged, indicating degeneration (Fig. 5); others were seen in the intervening connective tissue, without the tissue making any apparent reaction to their presence. Microfilariae were sometimes found in sections of skin, although none had been detected in life by immersing the skin slip in saline and waiting for larvae to migrate out.

During this work a patient was seen with lesions of the skin but no detectable microfilariae. The skin of the left thigh was greatly thickened, dry, and hard, and it contained many small flat lumps about 2 cm. across. A similar but less extreme condition was shown by the skin of both buttocks, the left leg, the right lower extremity, the feet, and the abdomen. In spite of careful search no microfilariae or nodules were discovered. Histologically, the corium of the skin contained circumscribed blocks of cells, most of which had large oval nuclei (? fibroblasts), but many eosinophils were also present, together with a few neutrophils and lymphocytes; no microfilariae were seen.

The man was put on hetrazan, and within 18 hours of the first dose he developed a typical allergic reaction, with a dense papular rash covering most of the trunk and lower limbs. Histologically, the corium contained numerous diffuse areas of intense inflammation. The centre of these areas was occupied by fibrinous exudate and great numbers of necrotic neutrophil polymorphs; towards the periphery there were eosinophils, neutrophils, and fibroblasts, together with a varying number of lymphocytes and monocytes. The intervening connective tissue of the corium was swollen and oedematous. The epithelial cells of the epidermis were separated by fluid. This reaction gradually settled down in about nine days, during which hetrazan was continued. In view of the intense reaction to hetrazan it seems probable that the man contained onchocerca microfilariae, to which he was sensitized, and their destruction by the drug fired off the allergic reaction; but in that case the reaction must be directed against the diffused products of degenerating microfilariae rather than against the presence of the larvae themselves, since the microfilariae were too few to be detected by histological examination.

Discussion

A bibliography concerning the treatment of onchocerciasis with hetrazan has been given by Hawking (1950). A histological study on the nodules and adult worms during such treatment was made by Ashburn, Burch, and Brady (1949). They found that nodules removed in less than 60 days after treatment showed no effect; in nodules removed at a later date there was possibly some suppression of the reproductive powers of the female worms, but this suppression was not certain. In the series described above no significant effect upon the adult worms in the nodules could be detected when a single course of hetrazan, lasting 15–27 days, had been given. It is possible that repeated courses might have been more effective, although they would not be practicable under the usual conditions of medical work in Africa.

A brief report on the histological changes in the skin has been given by Baez (1949), whose description is similar to the one given here but less complete. As was shown above, in untreated patients microfilariae often exist in the skin without provoking any histological reaction; if a reaction has been caused it is mild in degree and chronic in nature (fibroblasts, monocytes, etc.). When a small dose of hetrazan is given there is an intense acute reaction in the corium of the skin with great numbers of neutrophil and eosinophil polymorphonuclear leucocytes. This reaction is presumably due to the destruction of microfilariae, with the liberation of foreign protein which excites an allergic response in the sensitized patient. Often the foci of acute inflammation contain microfilariae, a circumstance which may or may not show derangement of the nuclei and other visible signs of

F. HAWKING: LEGENDS TO PLATE

FIG. 1.—Skin from untreated patient (*post mortem*). No reaction round microfilaria. (H. and E. $\times 600$.)

FIG. 2.—Eighteen hours after hetrazan, which produced an intense reaction. Microfilariae are seen in the corium surrounded by polymorphs, etc. Five living microfilariae emerged from this specimen. (H. and E. $\times 900$.)

FIG. 3.—Twenty-three hours after hetrazan, which produced a flat papular rash. Inflammation has produced small cyst in basal layers of epidermis, filled with polymorphs and eosinophils and centred round a microfilaria. Eighteen living microfilariae emerged. (H. and E. $\times 375$.)

FIG. 4.—After hetrazan for two days; moderate reaction subsiding. Microfilaria in deepest layer of epidermis surrounded by polymorphs. In corium (lower right) is a group of tiny nuclei from degenerating microfilaria. No living microfilariae emerged. (H. and E. $\times 714$.)

FIG. 5.—After hetrazan for five days. Groups of small nuclei from disintegrating microfilariae, with monocytes and fibroblasts. One living microfilaria emerged from this specimen. (H. and E. $\times 1,250$.)

degeneration. On the other hand, the distribution of inflammation often fails to coincide with that of microfilariae, for the inflammatory reaction may be widespread in cases where microfilariae are scanty or impossible to find, or, conversely, microfilariae can be seen histologically in connective tissue free from leucocytes; and living apparently normal microfilariae may emerge from samples of skin when examined in the fresh condition. The skin slips were taken within a few centimetres of nodules containing adult worms, and some of these microfilariae might have recently migrated out of the nodules.

The action of hetrazan on onchocerca may be compared with its action on the experimental infection of *Litomosoides* in cotton-rats and on *Wuchereria bancrofti* in man. In all cases it acts most promptly on the microfilariae, while its effect on the adult worms is much less marked. In previous papers (Hawking, Sewell, and Thurston, 1950; Hawking, 1950) the hypothesis has been put forward that the action of hetrazan upon microfilariae is due to an opsonin-like effect, so that the microfilariae become attractive to the macrophages of the reticulo-endothelium system, which accordingly fix them and then destroy them. The present observations would agree with this hypothesis if one makes the additional assumption that hetrazan acts more readily on some microfilariae than on others.

Thus the first effect of hetrazan would be to "opsonize" some of the microfilariae, so that the previously indifferent phagocytes of the skin become attracted to them and begin to destroy them. The first liberation of microfilarial products would excite the acute allergic inflammation described above, which would be more widespread than the microfilariae themselves. Meanwhile some microfilariae are postulated as being less readily affected by hetrazan (because they are newly emerged from adult worms, or for some other unknown reason), and these are the microfilariae which migrate from the skin slip in the fresh condition or which are found free from inflammatory reaction in the stained section. If the treatment with hetrazan is prolonged all microfilariae in the skin are eventually suppressed by it, for the skin slips no longer liberate microfilariae in the fresh state. But the adult female worms still remain alive, and when the administration of hetrazan is discontinued they gradually replenish the skin with fresh microfilariae. Unless means of destroying the adult worms can be found, treatment of onchocerciasis with hetrazan will give only temporary relief.

The allergic reaction provoked by hetrazan in patients with onchocerciasis may be used as an additional means of diagnosis. If the skin slips of a patient suspected of this infection reveal no microfilariae and he has no allergic reaction after taking hetrazan, it is unlikely that he contains onchocerca.

Summary

Patients infected with *Onchocerca volvulus* were treated with hetrazan, and subcutaneous nodules containing adult worms and samples of skin containing microfilariae were removed for histological examination.

There was no significant difference between nodules from five untreated patients and those from nine patients treated with hetrazan, and there was no evidence that hetrazan had produced any adverse effect upon the adult worms.

In skin samples from untreated patients, microfilariae in the corium usually did not excite any reaction; if reaction was caused it was mild in degree and chronic in nature.

In skin samples after treatment with hetrazan there were numerous foci of intense acute inflammation. Microfilariae were often present in these foci, and showed evidence of degeneration and destruction; but sometimes they were also present in normal connective tissue without exciting a reaction.

It is probable that the main action of hetrazan is to destroy microfilariae (probably by an opsonin-like effect), and that their destruction liberates some microfilarial product which excites an allergic reaction in the previously sensitized patient. Some microfilariae, however, are apparently less susceptible to the opsonin-like effect of hetrazan, and so they persist in the skin for long periods without doing visible harm.

The allergic reaction provoked by hetrazan in patients with onchocerciasis can be used as an additional means of diagnosis.

Grateful acknowledgments are due to the Director of Medical Services in Uganda and his colleagues for clinical facilities; to Mr. F. Higginson for histological preparations; and to Mr. M. R. Young, A.R.M.S., for the photography.

REFERENCES

- Ashburn, L. L., Burch, T. A., and Brady, F. J. (1949). *Bol. Ofic. sanit. panamer.*, 28, 1107.
Bacz, M. M. (1949). *Rev. Inst. Salub. Enferm. trop., México*, 10, 95.
Hawking, F. (1950). *Trans. roy. Soc. trop. Med. Hyg.*, 44, 153.
Sewell, P., and Thurston, J. P. (1950). *Brit. J. Pharmacol.*, 5, 217.
Hewitt, R. I., Kushner, S., Stewart, H. W., White, E., Wallace, W. S., and SubbaRow, Y. (1947). *J. Lab. clin. Med.*, 32, 1314.

FURTHER OBSERVATIONS ON CAPILLARY RESISTANCE AND ADRENOCORTICAL ACTIVITY

BY

H. N. ROBSON, M.B., F.R.C.P.Ed.

AND

J. J. R. DUTHIE, M.B., F.R.C.P.Ed.

(From the Departments of Medicine, Universities of
Aberdeen and Edinburgh, and the Rheumatic Unit, Northern
General Hospital, Edinburgh)

Earlier observations on changes in capillary resistance as measured by a negative-pressure method have already been reported (Robson and Duthie, 1950). It had been shown by Scarborough (1944) that the resistance of the capillaries in the skin rose sharply after a variety of surgical operations. This was confirmed by Robson (1949). As described previously, it was then shown that similar changes in resistance were caused by other types of non-specific stress such as x-ray irradiation, intravenous nitrogen mustard, hyperpyrexia, and the intravenous injection of T.A.B. vaccine. In view of the variety of "alarming" stimuli which gave rise to similar changes, it was felt that the common factor might be the activation of the pituitary-adrenal axis. This idea was supported by the observation that insulin-induced hypoglycaemia and the injection of adrenaline both caused a rise in resistance, since both these procedures stimulate the adrenal cortex, as evidenced by a fall in the number of circulating eosinophils.

The hypothesis was then tested more directly by studying the effects of adrenocorticotrophic hormone (A.C.T.H.). It was found that a single dose of 25 mg. of A.C.T.H. produced a significant rise in resistance in about four hours. Repeated injections, usually of the order of 25 mg. of A.C.T.H. eight-hourly, resulted in a marked elevation of capillary resistance, which was sustained during the period of administration. Such observations were made on patients suffering from a variety of conditions, including rheumatoid arthritis, ankylosing spondylitis, disseminated lupus erythematosus, and idiopathic