REFRESHER COURSE FOR GENERAL PRACTITIONERS

FRACTURES IN THE AGED

BY

RICHARD H. METCALFE, M.D., M.Chir., F.R.C.S.

Consultant Orthopaedic Surgeon, St. James' Hospital, London, S.W.12

The term "aged" is relative, depending on the physical and mental state of the patient, but in this article it will be taken to mean those over the age of 70. With the gradual increase in the expectation of life the problem of fractures in the aged has become more and more acute. Although from the surgical aspect the problem is partly solved and methods of treatment in the acute stages and during the period of rehabilitation are to a great extent standardized, the end-results of surgery still leave much to be desired. The major problems are sociological and psychological. The most urgent problem to-day is the disposal of these patients after their discharge from hospital.

Fractures in the aged occur with extreme ease, not only on account of the general osteoporosis and fragility of their bones, but also occasionally as a result of secondary malignant disease, often arising from some completely unsuspected or undiagnosed primary source. These fractures are mostly limited to certain regions of the body, undoubtedly the commonest type requiring admission to hospital being those at the upper end of the femur. Other common sites are the shaft of the femur, the wrist, the upper end of the humerus, and the spinal column. Fractures involving the knee, ankle, and elbow are relatively rare in the aged.

Fractures at the Upper End of the Femur

Fractures of the upper end of the femur are of extreme gravity in the aged. In women the fracture almost invariably occurs at the femoral neck and more commonly on the left side: in right-handed persons the right leg is apparently stronger than the left. With men fractures of the neck are rare, and the intertrochanteric fracture with complete avulsion of the lesser trochanter is most frequently encountered. Both these fractures can occur as a result of only slight violence. Sometimes there is impaction of the neck in old women and the actual fracture remains undiagnosed, but in most cases the shaft of the femur is pulled up by the gluteal muscles and deformity occurs at the site of the fracture. The classical signs are that the patient lies with the injured leg visibly shortened and with the foot completely externally rotated, owing to the fact that the centre of gravity of the femur is well outside the line passing through the centre of the hip and the knee-joints.

At this stage the patient suffers from severe shock and pain, and much relief can be obtained immediately if the rotation deformity is corrected so far as is possible. This can be done either by tying the feet together or by screwing a piece of wood about a foot (30 cm.) long to the heel of the shoe at right angles to the long axis of the foot and projecting laterally; these must be regarded as temporary measures only, while awaiting transfer to hospital. All these patients should be admitted to hospital as early as possible for radiography and the institution of treatment. Once this fracture is suspected, and while awaiting the patient's admission to hospital, the leg should be fixed by extension on to a Thomas splint. The amazing results of this form of fixation in the front line during the first world war are well known. During the second world war Charnley introduced the adjustable Thomas splint, which permits variation of both length and size of ring. Many thousands of these splints must be in store, and much unnecessary pain could be avoided if some were issued to general practitioners for these emergencies.

Hospital Regime

The admission of old people suffering from fractures at the upper end of the femur is fraught with difficulty. Their stay in hospital is necessarily prolonged, and treatment is best carried out in special wards set aside for this type of case. This is necessary because the treatment is highly specialized and both surgical and nursing staff must be enthusiasts. The wards should not be limited entirely to the aged, as a leavening of younger people and occasionally even older children gives added interest and pleasure to elderly patients.

The essence of treatment of these old people lies in the preservation of their morale, and the sight of patients recovering is most encouraging and helpful to new arrivals. A new case should therefore be placed in a bed next to a patient suffering from a similar injury who is well on the way to recovery. The healing of fractures depends on a full and adequate diet and a good vascular supply to the site of the fracture. The latter can be assured only if full mobility is encouraged as soon as possible after the fracture has been fixed. The best way to achieve this is by a daily physiotherapy class carried out in the ward (Fig. 1). The

FIG. 1.—Daily physiotherapy class.

physiotherapist should place herself in a raised position where all the patients can see her. Exercises are not limited to the fractured limbs but must mobilize all the limbs and the abdominal, chest, and back muscles. The patients enjoy these exercises immensely, and when the physiotherapist is on leave or for some reason unobtainable one of the younger patients is usually only too willing to take her place and supervise the exercises. In my ward all patients do these daily exercises, the only ones being excused being those aged over 90, and generally even these try to join in. Frequent baths with salt and soda in the water go a long way to loosen stiff joints once the ambulant state is reached.

The third essential is proper attention to meals. Many old people live alone in one room and feed on bread-and-margarine with an occasional cup of tea and but little else. Is it surprising, then, that their bones are so fragile and break so easily? If the fractures are ever to unite it is essential that these patients be provided with good food, attractively served, and be encouraged to eat all that is set before them. This can be done only with the help of an enthusiastic sister and nursing staff.

Surgical Treatment

Correction of the deformity of the fracture and fixation at the earliest possible date by pins or plates are the surest ways of relieving the patient of all pain and constitute the first stage to the restoration to full mobility. It must be

fully realized that, no matter how much steel-work in the form of pins, screws, nails, and plates is inserted into or on to a fractured bone, such treatment does not reduce the time of healing of a fracture by one minute. The reverse may unfortunately even be the case, for, if the steel-work results in holding the bone apart rather than well-impacted together, then union will be delayed or may never occur. When treatment by internal fixation was first used it was thought that the period of rest in bed could be considerably reduced. This unfortunately has not proved true, especially in fractures of the neck of the femur.

The blood supply to the head of the femur in old people is restricted almost entirely to that passing up the neck of the femur from the shaft; supply through the ligamentum teres is non-existent and that from the capsule negligible. Unless adequate time is allowed for the restoration of the blood supply across the fracture line avascular necrosis of the head of the femur will occur, and the pin will cut out, with resultant complications. Therefore a minimum period of 12 weeks' rest in bed is required for all fractures of either the upper end or the shaft of the femur in old people. Attempts are now being made to reduce this period of rest with the object of saving beds. Two methods are in use—the Künchner pin for fractures of the middle third of the shaft of the femur, and the replacement of the femoral head by a Judet's plastic prosthesis in sub-capital fractures of the neck of the femur. The first of these methods is firmly established, but the second is still in the experimental stage, and it is still too soon to say whether it is preferable to the usual fixation with a Smith-Petersen pin.

At the end of 12 weeks, if the appearance of the head of the femur is satisfactory, the patient gets up, wearing a bucket-top calliper, which to some extent takes the weight off the fracture line. This is worn until radiographs show full consolidation of the fracture, which does not usually occur until about six to nine months after the fracture.

Other Fractures in the Aged

Pathological Fractures.—Within the last year I have had under my care three pathological fractures of the upper end of the femur secondary to carcinoma of the breast. Internal fixation was impossible, and these patients have been treated by extension on a Thomas splint and the administration of dienoestrol, 5 mg. three times daily. The results have been remarkable: there have been bony union and almost complete disappearance of the secondary lesion, and the patients have in due course been able to walk. No untoward effects following this treatment have been observed. Dienoestrol is a synthetic oestrogen similar to stilboestrol but stronger, and consequently the dose required is smaller.

Fractures of the Spine.—The commonest fracture of the spine is a straight "concertina" compression of the bodies of one or more vertebrae, the result of marked generalized osteoporosis and minor trauma. Fixation in a plaster jacket is unnecessary and treatment by rest in bed with a firm pillow under the back is remarkably successful. Good consolidation occurs rapidly, especially if the patient is given a good diet and extension exercises for the erector spinae in the later stages of treatment. It is unnecessary to provide any supporting apparatus subsequently; this is not well tolerated.

Fractures of the Upper End of the Humerus.—The usual fracture is the transverse one just below the surgical neck. In many cases there is impaction of the shaft into the softened and osteoporotic head. These cases require no treatment other than rest in a sling, and should never be admitted to hospital. In other cases there is overlap of the upper end of the shaft upward in front of the head. This can be reduced by closed manipulation with great ease, as muscle spasm of the pectoralis major and deltoid is negligible in old people. A roller towel is placed round the chest to immobilize the trunk. The arm is then

abducted to 90 degrees under general anaesthesia and strong extension applied. It is then easy to push the shaft back under the head and reduction of the fracture can be achieved. Patients can then be treated as out-patients if the home conditions are suitable. It is by no means essential to obtain complete reduction, provided that the overlap and shortening are corrected.

Fractures of the Wrist.—The Colles fracture in an old person is a daily problem. Treatment should be carried out in the usual way by reduction under anaesthesia and fixation in plaster from below the elbow to the palm of the hand. Whereas in young people gross swelling occurs around the wrist and fixation during the first week should be by means of a posterior plaster slab only, in old people very little swelling occurs, and complete fixation in plaster at the time of reduction is possible. The blood supply to the lower end of the radius in old people is notoriously poor, and it is recommended that fixation in these cases should not be for less than 10 weeks. If one is tempted to remove the plaster earlier, displacement of the fracture only too often occurs with its associated disfigurement and poor function.

Anaesthesia

There is still considerable dispute on what is the best type of anaesthetic for open operations in old people. The main requirement is to prevent shock, and in my view the best anaesthetic is intravenous thiopentone followed by gas and oxygen, with a little "trilene" added from time to time as necessary. Local analgesics have been tried and found most unsatisfactory.

After Discharge from Hospital

The correct disposal of aged patients after discharge from hospital is most difficult. Many of them live alone; others live with relatives who are out at work all day. It is regrettable that the relatives of these unfortunate people often "write them off" and do everything possible to avoid having them back home once they have been admitted to hospital. When this happens the family doctor can give great assistance, as he will know the home conditions and any relatives who may be able to help. If it is found impossible to get the patient home, then unfortunately he must take his place among the chronic sick. In the past patients in chronic sick wards spent a large part of their time just lying in bed, but now treatment is much better, and in many cases physiotherapy departments have been opened. This has led to remarkable improvement in the patients' morale and mobility, and in many cases the relatives have been persuaded to take the patients back home again. Unfortunately the number of geriatric beds is quite inadequate and the "waiting-list" is enormous. Aged persons whose fractures have united but who still have a limited degree of mobility, and who have no suitable home to which they can return, can be well looked after in special homes for the aged, such as the Richard Cusden Homes in London. The homes are managed by matrons, and as much domestic work as possible is done by the residents; no nurses are needed, as the residents are not sick.

Back at Home

When the patient is in hospital it is comparatively easy to maintain his mobility. The beds are fitted with vertical bars and chains with handles on the end (Fig. 2). With these the patients can pull themselves up, and there is always help available when they wish to go to the bathroom or lavatory. At home one of the patient's main difficulties is getting up out of bed, out of a chair, or off the lavatory seat. For some years I have advised the provision of strong hooks, bolted into a ceiling joist and fitted with a strong rope and wooden handle, over the positions mentioned. It is remarkable how much this simple device can help.

Old people are often proud and independent and will refuse to go into a hospital for the chronic sick or a home for the aged. They prefer to return to their one room or

FIG 2.—Bed fitted with chain and handle.

to the home of relatives who are out all day. It is only after they have achieved their desire that they realize the terrible pangs of loneliness: patients attending hospital for follow-up constantly state their wish to be back in the ward with its bustle and happy companionship. Much is already achieved in some districts with the willing help of the Darby and Joan and similar clubs. If such a club is told that some old person is immobilized at home and alone for long

periods of the day arrangements are nearly always made for more active members of the club to pay visits. It may be possible to arrange through the medical officer of health of the local health authority for a "home help" as a temporary measure. Neighbours will often help with shopping, or there may be a "home meal service" in the district. A wireless set may also be a comfort, and there are societies which help towards the cost of providing these.

The services of occupational therapists are most useful in the treatment of fractures in old people. Patients vary considerably in their reaction to this form of treatment. Some are only too keen to carry out work, and it is often found they are more willing to do so if the work is for their own hospital. Other old people, who may never have had a proper rest in their lives before, are only too glad to lie quietly, and quite frankly cannot be bothered. Occupational therapy should not be forced on old people, but at the same time no patient in a fracture ward for the aged must be allowed to lie in bed like a log of wood.

Conclusions

In the treatment of fractures in the aged attention should be directed particularly to morale, mobility, and good meals.

Aged people suffering from fractures and requiring hospital treatment should be gathered together in special wards set aside for the treatment of such cases.

Every effort should be made to encourage the relatives and friends of old people with fractures to take them back into their own homes.

Patients who are unable to return home and who have perforce to be admitted into chronic sick hospitals should be given every opportunity for increasing their mobility by the aid of physiotherapy and occupational therapy.

The avoidance of extreme loneliness after the bustle and companionship of a ward needs urgent consideration.

Next Refresher Course Article.—"Serum Reactions and Serum Sensitivity Tests," by Drs. L. J. M. Laurent and H. J. Parish.

ROYAL COMMISSION ON MARRIAGE AND DIVORCE

ORAL EVIDENCE ON BEHALF OF B.M.A.

Representatives of the British Medical Association attended the meeting of the Royal Commission on Marriage and Divorce at Caxton Hall, Westminster, on May 28 to support orally the Association's memorandum of evidence, already presented. The memorandum was published as Appendix IV to the Annual Report of Council in the *Supplement* of April 19 (p. 188). The deputation was led by Dr. H. Guy Dain, and its members were Dr. Annis Gillie, Dr. J. A. Hobson, Dr. Doris Odium, and Dr. A. Walk, with Dr. E. E. Claxton (secretary).

The Chairman of the Royal Commission (Lord Morton*) said he wanted to ask a question about two proposals in the memorandum: (1) that it should be possible for a petition for divorce to be presented by either party or by agreement by both parties jointly, and (2) that in certain circumstances it should be possible after a time for the offending party himself or herself to sue for divorce. These proposals had caused considerable stir both inside and outside the British Medical Association, and he wanted to find out how far they represented the views of the Association's 65,000 members.

Misapprehensions about the Report

Dr. Dain said he was anxious to make a statement in view of the publicity which the document had received and the misapprehensions which had been aroused. The Council of the Association set up a special committee to prepare evidence, and that committee had endeavoured to answer certain of the problems which confronted the Commission, restricting itself closely to the terms of reference. It had been so successful in this last intention that it—and through it the Association—was now accused of wishing to make divorce more easy. It had not expressed any views on the sanctity of marriage as a permanent life contract, and not much was said about it in the document. But, unfortunately, a summary was presented in front of the main document, and people read the summary without reading the argument. The committee represented not only the British Medical Association but to an extent also the Medical Women's Federation and the Royal Medico-Psychological Association. But it had made no attempt to express the religious or ethical views of 65,000 doctors. It had addressed itself entirely to medical problems arising out of the divorce laws and to changes which they thought would make those laws more effective. It had accepted divorce as part of the structure of present-day society and had attempted to give the Commission advice on which improvements might be brought about. It laid stress on preventing divorce by various means which were suggested and spoke of methods whereby anomalies could be straightened out. It was due to the witnesses that this should be stated at the very beginning in view of the public misapprehension of the Association's attitude.

Lord Morton said the Secretary of the Association had stated that the Council had no objection to the publication of its proposals after the oral evidence had been given, but in fact the proposals were published and reached the Press before the witnesses appeared before them, and before there was any opportunity of elaboration or explanation by oral evidence. The proposals were seized upon by the Press for their news value, and it was widely thought that the Association had gone beyond the medical field and expressed a wider point of view. This had caused considerable perturbation; the Commission had received many letters from doctors who had disclaimed these views, and he had seen correspondence in the *British Medical Journal* to the same effect.

*Lord Morton of Henryton is a Lord of Appeal in Ordinary. He was created baron in 1947.