discovered that he can, then only is he plunged into deep water. Serum anaphylaxis can be fatal on rare occasions, but the experienced clinician will generally save his patient. On the other hand, serum sickness may be a troublesome but never a dangerous disease, and there are no fatalities except from ill-conceived treatment.

Conclusion

Whenever serum is injected, whatever the route, the patient should be kept under observation for at least 30 minutes afterwards, with a syringe and adrenaline 1 : 1,000 ready for emergency use.

Intradermal, conjunctival, and scratch tests for the detection of serum sensitivity are unreliable. The small "trial dose," given subcutaneously, is preferable, and should be used in all patients with a previous history of serum injections or allergic conditions.

Serum should never be given intravenously unless a preliminary intramuscular injection has been tolerated and all preparations have been made beforehand for treating shock.

Next Refresher Course Article.—"Clinical Effects of Exposure to Heat and Sunlight," by Professor B. G. Maegraith.

THE PROBLEM OF THE AGED SICK AS SEEN BY THE GENERAL PRACTITIONER*

BY

MORTIMER O'SULLIVAN, M.B., B.Ch.

From the general practitioner's point of view the aged sick are usually those persons who require more or less constant medical supervision and a considerable amount of personal attention. They are mostly visited in their homes, but those who can, even with an effort, attend during surgery hours are encouraged to do so. For the general practitioner there is no clear dividing line between the chronic sick proper and the aged infirm. This classification is of importance to us only when the question arises of removing them from their homes. Before and after they should become the responsibility of either the hospital or the local health authority the general practitioner has usually to look after them for a considerable time.

Speaking as a middle-aged general practitioner, one is thankful that all the aged do not come under one or the other category. Most of the over-seventies are hale and hearty. But when a doctor has worked in the same district for a number of years his patients grow up with him, with the result that the older he gets the more aged sick he has to look after.

In the last quarter of a century the face of medicine has changed. Osler's remark that "pneumonia is the friend of the aged" is no longer true. Since the advent of the antibiotics the chance of death from natural causes has decreased, and at the same time the improvement in anaesthesia and surgery has greatly reduced the risks of operation.

Types of Aged Sick Encountered

As is to be expected, most of the aged sick are suffering from some degenerative process. In my present list of

chronic cases I find the majority to be those of cerebral arteriosclerosis or of encephalopathy—patients with severe headaches, mental degeneration, irritability, retinal haemorrhages, senile cataract, transient paralysis, and a condition which, to me, seems to be on the increase, vertigo. The next largest group may be classified as coronary arteriosclerosis and hypertensive failure: this class varies from the impaired myocardium as the result of coronary thrombosis to the senile cardiac asthmatic. Next there are the arthritics—the chronic cases of rheumatoid arthritis and of osteoarthritis, both making their victims house-fast. Cases of chronic bronchitis and of pulmonary heart disease breaking down several times a year are always present. I have four cases of hemiplegia, three of disseminated sclerosis, two of paralysis agitans, and four of other forms of paralysis. There are five cases of inoperable carcinoma, four cases of chronic tuberculosis which require visiting, two cases of heart failure from rheumatic disease, one case of pituitary tumour, and one man with senile prostate who refuses operation and has had to be trained to catheterize himself, with all its attendant sequelae and complications. There are two mental cases, both women, who in spite of their delusions (they have been in hospital many times) make a laudable effort to run their homes but are always a worry. There are two cases of hysteria, in both of which one can produce a hemianaesthesia at will. I find that hysteria in the aged is a not uncommon symptom. Finally there are five senile cases, all over 90, with nothing much wrong except that they are physically and mentally enfeebled.

Reasons for the Problem

Perhaps Job was the first recorded case of a chronic sick person. Certainly there have always been aged sick. The problem has become acute (mainly in the urban areas), principally for the following reasons.

1. *People Live Longer.*—If you survive long enough you are almost certain to get on a chronic sick list. About ten years ago I assisted at an operation for the local removal of a carcinomatous breast in a woman of 96. The operation was necessary because the growth had begun to fungate. The patient survived for four years.

2. *Absence of the "Handy Woman."*—A generation ago there were in each of the poorer streets of a district one or two of this type of female. They attended at the time of birth, carrying pails of hot water. They would sit at night with a sick person. The more efficient gave an enema if required, and they finally did the washing and laying out after death. Very many of them were retired "gamps," and, although neither home helps nor nurses, combined the duties of both.

3. *Housing.*—The housing situation to-day reflects directly on the young and indirectly on the old. Gone are the days when the family had means and accommodation to care for aged relatives and servants past the age of service. Few young couples lucky enough to have a house are now able to start married life without having an aged relative thrust on them. As children come along there is no longer enough accommodation for them all; domestic friction arises, and so begins the trail of events which leads to the demand for hospital beds.

4. *Tendency to Avoid Responsibility.*—Because weekly contributions are paid to the State there is a feeling that it is the duty of the State to nurture and nurse, and that the care of the aged is no longer the responsibility of the family.

Home Treatment

I would like here to emphasize—and I think that most general practitioners would agree—that any service dealing with the aged should be so designed as to make it easier

*A paper read to the Middlesex Medical Society at the West Middlesex Hospital, Isleworth, June 14, 1951.

and more practicable to nurse this type of patient in his home rather than to endeavour to increase hospital accommodation. Hospitalization or local authority accommodation solves the problem for the relatives and the general practitioner, but in most cases not for the patient. Old people do not take kindly to removal from their usual surroundings. Many die quickly, and I have known some cases of dementia which resolved themselves on the return of the patients to their homes.

The problem of the aged sick has many aspects: the person may live alone or with another aged person or with relatives, and advice and treatment are influenced by the home conditions and the facilities available. Our problems may be divided into medical and domestic.

Medical Aspects

Occupational Therapy.—Many try to occupy themselves, but many more would benefit from expert training. The bedfast sick patient lives in a world limited by his room. Small events matter a lot, and the completion of a piece of knitting or tapestry is a cause for celebration.

Physiotherapy.—We all know of the long waiting-list at hospital for this branch of treatment, and there is usually a wait in hospital itself before treatment is administered. Is there any possibility of treatment in the home for the odd cases where transport causes distress and pain?

Investigations (x-ray examination, E.C.G., blood test, etc.).—One can become stale on a case because of frequent visits and not notice changes in the patient. Investigation at regular intervals would help the general practitioner to assess the patient's condition more accurately.

Foot Clinics.—Many of the aged infirm become the chronic sick and confined to the house because their feet give out. The old-age pensioner cannot afford the fees of a private chiropodist. The establishment of foot clinics would postpone the immobility of the aged.

Minor Problems such as Constipation and Insomnia.—It is very easy for an aged person to get constipated, and once the rectum is overloaded the feeble muscles are no longer able to expel the bolus. The procedure of manual removal (as many of the cases require) is unpleasant for the operator and the patient. To make sure that there is no recurrence the patient often uses aperients too energetically and so becomes incontinent. Not all the aged respond well to narcotics, and it is alarming to find the patient who has been given a sleeping draught mentally confused for the following few days.

Domestic Aspects

Shopping has to be done and food prepared, otherwise the nourishment is going to consist of bread and tea, with sooner or later the symptoms of malnutrition.

Washing of the Patient.—Nearly all old people living alone become dirty. The bathroom is cold or the preparation of the bath (often a zinc bath in the kitchen) is too much trouble. Similar problems are the washing of clothes and bed-linen, the cleaning of rooms, and attention to fires.

Social Intercourse.—Many of these people spend long hours alone. The value of the doctor's visit is only partly medical. The fifteen minutes spent with an old sick person is worth much more than a bottle of medicine.

Finance.—I feel that in some cases the relatives would be prepared to look after and care for an aged sick person if a contribution were made to the family budget. After all, if the patient has to occupy a hospital bed it costs the country about 15 guineas a week. In suitable cases anyone who takes on the arduous duties of nursing an aged sick relative should be compensated. Is it possible for a relative to be appointed a home-help with overtime pay for week-end work?

Suggested Improvements

1. *Admission to Hospitals or Homes During the Family Holiday.*—I think most children are anxious to look after the aged sick parent, but it is the continuity of care which gets the family down, and surely the young family is entitled to expect a little time to itself once a year. I realize the difficulty from the hospital side—once the patient is in hospital the relatives may not be so anxious to have him home.

2. *Supplies of Bed-clothes and Night Attire for the Incontinent Patient.*—An efficient laundry service would help to solve this problem.

3. *Male Nurses.*—While appreciating the very good work done by district nurses (and in our district we are very fortunate), we all know there are certain cases better looked after by men. One male nurse to every three or four female nurses would suffice, and I know from personal experience they could all work well together.

4. *Sick Visiting.*—To visit the sick is one of the corporal works of mercy. There is a great opportunity here for the churches. If more interest were shown in the aged sick there might be a chance of the younger people taking an interest in church affairs. The problem arises, whose duty it is to inform the appropriate vicar. Perhaps the general practitioner is here at fault for not doing so more often. While on this subject I would like to pay tribute to the Darby and Joan homes in our district. A lonely old man died recently, and I was pleased to find two friends from the home turning up to take care of the body and to look after his affairs.

5. *Increase in Number of District Nurses.*—A seasonal increase would be a great help. In the winter of 1950-1 these nurses had not a chance to deal with the number of calls on them. I would like also to make a plea for the provision of some home help for the week-end work.

6. *Priority Hospital Accommodation for Certain Dying Patients.*—I refer to the type of case that is in a coma, that is incontinent, and whose nursing is most distressing for relatives. This state of affairs may go on for weeks. Alternatively, the provision of sitters-in or nurses for night work during acute or terminal illness is suggested.

Conclusion

Finally, the general practitioner has often to deal with the problem that should not be a problem at all. An elderly sick person is put to bed for some minor illness, but when the illness is over the patient refuses to get up, perhaps because it is cold and there is not enough heating in the house. Very soon there is disharmony between the patient and those looking after him. The relatives first demand his admission to hospital, and in a few days the patient makes the same request. The general practitioner points out that the patient is now well and does not require hospital treatment. He is then accused of having a vested interest in keeping the patient out of hospital. Later there is the threat of a letter to the Minister of Health. The only solution to this problem is to send the patient to the out-patient department of the local hospital, where he will be told in no uncertain terms about his condition. He comes home with restored confidence in his relations, and realizes that, after all, his doctor is not the worst in the world. This problem is then solved for another twelve months.

Such are some of the problems which the average general practitioner has to face in dealing with the aged sick. Whatever the future of medicine in this country, the chronic sick and aged infirm will in the main be the responsibility of general practice. It is hard work, often harassing and occasionally satisfying. The few improvements suggested may help to solve the problem.